

Full wwPDB X-ray Structure Validation Report (i)

Sep 9, 2023 – 07:49 PM EDT

PDB ID : 4H4D

Title: IspH in complex with (E)-4-amino-3-methylbut-2-enyl diphosphate

Authors: Span, I.; Eisenreich, W.; Jauch, J.; Bacher, A.; Groll, M.

Deposited on : 2012-09-17

Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$		
R_{free}	130704	1509 (1.38-1.34)		
Clashscore	141614	1551 (1.38-1.34)		
Ramachandran outliers	138981	1530 (1.38-1.34)		
Sidechain outliers	138945	1530 (1.38-1.34)		
RSRZ outliers	127900	1487 (1.38-1.34)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	323	82%	11%	
1	В	323	81%	12%	



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 4-hydroxy-3-methylbut-2-enyl diphosphate reductase.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	309	Total 2383	C 1487	N 427	O 459	S 10	0	0	0
1	В	310	Total 2390	C 1492	N 428	O 460	S 10	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	expression tag	UNP P62623
A	-6	HIS	-	expression tag	UNP P62623
A	-5	HIS	-	expression tag	UNP P62623
A	-4	HIS	-	expression tag	UNP P62623
A	-3	HIS	-	expression tag	UNP P62623
A	-2	HIS	-	expression tag	UNP P62623
A	-1	GLY	-	expression tag	UNP P62623
A	0	SER	-	expression tag	UNP P62623
В	-7	HIS	-	expression tag	UNP P62623
В	-6	HIS	-	expression tag	UNP P62623
В	-5	HIS	-	expression tag	UNP P62623
В	-4	HIS	-	expression tag	UNP P62623
В	-3	HIS	-	expression tag	UNP P62623
В	-2	HIS	-	expression tag	UNP P62623
В	-1	GLY	-	expression tag	UNP P62623
В	0	SER	-	expression tag	UNP P62623


• Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe S 8 4 4	0	0
2	В	1	Total Fe S 8 4 4	0	0

• Molecule 3 is (2E)-4-amino-3-methylbut-2-en-1-yl trihydrogen diphosphate (three-letter code: 10E) (formula: $C_5H_{13}NO_7P_2$).



\mathbf{Mol}	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 15	C 5	N 1	O 7	P 2	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	D	1	Total	С	N	О	Р	0	0
)	Б	1	15	5	1	7	2	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	374	Total O 374 374	0	0
4	В	357	Total O 357 357	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	71.32Å 80.71Å 112.11Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 - 1.35	Depositor
rtesolution (A)	48.24 - 1.35	EDS
% Data completeness	97.6 (10.00-1.35)	Depositor
(in resolution range)	97.8 (48.24-1.35)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) > 1$	3.23 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
D D.	0.191 , 0.214	Depositor
R, R_{free}	0.190 , 0.212	DCC
R_{free} test set	6957 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.698	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 38.9	EDS
L-test for twinning ²	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5550	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 46.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1482e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 10E, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	1.10	5/2418 (0.2%)	1.54	$28/3273 \ (0.9\%)$	
1	В	1.06	$2/2425 \ (0.1\%)$	1.54	$29/3283 \ (0.9\%)$	
All	All	1.08	7/4843 (0.1%)	1.54	$57/6556 \ (0.9\%)$	

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
1	В	151	TRP	CB-CG	11.59	1.71	1.50
1	A	138	GLU	CD-OE1	6.93	1.33	1.25
1	A	162	SER	CB-OG	-6.67	1.33	1.42
1	В	184	ARG	CZ-NH2	5.70	1.40	1.33
1	A	34	TYR	CE1-CZ	-5.23	1.31	1.38
1	A	133	GLN	CD-OE1	5.14	1.35	1.24
1	A	230	ARG	CD-NE	-5.04	1.37	1.46

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
1	A	230	ARG	NE-CZ-NH1	-38.32	101.14	120.30
1	A	230	ARG	NE-CZ-NH2	33.70	137.15	120.30
1	В	281	ARG	NE-CZ-NH1	25.75	133.17	120.30
1	В	230	ARG	NE-CZ-NH1	-25.52	107.54	120.30
1	В	230	ARG	NE-CZ-NH2	21.45	131.03	120.30
1	В	281	ARG	NE-CZ-NH2	-18.36	111.12	120.30
1	В	184	ARG	NE-CZ-NH1	-15.08	112.76	120.30
1	В	9	ARG	NE-CZ-NH1	13.47	127.03	120.30
1	В	52	ARG	NE-CZ-NH2	12.13	126.36	120.30
1	A	230	ARG	CD-NE-CZ	11.98	140.37	123.60
1	В	52	ARG	NE-CZ-NH1	-10.02	115.29	120.30
1	A	43	ARG	NE-CZ-NH2	-9.53	115.53	120.30



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	230	ARG	CD-NE-CZ	9.23	136.53	123.60
1	A	17	ARG	NE-CZ-NH2	9.10	124.85	120.30
1	В	11	PHE	CB-CG-CD1	9.09	127.16	120.80
1	A	203	ARG	NE-CZ-NH1	-8.96	115.82	120.30
1	В	230	ARG	CG-CD-NE	-8.86	93.19	111.80
1	A	173	ASP	CB-CG-OD1	8.83	126.25	118.30
1	В	184	ARG	NE-CZ-NH2	8.52	124.56	120.30
1	A	230	ARG	CG-CD-NE	-8.37	94.23	111.80
1	В	141	MET	CG-SD-CE	8.27	113.44	100.20
1	A	17	ARG	NE-CZ-NH1	-8.26	116.17	120.30
1	В	281	ARG	CD-NE-CZ	8.01	134.81	123.60
1	В	192	ARG	NE-CZ-NH2	-7.87	116.37	120.30
1	A	281	ARG	NE-CZ-NH2	-7.75	116.42	120.30
1	В	241	ARG	NE-CZ-NH1	-7.57	116.51	120.30
1	В	43	ARG	NE-CZ-NH2	-7.45	116.58	120.30
1	В	9	ARG	NE-CZ-NH2	-7.43	116.58	120.30
1	A	9	ARG	NE-CZ-NH1	7.39	124.00	120.30
1	В	237	ARG	NE-CZ-NH2	7.38	123.99	120.30
1	A	141	MET	CG-SD-CE	7.16	111.66	100.20
1	В	285	LEU	CB-CG-CD1	6.62	122.26	111.00
1	В	203	ARG	NE-CZ-NH1	-6.31	117.15	120.30
1	A	43	ARG	NE-CZ-NH1	6.30	123.45	120.30
1	В	51	GLU	OE1-CD-OE2	-6.18	115.88	123.30
1	A	208	ARG	NE-CZ-NH1	-6.11	117.25	120.30
1	В	36	ARG	NE-CZ-NH2	-6.10	117.25	120.30
1	A	104	MET	CG-SD-CE	6.06	109.89	100.20
1	A	192	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	A	182	ARG	NE-CZ-NH1	-5.97	117.31	120.30
1	A	208	ARG	NE-CZ-NH2	5.80	123.20	120.30
1	A	16	ASP	CB-CG-OD2	-5.66	113.20	118.30
1	A	250	ASP	CB-CG-OD1	5.66	123.40	118.30
1	A	87	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	В	173	ASP	CB-CG-OD1	5.50	123.25	118.30
1	В	151	TRP	CA-CB-CG	5.49	124.14	113.70
1	В	285	LEU	CB-CG-CD2	5.42	120.22	111.00
1	A	50	ARG	NE-CZ-NH2	-5.40	117.60	120.30
1	A	138	GLU	CA-CB-CG	5.39	125.26	113.40
1	A	173	ASP	CB-CG-OD2	-5.32	113.51	118.30
1	A	176	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	В	52	ARG	CD-NE-CZ	5.11	130.75	123.60
1	A	1	MET	CG-SD-CE	-5.09	92.06	100.20
1	A	11	PHE	CB-CG-CD1	5.06	124.34	120.80



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	11	PHE	CB-CG-CD2	-5.05	117.27	120.80
1	A	11	PHE	CB-CG-CD2	-5.04	117.27	120.80
1	В	237	ARG	NE-CZ-NH1	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2383	0	2395	22	1
1	В	2390	0	2404	33	2
2	A	8	0	0	0	0
2	В	8	0	0	0	0
3	A	15	0	12	0	0
3	В	15	0	12	0	0
4	A	374	0	0	8	1
4	В	357	0	0	13	0
All	All	5550	0	4823	55	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:281:ARG:HD3	4:B:538:HOH:O	1.76	0.84
1:B:152:LYS:HD2	4:B:606:HOH:O	1.80	0.82
1:B:272:ASP:OD1	4:B:853:HOH:O	1.96	0.80
1:B:104:MET:CE	4:B:851:HOH:O	2.30	0.80
1:B:42:ASN:HD22	1:B:45:VAL:H	1.32	0.74
1:B:112:ARG:HG3	1:B:112:ARG:HH21	1.52	0.73
1:A:103:HIS:HD2	4:A:593:HOH:O	1.72	0.72
1:A:57:ILE:HD12	1:A:63:VAL:HG22	1.71	0.72
1:B:51:GLU:OE2	1:B:52:ARG:HD2	1.90	0.72



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)	
1:A:42:ASN:HD22	1:A:45:VAL:H	1.36	0.71	
1:B:103:HIS:HD2	4:B:566:HOH:O	1.75	0.69	
1:B:104:MET:HE2	4:B:851:HOH:O	1.91	0.68	
1:A:57:ILE:CD1	1:A:63:VAL:HG22	2.26	0.65	
1:A:276:GLN:NE2	4:A:635:HOH:O	2.30	0.65	
1:B:104:MET:HE1	4:B:851:HOH:O	1.93	0.64	
1:A:297:GLU:HG3	4:A:571:HOH:O	1.97	0.63	
1:A:294:GLU:HG3	4:A:819:HOH:O	1.98	0.63	
1:B:115:GLU:CD	1:B:155:VAL:HG13	2.20	0.62	
1:B:121:HIS:HD2	4:B:678:HOH:O	1.82	0.62	
4:A:571:HOH:O	1:B:306:LYS:HG2	2.00	0.60	
1:A:1:MET:N	4:A:587:HOH:O	2.36	0.58	
1:A:136:ASN:HD22	1:A:136:ASN:C	2.06	0.58	
1:B:241:ARG:HD3	4:B:649:HOH:O	2.03	0.58	
1:A:213:GLN:NE2	4:A:855:HOH:O	2.41	0.54	
1:B:68:ILE:HD13	1:B:90:THR:HB	1.91	0.53	
1:A:149:ASP:OD1	1:A:152:LYS:HE3	2.10	0.52	
1:B:184:ARG:NE	4:B:849:HOH:O	2.28	0.51	
1:A:149:ASP:HA	1:A:152:LYS:HE3	1.94	0.50	
1:B:51:GLU:OE2	1:B:52:ARG:CD	2.57	0.49	
1:B:115:GLU:OE2	1:B:155:VAL:CG1	2.60	0.49	
1:A:103:HIS:HE1	4:A:503:HOH:O	1.95	0.49	
1:A:136:ASN:ND2	1:A:138:GLU:H	2.10	0.49	
1:B:8:PRO:HG2	1:B:202:ASN:HB3	1.95	0.49	
1:A:42:ASN:HD21	1:A:44:TYR:HB3	1.77	0.49	
1:B:112:ARG:HH21	1:B:112:ARG:CG	2.22	0.48	
1:B:121:HIS:CD2	4:B:678:HOH:O	2.61	0.48	
1:B:115:GLU:OE1	1:B:155:VAL:HG13	2.15	0.46	
1:B:60:ILE:HD11	1:B:80:VAL:HG13	1.97	0.46	
1:A:136:ASN:HD22	1:A:138:GLU:H	1.65	0.45	
1:B:42:ASN:HD21	1:B:44:TYR:HB3	1.82	0.45	
1:B:103:HIS:HE1	4:B:507:HOH:O	2.00	0.44	
1:B:136:ASN:HD22	1:B:138:GLU:H	1.66	0.44	
1:B:249:LYS:HB2	1:B:249:LYS:HE2	1.33	0.43	
1:A:148:ASP:O	1:A:152:LYS:HG2	2.19	0.43	
1:B:136:ASN:HD22	1:B:136:ASN:C	2.23	0.42	
1:B:208:ARG:O	1:B:212:GLU:HG3	2.19	0.42	
1:A:8:PRO:HG2	1:A:202:ASN:HB3	2.01	0.42	
1:B:156:LYS:HE3	4:B:647:HOH:O	2.20	0.42	
1:B:230:ARG:HA	1:B:230:ARG:HD3	1.76	0.42	
1:A:59:GLN:HE21	1:A:59:GLN:HB3	1.57	0.41	



$\alpha \cdots$, r	•	
Continued	trom	mromonie	maaa
-	110116	DICULUUS	Duuc
	J	1	1

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	Clash overlap (Å)
1:A:217:VAL:O	1:A:242:ALA:HA	2.20	0.41
1:B:115:GLU:OE2	1:B:155:VAL:HG13	2.20	0.41
1:B:85:LYS:HB3	1:B:85:LYS:HE2	1.65	0.41
1:A:70:ILE:HA	1:A:92:PHE:O	2.19	0.41
1:A:34:TYR:HB3	1:A:57:ILE:HD11	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:276:GLN:NE2	4:A:635:HOH:O[4_445]	2.09	0.11
1:A:253:GLU:OE1	1:B:52:ARG:NH1[4_545]	2.16	0.04

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	307/323~(95%)	303 (99%)	4 (1%)	0	100	100
1	В	$308/323 \ (95\%)$	304 (99%)	4 (1%)	0	100	100
All	All	615/646~(95%)	607 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	257/270~(95%)	252 (98%)	5 (2%)	57 23
1	В	258/270 (96%)	245 (95%)	13 (5%)	24 2
All	All	515/540 (95%)	497 (96%)	18 (4%)	36 6

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	ILE
1	A	103	HIS
1	A	136	ASN
1	A	230	ARG
1	A	294	GLU
1	В	85	LYS
1	В	103	HIS
1	В	108	ARG
1	В	136	ASN
1	В	148	ASP
1	В	153	LEU
1	В	154	THR
1	В	159	GLU
1	В	230	ARG
1	В	249	LYS
1	В	281	ARG
1	В	285	LEU
1	В	299	ASN

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such side chains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	59	GLN
1	A	78	GLN
1	A	103	HIS
1	A	133	GLN
1	A	136	ASN
1	A	229	ASN
1	A	236	GLN
1	A	252	GLN
1	A	277	ASN
1	В	42	ASN
1	В	78	GLN



Continued from previous page...

Mol	Chain	Res	Type
1	В	103	HIS
1	В	121	HIS
1	В	133	GLN
1	В	136	ASN
1	В	277	ASN
1	В	283	GLN
1	В	299	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Cha		Chain	Chain Res	Res	Res	Link	Bo	Bond lengths			Bond angles		
Mol Type Chain	ries	nes				nes	nes	nes	Lilik	Counts	RMSZ	# Z > 2	Counts
3	10E	В	402	2	12,14,14	1.00	1 (8%)	13,20,20	1.32	2 (15%)			
3	10E	A	402	2	12,14,14	1.93	2 (16%)	13,20,20	1.11	1 (7%)			
2	SF4	A	401	1,3	0,12,12	-	-	-					
2	SF4	В	401	1,3	0,12,12	-	-	-					

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	10E	В	402	2	-	2/13/15/15	-
2	SF4	В	401	1,3	-	-	0/6/5/5
2	SF4	A	401	1,3	-	-	0/6/5/5
3	10E	A	402	2	-	1/13/15/15	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	A	402	10E	O7-C6	5.65	1.52	1.43
3	A	402	10E	C4-C3	-2.82	1.43	1.50
3	В	402	10E	O7-C6	2.20	1.47	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
3	В	402	10E	O15-P12-O14	2.65	117.76	107.64
3	В	402	10E	C6-C5-C3	-2.43	121.83	126.04
3	A	402	10E	P8-O11-P12	2.08	139.96	132.83

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	402	10E	P8-O11-P12-O15
3	A	402	10E	C5-C6-O7-P8
3	В	402	10E	C5-C6-O7-P8


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		$OWAB(A^2)$	Q < 0.9
1	A	309/323~(95%)	-0.18	5 (1%)	72 76	9, 16, 30, 45	0
1	В	310/323~(95%)	-0.15	1 (0%)	94 95	10, 17, 31, 51	0
All	All	619/646 (95%)	-0.17	6 (0%)	82 85	9, 16, 30, 51	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	88	ASP	3.6
1	В	310	VAL	2.9
1	A	309	ARG	2.4
1	A	137	PRO	2.2
1	A	289	GLU	2.1
1	A	65	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	SF4	A	401	8/8	0.98	0.07	9,10,11,17	0
2	SF4	В	401	8/8	0.98	0.07	9,10,11,17	0
3	10E	В	402	15/15	0.98	0.07	9,10,14,18	0
3	10E	A	402	15/15	0.99	0.06	8,10,13,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

