

wwPDB X-ray Structure Validation Summary Report (i)

May 14, 2020 - 08:52 am BST

PDB ID	:	2H5P
Title	:	Crystal structure of mStrawberry at pH 9.5
Authors	:	Shu, X.; Remington, S.J.
Deposited on		
$\operatorname{Resolution}$:	1.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\rm CCP4$:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R _{free}	130704	1232 (1.24-1.20)
Clashscore	141614	1294 (1.24-1.20)
Ramachandran outliers	138981	1251(1.24-1.20)
Sidechain outliers	138945	1250 (1.24-1.20)
RSRZ outliers	127900	1209 (1.24-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
			9% ■				
1	A	234	78%	14%	• 5%		

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	CRO	А	66	Х	-	-	-



2H5P

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2050 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called mStrawberry.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	223	Total 1804	$\begin{array}{c} \mathrm{C} \\ 1155 \end{array}$	N 295	O 343	S 11	0	7	0


• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	246	Total O 246 246	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: mStrawberry



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	48.60Å 43.44 Å 61.36 Å	Depositor
a, b, c, α , β , γ	90.00° 111.84° 90.00°	Depositor
Resolution (Å)	10.00 - 1.21	Depositor
Resolution (A)	31.29 - 1.21	EDS
% Data completeness	$97.1\ (10.00-1.21)$	Depositor
(in resolution range)	92.3 (31.29 - 1.21)	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.19 (at 1.21 \text{\AA})$	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.138 , 0.178	Depositor
n, n <i>free</i>	0.146 , 0.149	DCC
R_{free} test set	3520 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	12.9	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38 , 60.8	EDS
L-test for twinning ²	$< L > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2050	wwPDB-VP
Average B, all atoms $(Å^2)$	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.84% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



 $^{^1 {\}rm Intensities}$ estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.79	2/1855~(0.1%)	1.44	26/2495~(1.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	1	0

All (2) bond length outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	А	21	SER	CB-OG	-5.82	1.34	1.42
1	А	65	PHE	CA-C	5.28	1.66	1.52

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	216	ARG	NE-CZ-NH2	-15.47	112.56	120.30
1	А	216	ARG	NE-CZ-NH1	12.66	126.63	120.30
1	А	128[A]	MET	CG-SD-CE	12.44	120.11	100.20
1	А	128[B]	MET	CG-SD-CE	12.44	120.11	100.20
1	А	123	LYS	CB-CG-CD	9.54	136.40	111.60

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	А	66	CRO	CA1



There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1804	0	1767	20	0
2	А	246	0	0	6	0
All	All	2050	0	1767	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 20 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215[B]:GLU:OE1	2:A:361:HOH:O	1.86	0.93
1:A:92:LYS:HE2	2:A:537:HOH:O	1.91	0.71
1:A:199[A]:LEU:HD23	1:A:215[A]:GLU:HB2	1.75	0.68
1:A:108[A]:THR:HG22	2:A:346:HOH:O	1.99	0.62
1:A:139:LYS:HD2	2:A:541:HOH:O	2.00	0.60

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	225/234~(96%)	220~(98%)	5(2%)	0	100 100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	192/197~(98%)	185~(96%)	7~(4%)	35 5	

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	58	TRP
1	А	149	ARG
1	А	128[A]	MET
1	А	12[B]	MET
1	А	128[B]	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	B	ond leng	gths	B	ond ang	gles
	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
1	CRO	А	66	1	23,23,24	<mark>3.23</mark>	12 (52%)	30,32,34	2.94	10 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	А	66	1	1/1/6/8	5/12/31/32	0/2/2/2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	А	66	CRO	CA1-C1	-8.94	1.38	1.51
1	А	66	CRO	CG1-CB1	-4.97	1.37	1.51
1	А	66	CRO	OG1-CB1	4.82	1.56	1.43
1	А	66	CRO	CA1-N1	-4.65	1.33	1.47
1	А	66	CRO	CG2-CB2	-3.75	1.39	1.46

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$\mathbf{Ideal}(^{o})$
1	А	66	CRO	C1-CA1-N1	9.86	125.96	109.96
1	А	66	CRO	O2-C2-CA2	7.96	135.43	130.96
1	А	66	CRO	CA2-C2-N3	4.71	105.60	103.37
1	А	66	CRO	CA2-N2-C1	3.39	108.27	105.77
1	А	66	CRO	O2-C2-N3	-3.26	117.87	124.35

All (1) chirality outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atom
1	А	66	CRO	CA1

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	66	CRO	C1-CA1-CB1-CG1
1	А	66	CRO	C1-CA1-CB1-OG1
1	А	66	CRO	N2-C1-CA1-CB1
1	А	66	CRO	N3-C1-CA1-CB1
1	А	66	CRO	C2-CA2-CB2-CG2



There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$<$ RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	222/234~(94%)	0.34	3 (1%) 75 73	12, 18, 34, 59	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	А	6	ALA	3.9	
1	А	186	PRO	3.2	
1	А	199[A]	LEU	2.9	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	$B-factors(Å^2)$	$\mathbf{Q}{<}0.9$
1	CRO	А	66	22/23	0.97	0.09	$13,\!17,\!23,\!26$	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.



6.5 Other polymers (i)

There are no such residues in this entry.

