



wwPDB X-ray Structure Validation Summary Report

Sep 10, 2023 – 06:27 AM EDT

PDB ID : 4H5F
Title : Crystal structure of an amino acid ABC transporter substrate-binding protein from *Streptococcus pneumoniae* Canada MDR_19A bound to L-arginine, form 1
Authors : Stogios, P.J.; Wawrzak, Z.; Kudritska, M.; Minasov, G.; Yim, V.; Savchenko, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2012-09-18
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	 87% 11% ..
1	B	243	 91% 8% .
1	C	243	 80% 19% .
1	D	243	 80% 18% ..

Validation Pipeline (wwPDB-VP) : 2.35.1

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	B	305	-	-	X	-

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 8142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amino acid ABC superfamily ATP binding cassette transporter, binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	240	Total 1855	C 1167	N 300	O 381	S 7	0	9	0
1	B	240	Total 1824	C 1148	N 297	O 373	S 6	0	2	0
1	C	240	Total 1812	C 1140	N 295	O 371	S 6	0	0	0
1	D	239	Total 1803	C 1135	N 293	O 369	S 6	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	29	GLY	-	expression tag	UNP D6ZRZ2
A	136	HIS	ARG	SEE REMARK 999	UNP D6ZRZ2
B	29	GLY	-	expression tag	UNP D6ZRZ2
B	136	HIS	ARG	SEE REMARK 999	UNP D6ZRZ2
C	29	GLY	-	expression tag	UNP D6ZRZ2
C	136	HIS	ARG	SEE REMARK 999	UNP D6ZRZ2
D	29	GLY	-	expression tag	UNP D6ZRZ2
D	136	HIS	ARG	SEE REMARK 999	UNP D6ZRZ2

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0
4	C	1	Total C O 7 4 3	0	0
4	C	1	Total C O 7 4 3	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 7 4 3	0	0
4	C	1	Total C O 7 4 3	0	0
4	C	1	Total C O 7 4 3	0	0
4	D	1	Total C O 7 4 3	0	0
4	D	1	Total C O 7 4 3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 10 6 4	0	0
6	B	1	Total C O 10 6 4	0	0
6	B	1	Total C O 10 6 4	0	0
6	C	1	Total C O 10 6 4	0	0

- Molecule 7 is ARGinine (three-letter code: ARG) (formula: $C_6H_{15}N_4O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			12	6	4	2		
7	B	1	Total	C	N	O	0	0
			12	6	4	2		
7	C	1	Total	C	N	O	0	0
			12	6	4	2		
7	D	1	Total	C	N	O	0	0
			12	6	4	2		

- Molecule 8 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	C	1	Total	C	O	0	0
			4	2	2		
8	C	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total	C O	0	0
			4	2 2		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	157	Total	O	0	14
			171	171		
10	B	138	Total	O	0	13
			151	151		
10	C	114	Total	O	0	4
			118	118		
10	D	89	Total	O	0	5
			93	93		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Amino acid ABC superfamily ATP binding cassette transporter, binding protein




- Molecule 1: Amino acid ABC superfamily ATP binding cassette transporter, binding protein




- Molecule 1: Amino acid ABC superfamily ATP binding cassette transporter, binding protein



- Molecule 1: Amino acid ABC superfamily ATP binding cassette transporter, binding protein





4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	99.37Å 61.70Å 103.63Å 90.00° 118.04° 90.00°	Depositor
Resolution (Å)	91.46 – 1.90 91.46 – 1.90	Depositor EDS
% Data completeness (in resolution range)	94.8 (91.46-1.90) 86.4 (91.46-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 1.90Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.164 , 0.200 0.162 , 0.184	Depositor DCC
R_{free} test set	4107 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtrriage
Anisotropy	0.465	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 67.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.248 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8142	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO, PGE, GOL, PEG, ACT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/1887	0.55	0/2548
1	B	0.44	0/1847	0.55	0/2494
1	C	0.39	0/1832	0.53	0/2475
1	D	0.40	0/1823	0.56	0/2463
All	All	0.41	0/7389	0.55	0/9980

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1855	0	1884	18	0
1	B	1824	0	1852	16	0
1	C	1812	0	1835	32	0
1	D	1803	0	1827	25	0
2	A	15	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	1	0	0	0	0
4	A	56	0	71	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	28	0	36	6	0
4	C	35	0	41	2	0
4	D	14	0	16	2	0
5	A	18	0	24	0	0
5	C	12	0	16	3	0
5	D	6	0	8	0	0
6	A	10	0	14	1	0
6	B	20	0	28	0	0
6	C	10	0	14	2	0
7	A	12	0	12	0	0
7	B	12	0	12	0	0
7	C	12	0	12	0	0
7	D	12	0	12	0	0
8	B	16	0	12	0	0
8	C	8	0	6	0	0
8	D	4	0	3	0	0
9	B	4	0	6	1	0
10	A	171	0	0	9	0
10	B	151	0	0	2	0
10	C	118	0	0	6	0
10	D	93	0	0	3	0
All	All	8142	0	7741	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:189:GLN:HE22	4:B:306:PEG:H42	1.38	0.89
1:C:131:ILE:HG21	1:C:165:MET:HE3	1.53	0.89
1:B:114:ASP:HB2	9:B:303:EDO:H21	1.69	0.74
1:D:149:LEU:HD13	1:D:170:LEU:HD11	1.71	0.72
1:C:245:LYS:NZ	10:C:500:HOH:O	2.25	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/243 (102%)	241 (98%)	6 (2%)	0	100	100
1	B	240/243 (99%)	231 (96%)	9 (4%)	0	100	100
1	C	238/243 (98%)	233 (98%)	5 (2%)	0	100	100
1	D	237/243 (98%)	225 (95%)	10 (4%)	2 (1%)	19	9
All	All	962/972 (99%)	930 (97%)	30 (3%)	2 (0%)	47	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	211	ALA
1	D	212	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/199 (102%)	196 (96%)	8 (4%)	32	23
1	B	199/199 (100%)	195 (98%)	4 (2%)	55	51
1	C	197/199 (99%)	190 (96%)	7 (4%)	35	26
1	D	196/199 (98%)	185 (94%)	11 (6%)	21	11
All	All	796/796 (100%)	766 (96%)	30 (4%)	33	24

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	50	TYR
1	D	224	ASP
1	C	217	THR
1	D	236	SER
1	D	176	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	GLN
1	A	257	GLN
1	B	189	GLN
1	B	257	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 47 ligands modelled in this entry, 1 is monoatomic - leaving 46 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PGE	A	315	-	9,9,9	0.41	0	8,8,8	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	ARG	C	312	-	10,11,11	0.74	1 (10%)	11,13,13	1.07	1 (9%)
4	PEG	C	303	-	6,6,6	1.05	0	5,5,5	1.88	2 (40%)
8	ACT	B	308	-	3,3,3	0.79	0	3,3,3	1.29	0
4	PEG	C	305	-	6,6,6	1.06	0	5,5,5	1.86	2 (40%)
7	ARG	B	313	-	10,11,11	0.84	1 (10%)	11,13,13	0.99	1 (9%)
5	GOL	C	308	-	5,5,5	0.42	0	5,5,5	0.46	0
7	ARG	D	305	-	10,11,11	0.78	1 (10%)	11,13,13	1.06	2 (18%)
7	ARG	A	317	-	10,11,11	0.81	1 (10%)	11,13,13	1.22	2 (18%)
8	ACT	B	309	-	3,3,3	0.79	0	3,3,3	1.37	0
4	PEG	B	304	-	6,6,6	1.00	0	5,5,5	1.99	2 (40%)
9	EDO	B	303	-	3,3,3	0.42	0	2,2,2	0.46	0
8	ACT	B	302	-	3,3,3	0.76	0	3,3,3	1.52	0
2	SO4	A	301	-	4,4,4	0.15	0	6,6,6	0.28	0
4	PEG	A	311	-	6,6,6	1.09	0	5,5,5	2.68	3 (60%)
4	PEG	A	306	-	6,6,6	0.99	0	5,5,5	1.88	2 (40%)
5	GOL	A	312	-	5,5,5	0.37	0	5,5,5	0.28	0
2	SO4	B	301	-	4,4,4	0.14	0	6,6,6	0.16	0
4	PEG	A	308	-	6,6,6	0.97	0	5,5,5	2.75	3 (60%)
5	GOL	A	313	-	5,5,5	0.29	0	5,5,5	0.40	0
4	PEG	C	306	-	6,6,6	1.07	0	5,5,5	1.78	1 (20%)
4	PEG	B	306	-	6,6,6	1.14	0	5,5,5	1.42	0
6	PGE	B	312	-	9,9,9	0.70	0	8,8,8	0.51	0
8	ACT	C	310	-	3,3,3	0.79	0	3,3,3	1.42	0
4	PEG	A	305	-	6,6,6	0.99	0	5,5,5	1.92	1 (20%)
4	PEG	B	307	-	6,6,6	0.96	0	5,5,5	2.00	2 (40%)
4	PEG	B	305	-	6,6,6	0.96	0	5,5,5	2.05	2 (40%)
4	PEG	A	310	-	6,6,6	1.04	0	5,5,5	1.54	0
4	PEG	C	302	-	6,6,6	1.01	0	5,5,5	1.56	1 (20%)
5	GOL	C	307	-	5,5,5	0.36	0	5,5,5	0.34	0
4	PEG	A	307	-	6,6,6	1.12	0	5,5,5	2.60	3 (60%)
4	PEG	A	316	-	6,6,6	1.05	0	5,5,5	1.76	2 (40%)
8	ACT	B	310	-	3,3,3	0.72	0	3,3,3	1.39	0
6	PGE	B	311	-	9,9,9	0.57	0	8,8,8	0.63	0
2	SO4	A	302	-	4,4,4	0.16	0	6,6,6	0.25	0
2	SO4	C	301	-	4,4,4	0.13	0	6,6,6	0.18	0
4	PEG	D	301	-	6,6,6	1.17	0	5,5,5	1.57	2 (40%)
6	PGE	C	311	-	9,9,9	0.65	0	8,8,8	0.37	0
5	GOL	A	314	-	5,5,5	0.34	0	5,5,5	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PEG	C	304	-	6,6,6	1.04	0	5,5,5	2.04	2 (40%)
5	GOL	D	303	-	5,5,5	0.43	0	5,5,5	0.21	0
4	PEG	D	302	-	6,6,6	1.17	0	5,5,5	1.26	0
8	ACT	C	309	-	3,3,3	0.77	0	3,3,3	1.41	0
4	PEG	A	309	-	6,6,6	1.09	0	5,5,5	1.48	0
2	SO4	A	303	-	4,4,4	0.16	0	6,6,6	0.27	0
8	ACT	D	304	-	3,3,3	0.79	0	3,3,3	1.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	A	315	-	-	3/7/7/7	-
4	PEG	C	303	-	-	2/4/4/4	-
4	PEG	C	305	-	-	1/4/4/4	-
7	ARG	B	313	-	-	0/11/11/11	-
5	GOL	C	308	-	-	2/4/4/4	-
7	ARG	D	305	-	-	0/11/11/11	-
7	ARG	A	317	-	-	0/11/11/11	-
4	PEG	B	304	-	-	1/4/4/4	-
9	EDO	B	303	-	-	1/1/1/1	-
4	PEG	A	311	-	-	2/4/4/4	-
4	PEG	A	306	-	-	2/4/4/4	-
5	GOL	A	312	-	-	2/4/4/4	-
4	PEG	A	308	-	-	0/4/4/4	-
5	GOL	A	313	-	-	2/4/4/4	-
4	PEG	C	306	-	-	1/4/4/4	-
4	PEG	B	306	-	-	2/4/4/4	-
6	PGE	B	312	-	-	2/7/7/7	-
4	PEG	A	305	-	-	2/4/4/4	-
4	PEG	B	307	-	-	1/4/4/4	-
4	PEG	B	305	-	-	1/4/4/4	-
4	PEG	A	310	-	-	1/4/4/4	-
4	PEG	C	302	-	-	1/4/4/4	-
5	GOL	C	307	-	-	0/4/4/4	-
4	PEG	A	307	-	-	2/4/4/4	-
4	PEG	A	316	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	B	311	-	-	2/7/7/7	-
4	PEG	D	301	-	-	0/4/4/4	-
6	PGE	C	311	-	-	3/7/7/7	-
5	GOL	A	314	-	-	4/4/4/4	-
4	PEG	C	304	-	-	2/4/4/4	-
5	GOL	D	303	-	-	2/4/4/4	-
4	PEG	D	302	-	-	0/4/4/4	-
4	PEG	A	309	-	-	1/4/4/4	-
7	ARG	C	312	-	-	0/11/11/11	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	317	ARG	OXT-C	-2.27	1.23	1.30
7	B	313	ARG	OXT-C	-2.25	1.23	1.30
7	D	305	ARG	OXT-C	-2.21	1.23	1.30
7	C	312	ARG	OXT-C	-2.10	1.23	1.30

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	311	PEG	O2-C2-C1	4.57	130.14	110.07
4	A	308	PEG	O2-C2-C1	3.70	126.31	110.07
4	A	308	PEG	O2-C3-C4	3.59	125.86	110.07
4	A	307	PEG	O2-C2-C1	3.54	125.61	110.07
4	A	307	PEG	O2-C3-C4	3.28	124.48	110.07

There are no chirality outliers.

5 of 47 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	312	GOL	O1-C1-C2-C3
5	A	313	GOL	O1-C1-C2-O2
5	A	314	GOL	O1-C1-C2-C3
5	C	308	GOL	C1-C2-C3-O3
5	D	303	GOL	O1-C1-C2-O2

There are no ring outliers.

14 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	315	PGE	1	0
4	C	303	PEG	1	0
5	C	308	GOL	2	0
9	B	303	EDO	1	0
4	A	311	PEG	1	0
4	A	308	PEG	1	0
4	B	306	PEG	1	0
4	B	305	PEG	5	0
4	A	310	PEG	1	0
4	C	302	PEG	1	0
5	C	307	GOL	1	0
4	D	301	PEG	1	0
6	C	311	PGE	2	0
4	D	302	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/243 (98%)	-0.44	3 (1%) 77 79	16, 23, 59, 89	0
1	B	240/243 (98%)	-0.36	6 (2%) 57 60	17, 25, 63, 102	0
1	C	240/243 (98%)	-0.05	16 (6%) 17 20	19, 33, 78, 93	0
1	D	239/243 (98%)	0.18	27 (11%) 5 5	19, 36, 90, 118	0
All	All	959/972 (98%)	-0.17	52 (5%) 25 29	16, 28, 76, 118	0

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	211	ALA	7.4
1	D	147	THR	7.4
1	D	210	ASN	6.8
1	D	143	TYR	6.1
1	D	209	LYS	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	ACT	B	309	4/4	0.32	0.23	70,71,72,74	0
8	ACT	B	310	4/4	0.57	0.24	63,66,67,71	0
5	GOL	A	313	6/6	0.58	0.18	61,63,66,66	0
4	PEG	D	302	7/7	0.59	0.21	56,61,70,73	0
8	ACT	D	304	4/4	0.67	0.26	81,81,82,84	0
4	PEG	C	303	7/7	0.69	0.33	65,72,78,78	0
4	PEG	C	304	7/7	0.73	0.17	51,53,55,56	0
4	PEG	B	307	7/7	0.74	0.20	44,49,52,55	0
5	GOL	C	307	6/6	0.75	0.13	66,70,75,78	0
4	PEG	A	316	7/7	0.76	0.19	54,62,62,62	0
4	PEG	C	306	7/7	0.78	0.19	51,55,63,63	0
8	ACT	B	308	4/4	0.79	0.23	67,68,69,70	0
6	PGE	C	311	10/10	0.79	0.14	46,52,60,61	0
4	PEG	D	301	7/7	0.80	0.14	45,48,55,59	0
4	PEG	C	302	7/7	0.80	0.18	43,47,51,51	0
4	PEG	A	308	7/7	0.80	0.17	47,54,56,57	0
4	PEG	A	310	7/7	0.82	0.15	47,55,59,59	0
6	PGE	B	312	10/10	0.83	0.16	32,47,59,59	0
9	EDO	B	303	4/4	0.83	0.13	39,39,40,46	0
8	ACT	C	309	4/4	0.84	0.13	51,53,53,53	0
4	PEG	C	305	7/7	0.85	0.18	45,46,48,52	0
4	PEG	A	311	7/7	0.86	0.14	43,50,53,56	0
4	PEG	A	306	7/7	0.86	0.11	47,48,50,50	0
4	PEG	A	309	7/7	0.87	0.11	36,39,51,55	0
4	PEG	B	304	7/7	0.87	0.12	39,42,51,51	0
6	PGE	A	315	10/10	0.88	0.21	45,51,53,54	0
4	PEG	A	305	7/7	0.89	0.12	30,33,39,44	0
5	GOL	A	312	6/6	0.89	0.20	44,47,50,50	0
5	GOL	C	308	6/6	0.89	0.14	30,48,50,52	0
5	GOL	D	303	6/6	0.90	0.15	52,54,57,60	0
4	PEG	A	307	7/7	0.90	0.12	39,41,45,46	0
4	PEG	B	305	7/7	0.92	0.12	39,41,44,45	0
4	PEG	B	306	7/7	0.92	0.10	45,46,48,48	0
5	GOL	A	314	6/6	0.92	0.11	47,53,55,57	0
8	ACT	C	310	4/4	0.93	0.08	60,61,62,62	0
2	SO4	A	302	5/5	0.93	0.12	65,70,71,71	0
6	PGE	B	311	10/10	0.93	0.12	38,47,51,52	0
7	ARG	C	312	12/12	0.95	0.07	19,22,29,30	0
3	CL	A	304	1/1	0.96	0.04	40,40,40,40	0
7	ARG	D	305	12/12	0.97	0.07	21,25,29,31	0
8	ACT	B	302	4/4	0.97	0.09	39,41,43,44	0
2	SO4	A	303	5/5	0.97	0.10	43,48,52,53	0
7	ARG	A	317	12/12	0.98	0.09	16,16,18,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	ARG	B	313	12/12	0.98	0.08	16,17,21,21	0
2	SO4	A	301	5/5	0.98	0.09	41,42,44,44	0
2	SO4	C	301	5/5	0.98	0.07	43,47,49,52	0
2	SO4	B	301	5/5	0.99	0.11	38,41,46,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.