



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 12, 2022 – 06:11 PM JST

PDB ID : 5H5M
Title : Crystal structure of HMP-1 M domain
Authors : Kang, H.; Bang, I.; Weis, W.I.; Choi, H.J.
Deposited on : 2016-11-08
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.31.3
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	 2% 89% 5% 6%
1	B	381	 8% 76% 15% 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5785 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-catenin-like protein hmp-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	358	2831	1737	522	558	14	0	0	0
1	B	349	2756	1693	506	544	13	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	266	GLY	-	expression tag	UNP P90947
A	267	GLY	-	expression tag	UNP P90947
A	268	ILE	-	expression tag	UNP P90947
A	269	GLN	-	expression tag	UNP P90947
B	266	GLY	-	expression tag	UNP P90947
B	267	GLY	-	expression tag	UNP P90947
B	268	ILE	-	expression tag	UNP P90947
B	269	GLN	-	expression tag	UNP P90947


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	132	Total	O	0	0
			132	132		
2	B	66	Total	O	0	0
			66	66		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-catenin-like protein hmp-1



- Molecule 1: Alpha-catenin-like protein hmp-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.85Å 81.53Å 151.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.16 – 2.40 27.16 – 2.37	Depositor EDS
% Data completeness (in resolution range)	99.4 (27.16-2.40) 97.8 (27.16-2.37)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 2.36Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.203 , 0.248 0.207 , 0.248	Depositor DCC
R_{free} test set	1842 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtrriage
Anisotropy	0.109	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5785	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2865	0.40	0/3868
1	B	0.27	0/2787	0.46	0/3761
All	All	0.25	0/5652	0.43	0/7629

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2831	0	2801	11	0
1	B	2756	0	2731	41	0
2	A	132	0	0	1	0
2	B	66	0	0	2	0
All	All	5785	0	5532	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:270:GLY:HA3	1:B:433:GLN:HA	1.71	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:286:ASP:O	1:B:289:HIS:HB2	1.92	0.67
1:B:600:ARG:O	1:B:604:HIS:HB2	1.94	0.66
1:B:330:GLU:HG3	1:B:375:HIS:CD2	2.30	0.66
1:B:516:ASP:OD2	1:B:551:ARG:NH1	2.34	0.60
1:B:577:GLN:HB2	1:B:580:ARG:CZ	2.34	0.58
1:B:577:GLN:HB2	1:B:580:ARG:NH2	2.18	0.58
1:A:296:ARG:HG3	1:A:342:LEU:HD13	1.87	0.57
1:B:381:VAL:HG11	1:B:505:PHE:HA	1.87	0.56
1:B:500:THR:HG22	2:B:760:HOH:O	2.05	0.56
1:B:519:ARG:HG3	1:B:522:LYS:HE3	1.87	0.55
1:B:404:GLU:N	1:B:405:GLY:HA2	2.21	0.54
1:B:278:THR:O	1:B:282:ARG:HG2	2.09	0.53
1:A:404:GLU:N	1:A:405:GLY:HA2	2.24	0.52
1:A:623:ALA:O	1:A:626:LEU:HB2	2.10	0.51
1:B:614:GLU:HA	1:B:617:MET:HG2	1.93	0.49
1:B:292:ARG:O	1:B:292:ARG:HD3	2.11	0.49
1:B:273:ILE:HB	1:B:434:LEU:HA	1.94	0.49
1:A:323:ARG:HD2	1:A:382:ASP:OD1	2.13	0.48
1:B:547:SER:HB3	1:B:551:ARG:NH1	2.28	0.48
1:B:290:TYR:OH	1:B:292:ARG:NH2	2.39	0.47
1:B:577:GLN:HA	1:B:580:ARG:HG2	1.96	0.47
1:B:406:HIS:HB3	1:B:409:ASN:HB2	1.97	0.47
1:B:613:LYS:HD3	1:B:613:LYS:H	1.81	0.46
1:B:459:ALA:O	1:B:463:ILE:HG12	2.16	0.46
1:B:368:ARG:HA	1:B:368:ARG:HD3	1.70	0.46
1:B:496:LEU:O	1:B:500:THR:HG23	2.16	0.46
1:B:577:GLN:HA	1:B:580:ARG:CG	2.46	0.45
1:A:299:LEU:HD12	1:A:299:LEU:HA	1.85	0.45
1:B:611:LYS:O	1:B:615:GLU:HG2	2.17	0.45
1:A:295:ASP:O	1:A:299:LEU:HB2	2.17	0.44
1:B:377:ARG:O	1:B:381:VAL:HG23	2.18	0.43
1:B:381:VAL:HB	1:B:508:VAL:HG21	2.00	0.43
1:B:574:THR:O	1:B:577:GLN:HG2	2.19	0.42
1:B:358:ILE:HB	1:B:359:PRO:HD3	2.01	0.42
1:A:373:ARG:O	1:A:377:ARG:HG2	2.20	0.42
1:B:299:LEU:HD12	1:B:299:LEU:HA	1.89	0.42
1:B:358:ILE:O	1:B:362:ILE:HG12	2.19	0.42
1:A:300:GLU:O	1:A:304:GLU:HG2	2.21	0.41
1:B:404:GLU:H	1:B:405:GLY:HA2	1.84	0.41
1:B:296:ARG:NH2	1:B:339:GLN:OE1	2.53	0.41
1:A:343:THR:HG22	1:A:347:LYS:HE3	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:292:ARG:NH1	1:B:296:ARG:HH11	2.19	0.41
1:B:299:LEU:HD23	1:B:342:LEU:HD21	2.02	0.41
1:B:292:ARG:HH12	1:B:346:GLU:CD	2.24	0.41
1:B:332:ASN:ND2	2:B:703:HOH:O	2.50	0.41
1:B:517:CYS:O	1:B:521:ILE:HG12	2.21	0.41
1:A:319:THR:HA	2:A:715:HOH:O	2.20	0.40
1:B:505:PHE:CD2	1:B:558:VAL:HG21	2.56	0.40
1:B:472:LYS:O	1:B:476:GLU:HG3	2.20	0.40
1:A:273:ILE:HB	1:A:434:LEU:HA	2.02	0.40
1:B:379:ALA:O	1:B:382:ASP:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/381 (92%)	345 (98%)	7 (2%)	0	100	100
1	B	341/381 (90%)	330 (97%)	11 (3%)	0	100	100
All	All	693/762 (91%)	675 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/326 (94%)	307 (100%)	1 (0%)	92	97
1	B	300/326 (92%)	295 (98%)	5 (2%)	60	78
All	All	608/652 (93%)	602 (99%)	6 (1%)	76	88

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	641	ASN
1	B	271	ASP
1	B	404	GLU
1	B	411	ARG
1	B	540	THR
1	B	602	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	325	GLN
1	B	599	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	358/381 (93%)	-0.19	8 (2%) 62 60	19, 36, 60, 84	0
1	B	349/381 (91%)	0.39	31 (8%) 9 9	23, 45, 85, 111	0
All	All	707/762 (92%)	0.10	39 (5%) 25 24	19, 41, 78, 111	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	289	HIS	5.2
1	A	606	LEU	4.9
1	B	356	ASP	4.8
1	B	355	ASN	4.7
1	B	565	PHE	4.3
1	A	538	CYS	4.3
1	B	602	GLU	4.1
1	B	598	ALA	4.1
1	B	606	LEU	4.1
1	B	599	ASN	4.0
1	B	604	HIS	3.8
1	B	609	ASP	3.8
1	B	610	PRO	3.8
1	B	612	THR	3.6
1	B	291	ARG	3.5
1	B	470	THR	3.3
1	B	603	ALA	3.3
1	A	317	GLU	3.3
1	B	357	ASP	2.9
1	B	270	GLY	2.8
1	A	640	MET	2.8
1	B	522	LYS	2.8
1	B	404	GLU	2.8
1	B	577	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	292	ARG	2.6
1	B	596	ALA	2.5
1	B	293	GLY	2.5
1	A	641	ASN	2.5
1	A	404	GLU	2.5
1	B	567	GLN	2.5
1	B	524	ILE	2.5
1	A	603	ALA	2.4
1	B	588	ASP	2.3
1	B	568	ASN	2.2
1	B	306	ILE	2.2
1	B	346	GLU	2.2
1	B	317	GLU	2.2
1	B	580	ARG	2.1
1	A	351	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.