



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 06:07 am BST

PDB ID : 2H65  
Title : Crystal structure of caspase-3 with inhibitor Ac-VDVAD-Cho  
Authors : Fang, B.; Boross, P.I.; Tozser, J.; Weber, I.T.  
Deposited on : 2006-05-30  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	
1	C	146	
2	B	95	
2	D	95	
3	E	6	
3	F	6	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3938 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called caspase-3, p17 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	140	1112	688	201	214	9	0	0	0
1	C	141	1120	692	202	217	9	0	0	0

- Molecule 2 is a protein called caspase-3, p12 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	93	777	511	125	136	5	0	0	0
2	D	92	767	505	122	135	5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	278	HIS	-	EXPRESSION TAG	UNP P42574
D	278	HIS	-	EXPRESSION TAG	UNP P42574

- Molecule 3 is a protein called Ac-VDVAD-Cho.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	6	38	23	5	10	0	0	0
3	F	6	38	23	5	10	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	31	Total	O	0	0
			31	31		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	B	17	Total O 17 17	0	0
4	C	15	Total O 15 15	0	0
4	D	22	Total O 22 22	0	0
4	F	1	Total O 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: caspase-3, p17 subunit




- Molecule 1: caspase-3, p17 subunit




- Molecule 2: caspase-3, p12 subunit



- Molecule 2: caspase-3, p12 subunit



- Molecule 3: Ac-VDVAD-Cho





- Molecule 3: Ac-VDVAD-Cho



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.89Å 67.40Å 93.49Å 90.00° 100.80° 90.00°	Depositor
Resolution (Å)	10.00 – 2.30 37.95 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.0 (10.00-2.30) 91.5 (37.95-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.82 (at 2.10Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.209 , 0.246 0.206 , 0.239	Depositor DCC
$R_{free}$ test set	1329 reflections (3.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.9	Xtrriage
Anisotropy	0.609	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 42.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3938	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASJ, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/1127	0.62	0/1509
1	C	0.33	0/1135	0.58	0/1520
2	B	0.44	0/802	0.64	0/1083
2	D	0.42	0/791	0.61	0/1068
3	E	0.49	0/27	0.49	0/37
3	F	0.53	0/27	0.60	0/37
All	All	0.39	0/3909	0.61	0/5254

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1112	0	1109	21	0
1	C	1120	0	1113	27	0
2	B	777	0	754	21	0
2	D	767	0	747	14	0
3	E	38	0	35	1	0
3	F	38	0	35	2	0
4	A	31	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	17	0	0	0	0
4	C	15	0	0	0	0
4	D	22	0	0	2	0
4	F	1	0	0	0	0
All	All	3938	0	3793	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:MET:HG3	1:A:139:ILE:HG23	1.64	0.79
1:A:93:ARG:HB2	1:A:134:VAL:HG23	1.67	0.77
2:B:222:MET:CE	2:B:242:LYS:HD2	2.16	0.73
2:B:222:MET:HE1	2:B:242:LYS:HD2	1.70	0.72
1:C:119:LEU:HD23	1:C:161:GLN:HB3	1.72	0.71
1:A:50:ILE:HG23	1:A:91:LEU:HD12	1.73	0.70
1:A:42:PRO:HG2	1:A:110:LYS:HE2	1.76	0.68
1:C:100:MET:HG3	1:C:139:ILE:HG23	1.77	0.65
1:A:93:ARG:HB2	1:A:134:VAL:CG2	2.26	0.65
1:C:93:ARG:HB2	1:C:134:VAL:CG2	2.28	0.63
1:C:140:THR:HG21	2:D:195:TYR:HE2	1.64	0.63
2:D:271:LYS:HE2	4:D:382:HOH:O	1.98	0.62
1:A:74:LEU:HD13	1:A:117:VAL:HG11	1.84	0.59
2:D:207:ARG:HA	2:D:213:SER:HA	1.84	0.59
1:C:88:LYS:HB3	1:C:91:LEU:HD21	1.83	0.58
1:A:159:ILE:HD13	2:B:219:LEU:HD21	1.85	0.58
1:C:128:PHE:CE1	1:C:133:PRO:HG3	2.39	0.58
1:A:113:SER:HB3	1:A:155:PRO:HG2	1.86	0.57
1:A:57:LYS:CE	1:A:57:LYS:H	2.20	0.55
1:A:137:LYS:HA	1:A:137:LYS:HE2	1.88	0.55
2:B:207:ARG:HA	2:B:213:SER:HA	1.88	0.55
1:C:74:LEU:HD13	1:C:117:VAL:HG11	1.88	0.55
1:C:159:ILE:HD13	2:D:219:LEU:HD21	1.88	0.55
1:A:74:LEU:CD1	1:A:117:VAL:HG11	2.36	0.54
2:B:194:LEU:HD13	2:B:194:LEU:C	2.28	0.54
2:B:222:MET:HE2	2:B:242:LYS:HD2	1.90	0.54
2:B:194:LEU:HD13	2:B:195:TYR:N	2.23	0.53
1:C:53:LYS:HD3	1:C:68:ASP:OD2	2.09	0.52
1:C:141:ASN:HA	1:C:144:ARG:HG3	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:GLU:H	1:A:173:GLU:CD	2.13	0.52
1:C:99:LEU:O	1:C:103:VAL:HG23	2.10	0.52
1:A:57:LYS:H	1:A:57:LYS:HE2	1.75	0.51
2:D:231:GLU:OE1	2:D:233:MET:HB2	2.09	0.51
1:C:74:LEU:CD1	1:C:117:VAL:HG11	2.41	0.51
2:B:206:TRP:HH2	2:B:256:PHE:HB3	1.76	0.50
2:B:231:GLU:OE1	2:B:233:MET:HB2	2.13	0.49
2:D:210:LYS:HG3	2:D:211:ASP:N	2.28	0.49
1:A:37:TYR:CD1	1:A:155:PRO:HD3	2.47	0.49
1:C:149:ARG:HB3	1:C:149:ARG:NH1	2.28	0.49
1:C:91:LEU:HD12	1:C:95:GLU:HB3	1.94	0.49
2:D:240:ASN:OD1	2:D:263:PRO:HB2	2.13	0.48
1:A:151:LEU:O	1:A:154:LYS:HB2	2.14	0.48
2:D:210:LYS:HG3	2:D:211:ASP:OD2	2.14	0.48
1:A:135:ASP:OD2	1:A:138:LYS:HE2	2.13	0.47
1:C:163:CYS:N	3:F:6:ASJ:H	2.30	0.47
1:A:119:LEU:HD23	1:A:161:GLN:HB3	1.95	0.47
2:B:240:ASN:OD1	2:B:263:PRO:HB2	2.14	0.47
2:D:206:TRP:HH2	2:D:256:PHE:HB3	1.79	0.47
1:A:135:ASP:N	1:A:135:ASP:OD2	2.47	0.46
2:B:213:SER:OG	2:B:216:ILE:HG12	2.16	0.46
2:B:224:LYS:HZ2	2:B:224:LYS:HB2	1.82	0.45
2:D:224:LYS:HE2	2:D:224:LYS:HB2	1.75	0.45
1:A:168:LEU:HD22	2:B:259:LYS:HG3	1.99	0.44
2:B:194:LEU:C	2:B:194:LEU:CD1	2.86	0.44
2:B:266:VAL:HG12	2:B:268:MET:HE1	2.00	0.44
1:C:88:LYS:HB3	1:C:91:LEU:CD2	2.48	0.43
1:A:57:LYS:HD3	1:A:57:LYS:N	2.33	0.43
2:B:277:HIS:O	2:B:278:HIS:C	2.56	0.43
1:C:93:ARG:HB2	1:C:134:VAL:HG23	1.98	0.43
1:C:163:CYS:H	3:F:6:ASJ:H	1.84	0.43
1:C:39:MET:HG3	2:D:274:TYR:O	2.18	0.43
1:C:120:SER:HB3	1:C:127:ILE:HD11	1.99	0.42
1:C:37:TYR:CD2	1:C:155:PRO:HG3	2.54	0.42
2:B:191:ALA:O	2:B:192:ASP:HB2	2.19	0.42
1:C:93:ARG:HG2	1:C:94:GLU:OE1	2.20	0.42
2:D:213:SER:OG	2:D:216:ILE:HG12	2.20	0.42
2:D:271:LYS:CE	4:D:382:HOH:O	2.64	0.41
1:A:93:ARG:HG3	1:A:94:GLU:OE2	2.20	0.41
2:B:224:LYS:NZ	2:B:224:LYS:HB2	2.35	0.41
1:C:92:THR:OG1	1:C:95:GLU:HG3	2.19	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:268:MET:HB2	2:B:268:MET:HE3	1.77	0.41
2:B:204:TYR:CD2	3:E:5:ALA:HB1	2.56	0.41
1:C:42:PRO:HG2	1:C:110:LYS:HE2	2.03	0.41
1:C:56:HIS:HD2	1:C:57:LYS:H	1.69	0.41
1:C:56:HIS:HD2	1:C:57:LYS:N	2.19	0.40
2:B:270:THR:O	2:D:241:ARG:HB2	2.21	0.40
1:C:103:VAL:O	1:C:106:GLU:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/146 (94%)	131 (95%)	6 (4%)	1 (1%)	22	26
1	C	139/146 (95%)	129 (93%)	10 (7%)	0	100	100
2	B	91/95 (96%)	87 (96%)	3 (3%)	1 (1%)	14	15
2	D	90/95 (95%)	87 (97%)	3 (3%)	0	100	100
3	E	4/6 (67%)	4 (100%)	0	0	100	100
3	F	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
All	All	466/494 (94%)	441 (95%)	23 (5%)	2 (0%)	34	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	ASP
2	B	229	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/131 (96%)	121 (96%)	5 (4%)	31	44
1	C	127/131 (97%)	123 (97%)	4 (3%)	40	55
2	B	83/85 (98%)	79 (95%)	4 (5%)	25	36
2	D	82/85 (96%)	79 (96%)	3 (4%)	34	48
3	E	3/3 (100%)	3 (100%)	0	100	100
3	F	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	424/438 (97%)	407 (96%)	17 (4%)	31	44

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LYS
1	A	61	MET
1	A	124	GLU
1	A	137	LYS
1	A	173	GLU
2	B	222	MET
2	B	268	MET
2	B	273	LEU
2	B	278	HIS
1	C	34	ASP
1	C	102	ASP
1	C	147	ARG
1	C	157	LEU
2	D	190	GLU
2	D	209	SER
2	D	273	LEU
3	F	2	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	A	89	ASN
2	B	217	GLN
2	B	278	HIS
1	C	56	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ASJ	E	6	1,3	4,7,7	0.64	0	3,8,8	1.06	0
3	ASJ	F	6	1,3	4,7,7	0.54	0	3,8,8	1.31	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASJ	E	6	1,3	-	3/4/6/6	-
3	ASJ	F	6	1,3	-	4/4/6/6	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	6	ASJ	CB-CA-C	-2.24	108.15	112.21

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	6	ASJ	C-CA-CB-CG
3	F	6	ASJ	O-C-CA-N
3	F	6	ASJ	O-C-CA-CB
3	F	6	ASJ	C-CA-CB-CG
3	F	6	ASJ	N-CA-CB-CG
3	E	6	ASJ	N-CA-CB-CG
3	E	6	ASJ	O-C-CA-N

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	6	ASJ	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	140/146 (95%)	0.00	1 (0%) 87 91	12, 28, 42, 55	2 (1%)
1	C	141/146 (96%)	0.44	8 (5%) 23 30	16, 38, 59, 75	0
2	B	93/95 (97%)	0.12	2 (2%) 62 69	10, 20, 35, 62	0
2	D	92/95 (96%)	0.19	2 (2%) 62 69	13, 25, 42, 66	0
3	E	4/6 (66%)	-0.14	0 100 100	26, 29, 31, 31	0
3	F	4/6 (66%)	0.23	0 100 100	39, 40, 43, 43	0
All	All	474/494 (95%)	0.19	13 (2%) 54 62	10, 28, 51, 75	2 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	60	GLY	4.5
1	C	59	THR	4.2
1	C	57	LYS	3.7
1	C	61	MET	3.7
2	B	277	HIS	3.6
2	D	277	HIS	2.7
1	C	56	HIS	2.6
1	C	58	SER	2.5
1	C	135	ASP	2.3
2	D	268	MET	2.3
1	C	134	VAL	2.1
2	B	278	HIS	2.0
1	A	57	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ASJ	E	6	8/8	0.85	0.16	30,33,36,38	0
3	ASJ	F	6	8/8	0.90	0.13	40,44,46,48	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.