

Full wwPDB X-ray Structure Validation Report (i)

Aug 19, 2023 – 06:22 PM EDT

PDB ID : 2H6I

Title: W102T/Y365F Protein Farnesyltransferase Double Mutant Complexed with a

Geranylgeranylated DDPTASACVLS Peptide Product at 3.0A

Authors : Terry, K.L.; Beese, L.S.

Deposited on : 2006-05-31

Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	382	6	4%	18%	18%	
2	В	437		75%		19% 6%	
3	Р	11	27% 9	%	64%		
4	С	2		100%			



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 6019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein farnesyltransferase/geranylgeranyltransferase type I alpha subunit.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	315	Total 2683	C 1711	N 467	O 500	S 5	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	380	GLU	-	cloning artifact	UNP P49354
A	381	GLU	-	cloning artifact	UNP P49354
A	382	PHE	-	cloning artifact	UNP P49354

• Molecule 2 is a protein called Protein farnesyltransferase beta subunit.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	410	Total 3220	C 2058	N 551	O 589	S 22	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	602	THR	TRP	engineered mutation	UNP P49356
В	865	PHE	TYR	engineered mutation	UNP P49356

• Molecule 3 is a protein called farnesylated peptide.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	Р	4	Total 28	C 17	N 4	O 6	S 1	0	0	0

• Molecule 4 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.






Mol	Chain	Residues	At	oms		ZeroOcc	AltConf	Trace
4	С	2	Total 23	C 12	O 11	0	0	0

• Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Zn 1 1	0	0

 \bullet Molecule 6 is GERAN-8-YL GERAN (three-letter code: GER) (formula: $\mathrm{C}_{20}\mathrm{H}_{34}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	Р	1	Total C 20 20	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	17	Total O 17 17	0	0
7	В	24	Total O 24 24	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	Р	3	Total O 3 3	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein farnesyltransferase/geranylgeranyltransferase type I alpha subunit



• Molecule 2: Protein farnesyltransferase beta subunit



• Molecule 4: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



Chain C: 100%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants	178.13Å 178.13Å 64.90Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.81 - 3.00	Depositor
Resolution (A)	49.66 - 2.59	EDS
% Data completeness	89.9 (49.81-3.00)	Depositor
(in resolution range)	74.1 (49.66-2.59)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$< I/\sigma(I) > 1$	1.53 (at 2.58Å)	Xtriage
Refinement program	CNS	Depositor
D D	0.270 , 0.281	Depositor
R, R_{free}	0.286 , 0.296	DCC
R_{free} test set	1028 reflections (3.78%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35 , 4.1	EDS
L-test for twinning ²	$< L >=0.37, < L^2>=0.19$	Xtriage
Estimated twinning fraction	0.106 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.80	EDS
Total number of atoms	6019	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.81% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FRU, GLC, GER

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain Bond lengths		Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.54	0/2750	0.63	0/3735
2	В	0.60	0/3307	0.69	0/4493
3	Р	1.15	0/27	1.10	0/34
All	All	0.58	0/6084	0.66	0/8262

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2683	0	2601	41	1
2	В	3220	0	3144	56	1
3	Р	28	0	28	1	0
4	С	23	0	21	0	0
5	В	1	0	0	0	0
6	Р	20	0	32	5	0
7	A	17	0	0	2	0
7	В	24	0	0	3	0
7	Р	3	0	0	1	0
All	All	6019	0	5826	97	1



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 8.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
9.D.766. ADC.IIII01	9.D.010.CI N.H.O	1.34	overlap (Å)
2:B:766:ARG:HH21	2:B:818:GLN:HG2 7:P:1697:HOH:O		0.93
3:P:2008:LEU:HB2		1.76	0.85
2:B:602:THR:HG23	6:P:2010:GER:H201	1.66	0.76
1:A:335:ASN:O	1:A:339:GLU:HG3	1.87	0.74
2:B:764:ARG:NH1	7:B:1373:HOH:O	2.23	0.72
2:B:651:ALA:HB3	2:B:652:PRO:CD	2.22	0.70
2:B:577:LYS:HD3	2:B:846:PRO:O	1.92	0.69
2:B:766:ARG:NH2	2:B:818:GLN:HG2	2.06	0.67
1:A:316:VAL:O	1:A:320:GLU:HG3	1.94	0.67
2:B:702:ARG:HB3	6:P:2010:GER:H142	1.80	0.63
2:B:870:LEU:O	2:B:874:GLN:HG3	1.97	0.63
2:B:624:THR:O	2:B:628:GLN:HG3	1.99	0.63
2:B:739:ILE:HB	2:B:752:THR:HA	1.84	0.60
1:A:185:GLU:O	1:A:189:ILE:HG13	2.03	0.58
2:B:627:CYS:HB3	2:B:671:ILE:HD11	1.84	0.58
2:B:753:PHE:HA	2:B:807:LEU:HD21	1.85	0.58
1:A:263:THR:O	1:A:267:ILE:HG13	2.06	0.56
1:A:96:PHE:HA	1:A:126:LEU:HD13	1.88	0.56
1:A:237:TRP:CD1	1:A:275:SER:HB3	2.42	0.55
2:B:584:ARG:NH2	2:B:588:GLN:OE1	2.40	0.55
2:B:914:TYR:O	2:B:917:GLN:HB2	2.06	0.55
2:B:602:THR:HG23	6:P:2010:GER:C20	2.36	0.54
2:B:651:ALA:HB3	2:B:652:PRO:HD3	1.88	0.54
1:A:87:VAL:HG23	1:A:88:VAL:HG22	1.89	0.54
1:A:308:SER:HB3	7:A:1675:HOH:O	2.07	0.54
2:B:553:GLU:HB3	7:B:1589:HOH:O	2.07	0.54
1:A:266:MET:HA	1:A:266:MET:HE2	1.89	0.53
2:B:635:SER:OG	2:B:637:GLU:HG2	2.08	0.53
2:B:806:GLY:HA2	2:B:872:ILE:HD13	1.91	0.53
1:A:151:GLU:HG3	1:A:175:LEU:HD11	1.91	0.53
1:A:353:LYS:O	1:A:357:ARG:HG2	2.09	0.52
2:B:702:ARG:HD2	6:P:2010:GER:H142	1.92	0.52
2:B:837:GLU:HG3	2:B:887:LEU:HD21	1.90	0.52
1:A:156:THR:HG23	1:A:188:PHE:CE1	2.43	0.52
1:A:298:GLN:O	1:A:302:LEU:HD23	2.11	0.51
1:A:77:PRO:HB3	1:A:102:TYR:CE1	2.46	0.51
2:B:801:SER:O	2:B:805:ALA:HB3	2.11	0.50
	2.0001211.1120		1 0.00



 $Continued\ from\ previous\ page...$

Continuea from previ		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:303:GLN:N	1:A:304:PRO:HD2	2.25	0.50
2:B:577:LYS:NZ	7:B:1822:HOH:O	2.45	0.50
2:B:831:HIS:HB3	2:B:834:ALA:HB3	1.93	0.50
1:A:58:LEU:HD11	1:A:126:LEU:HG	1.93	0.50
1:A:109:ARG:HH11	1:A:109:ARG:CB	2.24	0.50
2:B:649:HIS:O	2:B:652:PRO:HG2	2.12	0.50
2:B:685:LYS:HD3	2:B:691:PHE:CZ	2.47	0.50
2:B:601:PRO:O	2:B:604:CYS:HB2	2.12	0.50
1:A:233:ASN:CG	1:A:236:VAL:HG23	2.32	0.49
2:B:881:MET:C	2:B:882:LEU:HD12	2.32	0.49
2:B:692:LEU:HD23	2:B:699:VAL:CG2	2.43	0.49
2:B:756:LEU:O	2:B:760:VAL:HG23	2.12	0.49
1:A:132:THR:HG23	2:B:647:TYR:CD1	2.47	0.48
2:B:658:ASN:O	2:B:662:ILE:HG13	2.13	0.48
2:B:837:GLU:HG3	2:B:887:LEU:CD2	2.44	0.47
1:A:138:ARG:HA	1:A:141:LEU:HD12	1.97	0.47
2:B:808:LEU:HB2	2:B:809:PRO:HD3	1.97	0.47
1:A:355:TYR:O	1:A:358:TYR:HB3	2.15	0.46
1:A:148:LEU:HB2	1:A:179:LEU:HD21	1.97	0.46
2:B:806:GLY:O	2:B:809:PRO:HD2	2.15	0.46
2:B:908:VAL:O	2:B:912:THR:HG23	2.16	0.46
1:A:349:ASP:OD1	1:A:352:ARG:NH2	2.48	0.46
2:B:832:GLN:O	2:B:836:GLN:HG3	2.16	0.46
2:B:763:LYS:HB2	2:B:763:LYS:NZ	2.30	0.45
1:A:344:LEU:HD13	1:A:356:TRP:CZ2	2.51	0.45
1:A:109:ARG:HB3	1:A:109:ARG:NH1	2.32	0.45
2:B:649:HIS:HB3	2:B:652:PRO:CG	2.47	0.45
2:B:729:ILE:O	2:B:732:CYS:HB2	2.17	0.45
2:B:835:LEU:HD23	2:B:873:ALA:HB2	1.98	0.45
2:B:534:ARG:HD3	2:B:784:PHE:CE1	2.52	0.45
1:A:328:ASP:O	1:A:329:ASN:HB2	2.16	0.45
2:B:797:ASP:HB3	2:B:800:TYR:HD1	1.82	0.44
2:B:534:ARG:HD3	2:B:784:PHE:CZ	2.53	0.43
1:A:221:GLN:NE2	7:A:1112:HOH:O	2.50	0.43
1:A:265:GLU:HA	1:A:265:GLU:OE2	2.18	0.43
2:B:678:LEU:HD13	2:B:716:ILE:O	2.19	0.43
2:B:852:ASP:HB3	2:B:856:LYS:HG3	2.00	0.43
2:B:898:HIS:CE1	2:B:900:VAL:HB	2.53	0.43
1:A:84:PRO:C	1:A:86:PRO:HD3	2.39	0.43
2:B:850:LEU:HB2	2:B:863:THR:HA	2.00	0.43
2:B:635:SER:OG	2:B:636:PRO:HD2	2.19	0.43



Continued from previous page...

Atom 1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
2:B:808:LEU:HD13	2:B:808:LEU:HA	1.88	0.42
6:P:2010:GER:H111	6:P:2010:GER:H91	1.94	0.42
1:A:112:ARG:NH1	1:A:143:SER:OG	2.52	0.42
1:A:219:GLU:O	1:A:223:VAL:HG23	2.19	0.42
1:A:256:LEU:O	1:A:260:VAL:HG23	2.20	0.42
1:A:198:LYS:NZ	2:B:793:ASN:O	2.53	0.42
1:A:231:VAL:HG22	1:A:231:VAL:O	2.20	0.42
1:A:274:GLU:HG2	1:A:278:ASN:ND2	2.34	0.41
1:A:116:ALA:O	1:A:119:LEU:HB3	2.19	0.41
1:A:109:ARG:CB	1:A:109:ARG:NH1	2.84	0.41
2:B:538:ASP:O	2:B:540:VAL:HG23	2.20	0.41
2:B:837:GLU:HB3	2:B:841:MET:CE	2.51	0.41
2:B:702:ARG:HA	2:B:754:CYS:SG	2.60	0.41
1:A:65:LEU:O	1:A:69:ARG:HG3	2.21	0.41
1:A:80:GLN:HB2	1:A:104:ARG:CZ	2.50	0.41
1:A:332:ASP:OD2	1:A:336:LYS:HE3	2.21	0.41
2:B:905:PRO:O	2:B:909:ILE:HG13	2.21	0.40
1:A:308:SER:O	1:A:312:ILE:HG12	2.21	0.40
2:B:681:LEU:HA	2:B:681:LEU:HD23	1.83	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:73:ALA:O	2:B:583:LYS:NZ[4_674]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	313/382 (82%)	294 (94%)	18 (6%)	1 (0%)	41 76



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	\mathbf{s}
2	В	408/437 (93%)	393 (96%)	14 (3%)	1 (0%)	47 82	
3	Р	2/11 (18%)	2 (100%)	0	0	100 100	
All	All	723/830 (87%)	689 (95%)	32 (4%)	2 (0%)	41 76	

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	217	ASP
2	В	921	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	A	294/344~(86%)	294 (100%)	0	100	100
2	В	346/370 (94%)	345 (100%)	1 (0%)	92	97
3	Р	4/9 (44%)	4 (100%)	0	100	100
All	All	644/723 (89%)	643 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	В	851	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	201	HIS
1	A	297	ASN
1	A	303	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Tiple	Link Bond lengths				Bond angles		
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	GLC	С	1	4	11,11,12	3.62	6 (54%)	15,15,17	1.58	2 (13%)	
4	FRU	С	2	4	11,12,12	1.58	2 (18%)	10,18,18	0.79	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	С	1	4	-	0/2/19/22	0/1/1/1
4	FRU	С	2	4	-	1/5/24/24	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
4	С	1	GLC	C2-C3	10.55	1.68	1.52
4	С	2	FRU	C1-C2	3.33	1.57	1.52
4	С	1	GLC	O5-C1	3.31	1.49	1.43
4	С	2	FRU	O2-C2	3.19	1.46	1.40
4	С	1	GLC	C4-C5	2.76	1.58	1.53
4	С	1	GLC	C4-C3	2.17	1.57	1.52



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$Ideal(\AA)$
4	С	1	GLC	C1-C2	2.11	1.57	1.52
4	С	1	GLC	O5-C5	2.08	1.47	1.43

All (2) bond angle outliers are listed below:

	Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
ſ	4	С	1	GLC	C1-O5-C5	3.88	117.44	112.19
	4	С	1	GLC	C1-C2-C3	-3.42	105.46	109.67

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	С	2	FRU	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain	Res	Link	Bond lengths			\mathbf{B}_{0}	ond ang	les
				nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	6	GER	Р	2010	3	19,19,19	0.80	0	22,22,22	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GER	Р	2010	3	-	7/20/20/20	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	Р	2010	GER	C3-C5-C6-C7
6	P	2010	GER	C11-C10-C8-C9
6	P	2010	GER	C11-C10-C8-C7
6	P	2010	GER	C14-C13-C15-C16
6	P	2010	GER	C13-C15-C16-C17
6	P	2010	GER	C12-C13-C15-C16
6	P	2010	GER	C10-C11-C12-C13


There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	P	2010	GER	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2	$OWAB(A^2)$	Q < 0.9
1	A	315/382 (82%)	-0.09	0 100 100	34, 42, 57, 71	0
2	В	410/437 (93%)	-0.15	2 (0%) 91 75	31, 37, 49, 58	0
3	Р	4/11 (36%)	-0.18	0 100 100	41, 45, 50, 53	0
All	All	729/830 (87%)	-0.12	2 (0%) 94 84	31, 39, 52, 71	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	В	921	PRO	2.2
2	В	878	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	GLC	С	1	11/12	0.84	0.31	46,47,47,48	0
4	FRU	С	2	12/12	0.91	0.32	43,45,46,46	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
6	GER	Р	2010	20/20	0.76	0.39	52,72,74,75	0
5	ZN	В	1001	1/1	0.98	0.10	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

