



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 01:11 am BST

PDB ID : 5H87  
Title : Crystal structure of mRojoA mutant - P63H - W143S  
Authors : Pandelieva, A.T.; Tremblay, V.; Sarvan, S.; Chica, R.A.; Couture, J.-F.  
Deposited on : 2015-12-23  
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	242	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">2%      79%      8% • 12%</p>
1	B	242	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">2%      79%      9% • 12%</p>
1	C	242	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 78%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">5%      78%      10% • 12%</p>
1	D	242	<div style="display: flex; align-items: center;"> <div style="width: 8%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 12%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">8%      79%      9% 12%</p>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7145 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called mRojoA fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	214	Total 1732	C 1103	N 290	O 331	S 8	0	2	0
1	B	214	Total 1732	C 1103	N 290	O 331	S 8	1	2	0
1	C	214	Total 1732	C 1103	N 290	O 331	S 8	2	2	0
1	D	214	Total 1732	C 1103	N 290	O 331	S 8	0	2	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	65	Total 65	O 65	0	0
2	B	46	Total 46	O 46	0	0
2	C	78	Total 78	O 78	0	0
2	D	28	Total 28	O 28	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.67Å 91.51Å 84.61Å 90.00° 108.72° 90.00°	Depositor
Resolution (Å)	32.80 – 2.24 30.80 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.3 (32.80-2.24) 99.8 (30.80-2.24)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.27 (at 2.24Å)	Xtrriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.181 , 0.234 0.191 , 0.244	Depositor DCC
$R_{free}$ test set	2134 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.8	Xtrriage
Anisotropy	0.452	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7145	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CH6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/1757	0.75	1/2363 (0.0%)
1	B	0.51	0/1757	0.74	0/2363
1	C	0.53	0/1757	0.75	0/2363
1	D	0.50	0/1757	0.71	0/2363
All	All	0.52	0/7028	0.74	1/9452 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	197	TYR	CB-CA-C	-5.86	98.68	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1732	0	1679	11	0
1	B	1732	0	1679	10	0
1	C	1732	0	1679	11	0
1	D	1732	0	1679	6	0
2	A	65	0	0	0	0
2	B	46	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	78	0	0	0	0
2	D	28	0	0	0	0
All	All	7145	0	6716	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:CH6:HD1	1:A:215:GLU:OE1	1.87	0.75
1:D:42:GLN:HE22	1:D:69:SER:HB3	1.60	0.66
1:A:127:THR:HB	1:C:94:GLU:HB3	1.83	0.60
1:A:42:GLN:HE22	1:A:69:SER:HB3	1.68	0.59
1:C:153:GLU:HB3	1:C:158:LYS:HD2	1.84	0.59
1:D:80:PRO:HD2	1:D:190:PRO:HB3	1.85	0.59
1:A:78:ASP:OD1	1:A:221:HIS:CE1	2.60	0.55
1:C:149:ARG:HH12	1:C:192:ALA:HB1	1.72	0.55
1:B:94:GLU:HG3	1:B:108:THR:HG22	1.89	0.55
1:B:99:PHE:CD2	1:B:175:ALA:HB2	2.43	0.54
1:D:28:GLU:HG3	1:D:50:LYS:HD3	1.90	0.54
1:D:41:THR:HG22	1:D:216:ARG:HG2	1.95	0.48
1:A:99:PHE:CD2	1:A:175:ALA:HB2	2.49	0.48
1:D:109:GLN:HG3	1:D:122:VAL:HG22	1.95	0.48
1:C:141:MET:SD	1:C:168:LYS:HA	2.54	0.48
1:B:53:PRO:HB3	1:B:207:ASP:HB2	1.97	0.47
1:C:59:ASP:HB3	1:C:165:LEU:HD22	1.96	0.47
1:C:157:LEU:HD13	1:C:190:PRO:HD2	1.97	0.47
1:C:41:THR:HG22	1:C:216:ARG:HG3	1.97	0.46
1:B:153:GLU:HB3	1:B:158:LYS:HG3	1.97	0.46
1:B:148:GLU:HG3	1:B:161:ILE:HG12	1.96	0.45
1:C:143:SER:HB3	1:C:165:LEU:HD23	1.97	0.45
1:B:66:CH6:HB12	1:B:215:GLU:OE1	2.17	0.45
1:C:42:GLN:HE22	1:C:69:SER:HB3	1.81	0.45
1:A:78:ASP:OD1	1:A:221:HIS:HE1	2.00	0.44
1:B:42:GLN:HE22	1:B:69:SER:HB3	1.83	0.44
1:B:149:ARG:HH12	1:B:192:ALA:HB1	1.84	0.43
1:A:171:GLY:HA2	1:B:100:GLU:OE2	2.19	0.43
1:B:145:ALA:HB2	1:B:198:LYS:HG3	2.02	0.42
1:C:148:GLU:HG3	1:C:161:ILE:HG12	2.02	0.42
1:D:99:PHE:CD2	1:D:175:ALA:HB2	2.56	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:GLN:HG3	1:A:124:LEU:HD12	2.02	0.41
1:A:70:LYS:HD3	1:A:197:TYR:OH	2.20	0.41
1:A:106:THR:OG1	1:A:125:HIS:HB2	2.21	0.40
1:A:66:CH6:HD2	1:A:161:ILE:HD13	2.03	0.40
1:C:109:GLN:HG3	1:C:122:VAL:HG22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/242 (87%)	209 (99%)	2 (1%)	0	100	100
1	B	211/242 (87%)	207 (98%)	4 (2%)	0	100	100
1	C	211/242 (87%)	208 (99%)	3 (1%)	0	100	100
1	D	211/242 (87%)	199 (94%)	10 (5%)	2 (1%)	17	13
All	All	844/968 (87%)	823 (98%)	19 (2%)	2 (0%)	47	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	115	ASP
1	D	169	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/204 (90%)	174 (95%)	10 (5%)	22	20
1	B	184/204 (90%)	177 (96%)	7 (4%)	33	36
1	C	184/204 (90%)	177 (96%)	7 (4%)	33	36
1	D	184/204 (90%)	175 (95%)	9 (5%)	25	25
All	All	736/816 (90%)	703 (96%)	33 (4%)	27	29

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ILE
1	A	26	GLU
1	A	89	GLU
1	A	125	HIS
1	A	127	THR
1	A	144	GLU
1	A	149	ARG
1	A	161	ILE
1	A	174	ASP
1	A	199	LEU
1	B	10	GLU
1	B	89	GLU
1	B	138	LYS
1	B	149	ARG
1	B	161	ILE
1	B	185	LYS
1	B	188	GLN
1	C	34	GLU
1	C	96	VAL
1	C	115	ASP
1	C	149	ARG
1	C	161	ILE
1	C	203	SER
1	C	215	GLU
1	D	89	GLU
1	D	114	GLN
1	D	132	ASP
1	D	147	SER
1	D	149	ARG
1	D	161	ILE
1	D	215	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	216	ARG
1	D	220	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	194	ASN
1	A	196	ASN
1	A	221	HIS
1	C	125	HIS
1	D	42	GLN
1	D	137	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CH6	A	66	1	24,24,25	0.39	0	28,32,34	0.92	2 (7%)
1	CH6	C	66	1	24,24,25	0.53	0	28,32,34	0.83	2 (7%)
1	CH6	B	66	1	24,24,25	0.48	0	28,32,34	0.82	2 (7%)
1	CH6	D	66	1	24,24,25	0.44	0	28,32,34	0.67	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CH6	A	66	1	-	0/12/31/32	0/2/2/2
1	CH6	C	66	1	-	1/12/31/32	0/2/2/2
1	CH6	B	66	1	-	1/12/31/32	0/2/2/2
1	CH6	D	66	1	-	2/12/31/32	0/2/2/2

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CH6	O3-C3-CA3	-2.83	117.85	126.39
1	C	66	CH6	O3-C3-CA3	-2.77	118.03	126.39
1	D	66	CH6	N3-C1-N2	2.67	113.31	111.45
1	A	66	CH6	N3-C1-N2	2.30	113.04	111.45
1	B	66	CH6	O3-C3-CA3	-2.23	119.66	126.39
1	C	66	CH6	N3-C1-N2	2.22	112.99	111.45
1	B	66	CH6	N3-C1-N2	2.11	112.91	111.45

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	66	CH6	CA1-CB1-CG1-SD
1	B	66	CH6	C1-CA1-CB1-CG1
1	D	66	CH6	C1-CA1-CB1-CG1
1	D	66	CH6	C3-CA3-N3-C2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	CH6	2	0
1	B	66	CH6	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	213/242 (88%)	0.09	6 (2%) 53 53	38, 53, 71, 91	0
1	B	213/242 (88%)	0.17	4 (1%) 66 68	36, 55, 75, 88	1 (0%)
1	C	213/242 (88%)	0.22	11 (5%) 27 26	38, 54, 75, 100	2 (0%)
1	D	213/242 (88%)	0.50	19 (8%) 9 9	38, 65, 90, 115	0
All	All	852/968 (88%)	0.24	40 (4%) 31 31	36, 56, 82, 115	3 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	168	LYS	6.4
1	C	169	ASP	4.8
1	C	115	ASP	4.4
1	D	197	TYR	4.2
1	C	107	VAL	3.7
1	D	172	HIS	3.4
1	A	169	ASP	3.2
1	C	149	ARG	3.0
1	D	149	ARG	3.0
1	C	222	SER	2.9
1	B	132	ASP	2.9
1	D	115	ASP	2.9
1	D	74	LYS	2.8
1	C	108	THR	2.8
1	B	222	SER	2.7
1	B	105	VAL	2.7
1	D	79	ILE	2.7
1	D	192	ALA	2.6
1	C	105	VAL	2.6
1	D	221	HIS	2.6
1	D	77	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	124	LEU	2.5
1	D	154	ASP	2.5
1	D	199	LEU	2.4
1	A	197	TYR	2.4
1	C	170	GLY	2.4
1	A	172	HIS	2.3
1	D	193	TYR	2.3
1	C	124	LEU	2.3
1	B	107	VAL	2.2
1	D	190	PRO	2.2
1	D	100	GLU	2.1
1	D	171	GLY	2.1
1	D	153	GLU	2.1
1	A	107	VAL	2.1
1	D	152	PRO	2.1
1	C	93	TRP	2.1
1	D	7	ILE	2.1
1	D	206	GLU	2.1
1	A	106	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CH6	B	66	23/24	0.87	0.23	67,78,98,99	0
1	CH6	D	66	23/24	0.89	0.26	75,85,105,110	0
1	CH6	A	66	23/24	0.90	0.23	62,73,97,97	0
1	CH6	C	66	23/24	0.94	0.20	58,68,86,87	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.