

# Full wwPDB X-ray Structure Validation Report (i)

#### May 21, 2024 – 03:44 AM EDT

PDB ID : 7H80

Title: Group deposition for crystallographic fragment screening of Chikungunya

virus nsP3 macrodomain – Crystal structure of Chikungunya virus nsP3

macrodomain in complex with Z57127349 (CHIKV\_MacB-x1112)

Authors: Aschenbrenner, J.C.; Fairhead, M.; Godoy, A.S.; Balcomb, B.H.; Capkin,

E.; Chandran, A.V.; Dolci, I.; Golding, M.; Koekemoer, L.; Lithgo, R.M.; Marples, P.G.; Ni, X.; Oliva, G.; Thompson, W.; Tomlinson, C.W.E.; Wild,

C.; Winokan, M.; Xavier, M.-A.E.; Fearon, D.; von Delft, F.

Deposited on : 2024-04-26

Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.2

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158


CCP4 : 7.0.044 (Gargrove)

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	٨	1.00	13%	_
1	А	163	98%	•
			21%	
1	В	163	91%	9%
			13%	
1	С	163	91%	9%

Continued on next page...

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36.2



Continued from previous page...

Mol	Chain	Length	Quality of chain	
			22%	
1	D	163	93%	• •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DMS	A	208	-	-	-	X



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Non-structural protein 3.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	163	Total	С	N	О	S	0	3	0
1	A	105	1270	791	221	248	10	0	3	0
1	В	163	Total	С	N	О	S	0	2	0
1	Ъ	105	1263	787	220	247	9	0		
1	С	163	Total	С	N	О	S	0	3	0
1		105	1274	794	225	246	9	0	3	0
1	D	159	Total	С	N	О	S	0	2	0
1	ש	109	1238	772	216	242	8	U		U

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q8JUX6
A	-1	ALA	-	expression tag	UNP Q8JUX6
A	0	MET	-	expression tag	UNP Q8JUX6
A	77	THR	SER	conflict	UNP Q8JUX6
В	-2	GLY	-	expression tag	UNP Q8JUX6
В	-1	ALA	-	expression tag	UNP Q8JUX6
В	0	MET	-	expression tag	UNP Q8JUX6
В	77	THR	SER	conflict	UNP Q8JUX6
С	-2	GLY	-	expression tag	UNP Q8JUX6
С	-1	ALA	-	expression tag	UNP Q8JUX6
С	0	MET	-	expression tag	UNP Q8JUX6
С	77	THR	SER	conflict	UNP Q8JUX6
D	-2	GLY	-	expression tag	UNP Q8JUX6
D	-1	ALA	-	expression tag	UNP Q8JUX6
D	0	MET	-	expression tag	UNP Q8JUX6
D	77	THR	SER	conflict	UNP Q8JUX6

• Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O S 4 2 1 1	0	0
2	A	1	Total C O S 4 2 1 1	0	0
2	A	1	Total C O S 4 2 1 1	0	0
2	A	1	Total C O S 4 2 1 1	0	0
2	С	1	Total C O S 4 2 1 1	0	0
2	С	1	Total C O S 4 2 1 1	0	0
2	D	1	Total C O S 4 2 1 1	0	0
2	D	1	Total C O S 4 2 1 1	0	0

 $\bullet$  Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1, 3-DIOL (three-letter code: TRS) (formula:  $\rm C_4H_{12}NO_3).$ 






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 8		N 1		0	0
3	D	1	Total 8	C 4	N 1	O 3	0	0

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	С	1	Total Cl 1 1	0	0

• Molecule 5 is succinimide (three-letter code: SZB) (formula:  $C_4H_3NO_2$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 7 4 1 2	0	0
5	A	1	Total C N O 7 4 1 2	0	0
5	В	1	Total C N O 7 4 1 2	0	0
5	В	1	Total C N O 7 4 1 2	0	0
5	С	1	Total C N O 7 4 1 2	0	0
5	С	1	Total C N O 7 4 1 2	0	0
5	D	1	Total C N O 7 4 1 2	0	0

#### • Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	165	Total O 165 165	0	0
6	В	99	Total O 99 99	0	0
6	С	134	Total O 134 134	0	0
6	D	129	Total O 129 129	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Non-structural protein 3





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants	87.23Å 87.23Å 85.16Å	D: t
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
Resolution (Å)	56.58 - 1.78	Depositor
Resolution (A)	56.51 - 1.78	EDS
% Data completeness	99.4 (56.58-1.78)	Depositor
(in resolution range)	99.4 (56.51-1.78)	EDS
$R_{merge}$	0.19	Depositor
$R_{sum}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.04 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC5	Depositor
υ .	0.193 , $0.244$	Depositor
$R, R_{free}$	0.230 , $0.266$	DCC
$R_{free}$ test set	3528 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.2	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.25 , 42.2	EDS
L-test for twinning <sup>2</sup>	$< L >=0.51, < L^2>=0.35$	Xtriage
	0.000 for -h,-k,l	
Estimated twinning fraction	0.042  for h,-h-k,-l	Xtriage
	0.003  for -k,-h,-l	
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5671	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 24.22 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.0012e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SZB, DMS, TRS, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.73	0/1293	0.82	0/1750	
1	В	0.68	0/1286	0.79	0/1741	
1	С	0.70	0/1297	0.80	0/1754	
1	D	0.71	0/1261	0.80	0/1707	
All	All	0.70	0/5137	0.80	0/6952	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1270	0	1259	2	0
1	В	1263	0	1253	10	0
1	С	1274	0	1272	7	0
1	D	1238	0	1228	5	0
2	A	16	0	24	2	0
2	С	8	0	12	1	0
2	D	8	0	12	0	0
3	A	8	0	12	0	0
3	D	8	0	12	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
4	A	1	0	0	0	0
4	С	1	0	0	1	0
5	A	14	0	0	0	0
5	В	14	0	0	0	0
5	С	14	0	0	0	0
5	D	7	0	0	0	0
6	A	165	0	0	3	0
6	В	99	0	0	1	0
6	С	134	0	0	2	0
6	D	129	0	0	0	0
All	All	5671	0	5084	27	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:201:DMS:H12	6:A:426:HOH:O	2.00	0.60
1:A:5:ARG:HD3	6:A:342:HOH:O	2.06	0.55
2:C:202:DMS:H22	6:C:348:HOH:O	2.07	0.53
1:B:121:LEU:C	1:B:121:LEU:HD23	2.29	0.52
1:B:143:CYS:O	1:B:144:ARG:HD2	2.10	0.51
1:D:2:PRO:HG3	1:D:133:ASP:HA	1.93	0.50
1:C:121:LEU:C	1:C:121:LEU:HD23	2.32	0.49
1:A:157:GLN:NE2	1:A:157:GLN:HA	2.28	0.48
1:B:9:MET:CE	1:B:13:LYS:O	2.63	0.47
1:B:103:SER:HA	1:B:138:ASP:O	2.16	0.46
1:C:38:TYR:HA	1:C:45:PHE:CE1	2.52	0.45
1:D:121:LEU:C	1:D:121:LEU:HD23	2.38	0.44
1:B:10:ASP:HB3	1:B:13:LYS:HD2	1.99	0.44
1:D:84:GLU:HA	1:D:84:GLU:OE1	2.18	0.44
1:B:13:LYS:HD3	4:C:201:CL:CL	2.55	0.44
1:B:95:GLU:HG2	6:B:386:HOH:O	2.17	0.44
1:C:140:VAL:HG12	1:C:142:TYR:CE1	2.54	0.43
1:C:159:ARG:HD3	6:C:407:HOH:O	2.19	0.42
1:B:11:ILE:HG21	1:B:33:VAL:HG13	2.01	0.42
1:B:8:ARG:CD	1:B:146:LYS:HG3	2.48	0.42
1:C:103:SER:HA	1:C:138:ASP:O	2.20	0.42
1:D:121:LEU:HD12	1:D:148:TRP:CD2	2.55	0.42
2:A:201:DMS:C1	6:A:426:HOH:O	2.63	0.41



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:C:73:PHE:CE1	1:C:81:GLY:HA3	2.56	0.41
1:B:9:MET:HE1	1:B:13:LYS:O	2.19	0.40
1:D:8:ARG:HA	1:D:143:CYS:O	2.22	0.40
1:C:21:ASN:OD1	1:C:34:CYS:HA	2.22	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	164/163 (101%)	163 (99%)	1 (1%)	0	100	100
1	В	163/163 (100%)	160 (98%)	3 (2%)	0	100	100
1	С	164/163 (101%)	161 (98%)	3 (2%)	0	100	100
1	D	159/163 (98%)	157 (99%)	2 (1%)	0	100	100
All	All	650/652 (100%)	641 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	137/134 (102%)	136 (99%)	1 (1%)	84 79		
1	В	136/134 (102%)	134 (98%)	2 (2%)	65 53		



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	137/134 (102%)	135 (98%)	2 (2%)	65	53
1	D	135/134 (101%)	135 (100%)	0	100	100
All	All	545/536 (102%)	540 (99%)	5 (1%)	78	72

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	LYS
1	В	31	ASP
1	В	153	SER
1	С	26	ARG
1	С	146	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	157	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	gles
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	DMS	С	202	-	3,3,3	0.30	0	3,3,3	0.13	0
5	SZB	A	206	_	7,7,7	3.02	1 (14%)	8,9,9	3.21	4 (50%)
5	SZB	В	202	-	7,7,7	2.99	1 (14%)	8,9,9	3.18	4 (50%)
2	DMS	A	208	-	3,3,3	0.23	0	3,3,3	0.12	0
3	TRS	A	202	-	7,7,7	0.30	0	9,9,9	0.47	0
2	DMS	D	202	-	3,3,3	0.48	0	3,3,3	0.27	0
3	TRS	D	201	-	7,7,7	0.30	0	9,9,9	0.40	0
5	SZB	С	204	-	7,7,7	3.08	1 (14%)	8,9,9	3.24	4 (50%)
2	DMS	A	203	-	3,3,3	0.29	0	3,3,3	0.10	0
5	SZB	A	207	-	7,7,7	3.06	1 (14%)	8,9,9	3.22	4 (50%)
5	SZB	D	204	-	7,7,7	3.10	1 (14%)	8,9,9	3.21	4 (50%)
2	DMS	D	203	-	3,3,3	0.30	0	3,3,3	0.08	0
2	DMS	A	204	-	3,3,3	0.29	0	3,3,3	0.09	0
5	SZB	В	201	-	7,7,7	3.02	1 (14%)	8,9,9	3.25	4 (50%)
2	DMS	С	205	-	3,3,3	0.16	0	3,3,3	0.14	0
2	DMS	A	201	-	3,3,3	0.41	0	3,3,3	0.31	0
5	SZB	С	203	-	7,7,7	3.08	1 (14%)	8,9,9	3.24	4 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SZB	A	206	-	-	-	0/1/1/1
5	SZB	В	202	-	-	-	0/1/1/1
3	TRS	D	201	-	-	6/9/9/9	-
5	SZB	С	204	-	-	-	0/1/1/1
5	SZB	A	207	-	-	-	0/1/1/1
5	SZB	D	204	-	-	-	0/1/1/1
5	SZB	В	201	-	-	-	0/1/1/1
5	SZB	С	203	-	-	-	0/1/1/1
3	TRS	A	202	-	-	4/9/9/9	-

All (7) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$Ideal(\AA)$
5	D	204	SZB	C1-C2	7.92	1.52	1.35
5	С	203	SZB	C1-C2	7.82	1.52	1.35
5	С	204	SZB	C1-C2	7.79	1.52	1.35
5	A	207	SZB	C1-C2	7.77	1.52	1.35
5	В	201	SZB	C1-C2	7.66	1.52	1.35
5	A	206	SZB	C1-C2	7.65	1.52	1.35
5	В	202	SZB	C1-C2	7.60	1.52	1.35

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	A	207	SZB	C2-C1-C	-5.52	104.85	108.61
5	С	204	SZB	C2-C1-C	-5.48	104.88	108.61
5	В	201	SZB	C2-C1-C	-5.47	104.89	108.61
5	С	203	SZB	C2-C1-C	-5.43	104.92	108.61
5	С	203	SZB	C1-C2-C3	-5.43	104.92	108.61
5	С	204	SZB	C1-C2-C3	-5.43	104.92	108.61
5	В	202	SZB	C1-C2-C3	-5.39	104.94	108.61
5	A	206	SZB	C1-C2-C3	-5.38	104.95	108.61
5	A	206	SZB	C2-C1-C	-5.37	104.96	108.61
5	В	201	SZB	C1-C2-C3	-5.37	104.96	108.61
5	A	207	SZB	C1-C2-C3	-5.36	104.96	108.61
5	D	204	SZB	C2-C1-C	-5.33	104.99	108.61
5	В	202	SZB	C2-C1-C	-5.31	105.00	108.61
5	D	204	SZB	C1-C2-C3	-5.31	105.00	108.61
5	С	203	SZB	C2-C3-N	3.19	108.29	106.05
5	В	201	SZB	C2-C3-N	3.16	108.27	106.05
5	В	201	SZB	C1-C-N	3.15	108.26	106.05
5	A	206	SZB	C2-C3-N	3.11	108.23	106.05
5	С	204	SZB	C1-C-N	3.10	108.23	106.05
5	С	204	SZB	C2-C3-N	3.10	108.22	106.05
5	A	206	SZB	C1-C-N	3.10	108.22	106.05
5	A	207	SZB	C1-C-N	3.08	108.22	106.05
5	D	204	SZB	C2-C3-N	3.07	108.21	106.05
5	С	203	SZB	C1-C-N	3.05	108.19	106.05
5	D	204	SZB	C1-C-N	3.05	108.19	106.05
5	В	202	SZB	C2-C3-N	3.04	108.19	106.05
5	A	207	SZB	C2-C3-N	3.01	108.16	106.05
5	В	202	SZB	C1-C-N	3.00	108.16	106.05

There are no chirality outliers.

All (10) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	202	TRS	N-C-C3-O3
3	A	202	TRS	C2-C-C3-O3
3	D	201	TRS	N-C-C1-O1
3	A	202	TRS	C1-C-C3-O3
3	D	201	TRS	C2-C-C1-O1
3	D	201	TRS	C3-C-C1-O1
3	D	201	TRS	C3-C-C2-O2
3	A	202	TRS	C3-C-C1-O1
3	D	201	TRS	N-C-C2-O2
3	D	201	TRS	C1-C-C3-O3


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	202	DMS	1	0
2	A	201	DMS	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2		$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	A	163/163 (100%)	0.23	22 (13%) 3	3	21, 28, 45, 53	23 (14%)
1	В	163/163 (100%)	0.76	34 (20%) 1	0	23, 34, 57, 72	34 (20%)
1	С	163/163 (100%)	0.32	22 (13%) 3	3	25, 34, 50, 70	29 (17%)
1	D	159/163 (97%)	0.69	36 (22%) 0	0	26, 33, 53, 77	34 (21%)
All	All	648/652 (99%)	0.50	114 (17%) 1	1	21, 32, 52, 77	120 (18%)

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106[A]	ILE	9.7
1	В	106[A]	ILE	8.4
1	С	106[A]	ILE	8.4
1	A	104	VAL	8.3
1	D	152	ILE	8.2
1	В	41	TRP	7.9
1	В	129	PHE	7.4
1	D	63	TYR	7.4
1	В	148	TRP	7.0
1	В	6	VAL	6.9
1	D	135	THR	6.9
1	D	143	CYS	6.7
1	A	103	SER	6.6
1	A	130	THR	6.2
1	В	158	MET	6.1
1	D	106[A]	ILE	6.0
1	D	62	THR	5.4
1	D	99	LEU	5.4
1	A	60	CYS	5.3
1	В	33	VAL	5.3
1	D	113	VAL	5.2



Continued from previous page...

Mol	nued fron Chain	$oxed{\mathbf{Res}}$	Type	RSRZ
1	В	60	CYS	5.2
1	В	42	PRO	5.0
1	В	130	THR	4.9
1	C	143	CYS	4.9
1	В	7	LYS	4.8
1	A	158	MET	4.8
1	В	113	VAL	4.8
1	В	61	GLY	4.8
1	D	3[A]	SER	4.7
1	В	111	THR	4.7
1	D	130	THR	4.7
1	В	150	LYS	4.7
1	D	103	SER	4.6
1	D	153	SER	4.5
1	D	111	THR	4.4
1	В	59	MET	4.4
1	D	144	ARG	4.3
1	A	113	VAL	4.3
1	В	110	SER	4.2
1	C	9	MET	4.2
1	D	150	LYS	4.2
1	D	136	ASP	4.2
1	D	59	MET	4.1
1	A	145[A]	ASP	4.1
1	В	112	GLY	4.0
1	В	16	GLU	4.0
1	A	112	GLY	4.0
1	В	147	GLU	4.0
1	В	44	SER	4.0
1	C	148	TRP	3.9
1	A	150	LYS	3.9
1	A	111	THR	3.9
1	D	90	ARG	3.8
1	A	160	THR	3.8
1	С	26	ARG	3.8
1	В	31	ASP	3.8
1	A	159	ARG	3.7
1	D	118	LYS	3.7
1	D	110	SER	3.7
1	D	147	GLU	3.7
1	A	157	GLN	3.6
1	В	91[A]	GLU	3.5
	1	L J	l .	



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Res	Type	RSRZ
1	D	112	GLY	3.5
1	С	158	MET	3.5
1	A	151	LYS	3.5
1	A	134	SER	3.4
1	A	62	THR	3.4
1	D	98	ARG	3.4
1	В	157	GLN	3.4
1	В	5	ARG	3.4
1	В	40	LYS	3.4
1	D	102	ASN	3.3
1	D	151	LYS	3.3
1	В	43	GLU	3.3
1	В	15	ASP	3.3
1	D	119	ASP	3.3
1	С	33	VAL	3.3
1	С	8	ARG	3.2
1	D	146	LYS	3.2
1	D	145	ASP	3.1
1	A	59[A]	MET	3.1
1	В	160	THR	3.1
1	В	159	ARG	3.0
1	С	111	THR	3.0
1	С	13[A]	LYS	3.0
1	D	149	GLU	3.0
1	D	91	GLU	3.0
1	В	149	GLU	2.8
1	D	79	SER	2.8
1	A	110	SER	2.8
1	D	117	GLY	2.8
1	В	13	LYS	2.7
1	В	83	ARG	2.7
1	D	160	THR	2.6
1	A	7	LYS	2.6
1	С	127	HIS	2.5
1	D	94	LYS	2.5
1	С	160	THR	2.5
1	C C C	90[A]	ARG	2.4
1	С	7	LYS	2.4
1	C	-2	GLY	2.4
1	C C	159	ARG	2.4
1		157	GLN	2.4
1	С	113	VAL	2.4



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	10	ASP	2.4
1	В	62	THR	2.4
1	A	144	ARG	2.4
1	D	158	MET	2.4
1	С	112	GLY	2.3
1	D	31	ASP	2.2
1	С	153	SER	2.1
1	С	144	ARG	2.1
1	С	146	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	TRS	D	201	8/8	0.57	0.28	50,63,65,70	0
5	SZB	A	207	7/7	0.68	0.28	28,29,30,30	7
2	DMS	С	202	4/4	0.73	0.30	56,59,65,70	0
2	DMS	A	208	4/4	0.78	0.49	115,116,117,120	4
5	SZB	В	202	7/7	0.79	0.17	30,31,31,31	7
2	DMS	D	203	4/4	0.86	0.29	74,83,92,93	0
5	SZB	D	204	7/7	0.86	0.17	17,18,18,18	7
3	TRS	A	202	8/8	0.87	0.13	44,55,59,64	0
2	DMS	A	203	4/4	0.89	0.42	64,75,77,79	0
5	SZB	С	203	7/7	0.89	0.14	26,27,27,28	7
2	DMS	С	205	4/4	0.89	0.17	64,64,65,65	4
5	SZB	С	204	7/7	0.90	0.11	26,29,30,30	7
4	CL	С	201	1/1	0.93	0.09	73,73,73,73	0
5	SZB	A	206	7/7	0.93	0.12	15,15,16,16	7




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	DMS	A	201	4/4	0.93	0.28	50,53,59,62	0
5	SZB	В	201	7/7	0.93	0.20	18,19,19,20	7
2	DMS	A	204	4/4	0.94	0.13	35,40,40,42	4
2	DMS	D	202	4/4	0.95	0.18	41,45,47,51	0
4	CL	A	205	1/1	0.95	0.16	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.












#### Electron density around SZB D 204:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around SZB C 203:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)
















## 6.5 Other polymers (i)

There are no such residues in this entry.

