

Full wwPDB X-ray Structure Validation Report (i)

May 20, 2024 – 08:32 PM EDT

PDB ID : 7H8Y

Title: Group deposition for crystallographic fragment screening of Chikungunya

virus nsP3 macrodomain – Crystal structure of Chikungunya virus nsP3

macrodomain in complex with Z57127903 (CHIKV_MacB-x1331)

Authors: Aschenbrenner, J.C.; Fairhead, M.; Godoy, A.S.; Balcomb, B.H.; Capkin,

E.; Chandran, A.V.; Dolci, I.; Golding, M.; Koekemoer, L.; Lithgo, R.M.; Marples, P.G.; Ni, X.; Oliva, G.; Thompson, W.; Tomlinson, C.W.E.; Wild,

C.; Winokan, M.; Xavier, M.-A.E.; Fearon, D.; von Delft, F.

Deposited on : 2024-04-26

Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.2

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158


CCP4 : 7.0.044 (Gargrove)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain	
1	A	163	93%	6% •
1	В	163	90%	7% ••
1	С	163	88%	11% •
1	D	163	94%	

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36.2



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Non-structural protein 3.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	163	Total	С	N	О	S	0	2	0
1	A	105	1262	787	220	245	10	0	<u> </u>	
1	В	159	Total	С	N	О	S	0	2	0
1	Ъ	159	1241	774	216	243	8	0	2	0
1	С	163	Total	С	N	О	S	0	3	0
1		105	1271	793	222	246	10	0	3	U
1	D	159	Total	С	N	О	S	0	2	0
1	ע	159	1238	772	216	242	8	0	<u> </u>	U

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q8JUX6
A	-1	ALA	-	expression tag	UNP Q8JUX6
A	0	MET	-	expression tag	UNP Q8JUX6
A	77	THR	SER	conflict	UNP Q8JUX6
В	-2	GLY	-	expression tag	UNP Q8JUX6
В	-1	ALA	-	expression tag	UNP Q8JUX6
В	0	MET	_	expression tag	UNP Q8JUX6
В	77	THR	SER	conflict	UNP Q8JUX6
С	-2	GLY	-	expression tag	UNP Q8JUX6
С	-1	ALA	-	expression tag	UNP Q8JUX6
С	0	MET	-	expression tag	UNP Q8JUX6
С	77	THR	SER	conflict	UNP Q8JUX6
D	-2	GLY	-	expression tag	UNP Q8JUX6
D	-1	ALA	-	expression tag	UNP Q8JUX6
D	0	MET	-	expression tag	UNP Q8JUX6
D	77	THR	SER	conflict	UNP Q8JUX6

• Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1		С	О	S	0	0
	7.1	1	4	2	1	1	0	<u> </u>
2	A	1		С	Ο	S	0	0
	11	1	4	2	1	1	Ü	Ü
2	A	1		С	Ο	S	0	0
	11	1	4	2	1	1	Ü	
2	A	1		С	Ο	S	0	0
	11	1	4	2	1	1	Ü	
2	В	1		С	Ο	S	0	0
		-	4	2	1	1	Ü	
2	В	1		С	Ο	S	0	0
		-	4	2	1	1	Ű	
2	С	1		С	Ο	S	0	0
		-	4	2	1	1	Ŭ	
2	D	1		С	Ο	\mathbf{S}	0	0
		-	4	2	1	1	Ü	
2	D	1		С	O	S	0	0
		_	4	2	1	1	J	
2	D	1		С	Ο	S	0	0
		1	4	2	1	1		<u> </u>

 \bullet Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	В	1	Total Cl 1 1	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	С	1	Total Cl 1 1	0	0
3	D	1	Total Cl 1 1	0	0

• Molecule 4 is ISATIN (three-letter code: ISN) (formula: $C_8H_5NO_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 11 8 1 2	0	0
4	В	1	Total C N O 11 8 1 2	0	0
4	В	1	Total C N O 11 8 1 2	0	0
4	С	1	Total C N O 11 8 1 2	0	0
4	D	1	Total C N O 11 8 1 2	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	80	Total O 80 80	0	0
5	В	52	Total O 52 52	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	С	58	Total O 58 58	0	0
5	D	50	Total O 50 50	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Non-structural protein 3





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants	87.46Å 87.46Å 84.60Å	D t
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	56.49 - 1.91	Depositor
Resolution (A)	56.43 - 1.91	EDS
% Data completeness	99.9 (56.49-1.91)	Depositor
(in resolution range)	99.9 (56.43-1.91)	EDS
R_{merge}	0.11	Depositor
R_{sum}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.02 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC5	Depositor
υ .	0.203 , 0.258	Depositor
R, R_{free}	0.291 , 0.309	DCC
R_{free} test set	2891 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.28, 60.3	EDS
L-test for twinning ²	$< L >=0.51, < L^2>=0.34$	Xtriage
	0.000 for -h,-k,l	
Estimated twinning fraction	0.032 for h,-h-k,-l	Xtriage
	0.000 for -k,-h,-l	
F_o, F_c correlation	0.90	EDS
Total number of atoms	5351	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 21.59 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.7956e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, CL, ISN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.73	0/1285	0.78	0/1739	
1	В	0.70	0/1264	0.80	0/1711	
1	С	0.74	0/1294	0.77	0/1750	
1	D	0.72	0/1261	0.80	0/1707	
All	All	0.72	0/5104	0.79	0/6907	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1262	0	1256	6	0
1	В	1241	0	1229	8	0
1	С	1271	0	1268	13	0
1	D	1238	0	1228	4	0
2	A	16	0	24	0	0
2	В	8	0	12	0	0
2	С	4	0	6	0	0
2	D	12	0	18	0	0
3	A	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	1	0	0	0	0
3	С	1	0	0	1	0
3	D	1	0	0	0	0
4	A	11	0	5	0	0
4	В	22	0	10	2	0
4	С	11	0	5	0	0
4	D	11	0	5	0	0
5	A	80	0	0	0	0
5	В	52	0	0	0	0
5	С	58	0	0	3	0
5	D	50	0	0	2	0
All	All	5351	0	5066	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:B:33:VAL:O	1:B:37:VAL:HG23	1.95	0.66
1:B:24:ASN:HB2	1:B:25:PRO:HD2	1.87	0.56
1:C:83:ARG:NE	5:C:303:HOH:O	2.40	0.55
1:C:114:TYR:HB2	3:C:202:CL:CL	2.46	0.52
1:C:122:THR:HB	5:C:333:HOH:O	2.09	0.52
1:B:60:CYS:SG	4:B:201:ISN:C7	2.97	0.52
1:C:121:LEU:C	1:C:121:LEU:HD23	2.30	0.51
1:A:43:GLU:OE1	1:C:83:ARG:NH2	2.41	0.51
1:A:21:ASN:OD1	1:A:34:CYS:HA	2.14	0.47
1:C:55:ALA:HA	1:C:67:HIS:O	2.15	0.46
1:D:90:ARG:HG3	5:D:335:HOH:O	2.14	0.46
1:C:2:PRO:HB2	1:D:26:ARG:HG2	1.98	0.46
1:B:128:LEU:C	1:B:128:LEU:HD23	2.37	0.45
1:C:76:TYR:HA	5:C:320:HOH:O	2.17	0.45
1:D:24:ASN:ND2	5:D:305:HOH:O	2.50	0.45
1:C:103:SER:HA	1:C:138:ASP:O	2.18	0.44
1:A:9:MET:CE	1:A:13:LYS:HB2	2.48	0.43
1:B:55:ALA:HA	1:B:67:HIS:O	2.19	0.43
1:D:33:VAL:O	1:D:37:VAL:HG23	2.19	0.43
1:B:24:ASN:HD21	1:B:28:LEU:HB2	1.84	0.42
1:B:145:ASP:OD2	1:B:147:GLU:HB2	2.18	0.42
1:A:87:ALA:O	1:A:91:GLU:HG2	2.19	0.42



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:128:LEU:C	1:A:128:LEU:HD23	2.40	0.42
1:C:12:ALA:HA	1:C:41:TRP:CH2	2.54	0.42
1:A:55:ALA:HA	1:A:67:HIS:O	2.20	0.41
1:C:50:THR:O	1:C:69:VAL:HG21	2.21	0.41
1:C:74:SER:C	1:C:75:ASN:HD22	2.25	0.40
1:B:60:CYS:HG	4:B:201:ISN:C7	2.35	0.40
1:C:121:LEU:HD23	1:C:121:LEU:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	163/163 (100%)	158 (97%)	5 (3%)	0	100 100
1	В	159/163~(98%)	154 (97%)	5 (3%)	0	100 100
1	С	164/163 (101%)	158 (96%)	6 (4%)	0	100 100
1	D	159/163~(98%)	156 (98%)	3 (2%)	0	100 100
All	All	$645/652 \ (99\%)$	626 (97%)	19 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	136/134 (102%)	134 (98%)	2 (2%)	65 61
1	В	135/134 (101%)	133 (98%)	2 (2%)	65 61
1	\mathbf{C}	137/134 (102%)	131 (96%)	6 (4%)	28 18
1	D	135/134 (101%)	134 (99%)	1 (1%)	84 83
All	All	543/536 (101%)	532 (98%)	11 (2%)	53 49

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	ARG
1	A	43	GLU
1	В	34	CYS
1	В	60	CYS
1	С	0	MET
1	С	26	ARG
1	С	35	LYS
1	С	75	ASN
1	С	83	ARG
1	С	150	LYS
1	D	39	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	47	ASN
1	С	75	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 19 ligands modelled in this entry, 4 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ISN	D	205	-	12,12,12	1.81	2 (16%)	17,17,17	1.86	3 (17%)
4	ISN	В	205	-	12,12,12	1.74	2 (16%)	17,17,17	1.85	3 (17%)
2	DMS	A	203	-	3,3,3	0.22	0	3,3,3	0.10	0
2	DMS	В	202	-	3,3,3	0.23	0	3,3,3	0.11	0
2	DMS	D	201	-	3,3,3	0.25	0	3,3,3	0.10	0
2	DMS	D	203	-	3,3,3	0.27	0	3,3,3	0.07	0
4	ISN	A	206	-	12,12,12	2.08	3 (25%)	17,17,17	2.37	5 (29%)
2	DMS	С	201	-	3,3,3	0.22	0	3,3,3	0.07	0
4	ISN	В	201	-	12,12,12	1.59	3 (25%)	17,17,17	2.15	5 (29%)
2	DMS	В	203	-	3,3,3	0.16	0	3,3,3	0.28	0
4	ISN	С	203	-	12,12,12	1.68	3 (25%)	17,17,17	1.95	4 (23%)
2	DMS	D	202	-	3,3,3	0.13	0	3,3,3	0.17	0
2	DMS	A	204	-	3,3,3	0.35	0	3,3,3	0.13	0
2	DMS	A	201	-	3,3,3	0.11	0	3,3,3	0.15	0
2	DMS	A	202	-	3,3,3	0.19	0	3,3,3	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ISN	D	205	-	-	-	0/2/2/2
4	ISN	В	205	-	-	-	0/2/2/2
4	ISN	A	206	-	-	-	0/2/2/2
4	ISN	В	201	-	-	-	0/2/2/2
4	ISN	С	203	-	-	-	0/2/2/2

All (13) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(A)
4	A	206	ISN	C10-N1	5.89	1.42	1.36
4	D	205	ISN	C10-N1	4.52	1.41	1.36
4	В	205	ISN	C10-N1	4.51	1.41	1.36
4	С	203	ISN	C10-N1	4.18	1.40	1.36
4	В	201	ISN	C10-N1	3.83	1.40	1.36
4	D	205	ISN	C10-C7	-3.66	1.51	1.56
4	В	205	ISN	C10-C7	-3.28	1.51	1.56
4	С	203	ISN	C10-C7	-2.88	1.52	1.56
4	В	201	ISN	C10-C7	-2.80	1.52	1.56
4	A	206	ISN	C10-C7	-2.78	1.52	1.56
4	A	206	ISN	C1-C7	-2.36	1.43	1.48
4	В	201	ISN	C1-C7	-2.28	1.43	1.48
4	С	203	ISN	C1-C7	-2.16	1.44	1.48

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
4	A	206	ISN	C6-C1-C7	6.21	111.11	107.30
4	D	205	ISN	O11-C10-C7	5.36	130.49	125.86
4	В	201	ISN	O11-C10-C7	5.34	130.47	125.86
4	В	201	ISN	C6-C1-C7	5.02	110.38	107.30
4	С	203	ISN	C6-C1-C7	4.88	110.29	107.30
4	В	205	ISN	O11-C10-C7	4.73	129.94	125.86
4	В	205	ISN	C6-C1-C7	4.58	110.11	107.30
4	С	203	ISN	O11-C10-C7	4.54	129.78	125.86
4	A	206	ISN	O11-C10-C7	4.44	129.69	125.86
4	D	205	ISN	C6-C1-C7	4.24	109.90	107.30
4	A	206	ISN	C1-C7-C10	-3.62	103.44	104.99
4	A	206	ISN	O8-C7-C10	3.46	126.65	123.73
4	В	201	ISN	O8-C7-C10	3.16	126.39	123.73
4	С	203	ISN	O8-C7-C10	2.90	126.17	123.73
4	В	201	ISN	C1-C7-C10	-2.50	103.92	104.99
4	С	203	ISN	C1-C7-C10	-2.21	104.05	104.99
4	В	201	ISN	O11-C10-N1	-2.20	122.86	126.36
4	A	206	ISN	C6-N1-C10	-2.12	110.04	111.38
4	D	205	ISN	O8-C7-C10	2.11	125.51	123.73
4	В	205	ISN	C1-C7-C10	-2.11	104.09	104.99

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 2 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	201	ISN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.












Electron density around ISN B 205:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ISN C 203:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

