



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 11:02 AM BST

PDB ID : 4HAQ
Title : Crystal Structure of a GH7 family cellobiohydrolase from *Limnoria quadripunctata* in complex with cellobiose and cellotriose
Authors : Martin, R.N.A.; McGeehan, J.E.; Streeter, S.D.; Cragg, S.M.; Guille, M.J.; Schnorr, K.M.; Kern, M.; Bruce, N.C.; McQueen-Mason, S.J.
Deposited on : 2012-09-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	431	
1	B	431	
2	C	2	
2	D	2	
3	E	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TRS	A	503	-	X	-	-

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 7207 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GH7 family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	431	3274	2011	529	707	27	0	5	0
1	B	431	3278	2014	529	708	27	0	6	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	PCA	-	expression tag	UNP D4HRL0
B	23	PCA	-	expression tag	UNP D4HRL0

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	23	12	11	0	0	0
2	D	2	23	12	11	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	E	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	Ca	0	0
			2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	282	Total	O	0	0
			282	282		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	282	Total 282	O 282	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GH7 family protein




- Molecule 1: GH7 family protein



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



- Molecule 3: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain E:

100%

GLA1
BGC2
BGC3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	52.39Å 54.27Å 72.67Å 94.39° 96.33° 94.99°	Depositor
Resolution (Å)	44.62 – 1.90 44.62 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.0 (44.62-1.90) 96.0 (44.62-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 1.89Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.165 , 0.217 0.165 , 0.216	Depositor DCC
R_{free} test set	3030 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	12.1	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.019 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7207	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BGC, CA, GLC, TRS, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	1/3356 (0.0%)	0.98	7/4568 (0.2%)
1	B	0.98	1/3363 (0.0%)	1.01	9/4578 (0.2%)
All	All	0.95	2/6719 (0.0%)	0.99	16/9146 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	304	SER	CB-OG	-5.23	1.35	1.42
1	A	246	TYR	CG-CD1	5.13	1.45	1.39

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	134	ASP	CB-CG-OD1	5.96	123.67	118.30
1	B	40	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	A	134	ASP	CB-CG-OD1	5.74	123.47	118.30
1	B	301	ASP	CB-CG-OD1	5.74	123.47	118.30
1	A	128	ARG	NE-CZ-NH1	-5.71	117.45	120.30
1	B	69	ASP	CB-CG-OD1	-5.61	113.25	118.30
1	B	348	ASP	CB-CG-OD1	5.60	123.34	118.30
1	B	69	ASP	CB-CG-OD2	5.36	123.12	118.30
1	A	219	ASP	CB-CG-OD1	5.32	123.08	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	353	ASP	CB-CG-OD1	5.25	123.02	118.30
1	A	136	ASP	CB-CG-OD1	-5.24	113.59	118.30
1	A	94	ASP	CB-CG-OD1	5.20	122.98	118.30
1	B	153	ASP	CB-CG-OD1	5.18	122.96	118.30
1	B	279	ASP	CB-CG-OD2	-5.16	113.65	118.30
1	B	393	ASP	CB-CG-OD1	5.14	122.92	118.30
1	A	153	ASP	CB-CG-OD1	5.12	122.91	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	452	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3274	0	2929	19	0
1	B	3278	0	2936	24	0
2	C	23	0	21	0	0
2	D	23	0	21	5	0
3	E	34	0	30	2	0
4	A	1	0	0	0	0
5	A	8	0	12	3	0
6	B	2	0	0	0	0
7	A	282	0	0	3	0
7	B	282	0	0	10	0
All	All	7207	0	5949	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:85:THR:HB	7:B:733:HOH:O	0.93	1.10
2:D:1:BGC:H6C2	2:D:2:BGC:O5	1.57	1.03
2:D:1:BGC:C6	2:D:2:BGC:O5	2.28	0.81
1:B:218:ASN:HD22	1:B:218:ASN:H	1.30	0.79
1:A:412:GLU:OE1	5:A:503:TRS:C	2.32	0.77
1:B:85:THR:CB	7:B:733:HOH:O	1.70	0.76
3:E:1:GLC:H62	3:E:2:BGC:O2	1.89	0.71
1:B:85:THR:HG23	7:B:686:HOH:O	1.91	0.70
1:B:82:ASP:OD1	1:B:207:LYS:HE2	1.92	0.69
1:A:110:LYS:HD3	7:A:859:HOH:O	1.92	0.68
1:B:85:THR:CG2	7:B:733:HOH:O	2.18	0.68
1:B:85:THR:CG2	7:B:686:HOH:O	2.43	0.66
1:B:218:ASN:ND2	1:B:218:ASN:H	1.97	0.63
1:B:218:ASN:HD22	1:B:218:ASN:N	1.95	0.62
1:A:227[B]:MET:SD	1:A:260:THR:CG2	2.89	0.61
7:B:871:HOH:O	2:D:2:BGC:C4	2.49	0.59
1:B:158:PRO:HD2	1:B:161:LEU:HD12	1.86	0.56
1:B:82:ASP:O	1:B:85:THR:HG22	2.07	0.54
1:B:293:PHE:CD1	1:B:336:ALA:HB2	2.47	0.49
1:A:352:GLU:O	1:A:356:VAL:HG23	2.13	0.49
3:E:2:BGC:O3	3:E:3:BGC:O5	2.23	0.49
2:D:1:BGC:H6C2	2:D:2:BGC:C5	2.42	0.49
1:A:227[B]:MET:SD	1:A:260:THR:HG22	2.54	0.48
1:A:363:GLY:O	5:A:503:TRS:H11	2.14	0.48
1:A:362:PHE:O	5:A:503:TRS:H12	2.14	0.48
1:B:120:GLU:HB2	7:B:856:HOH:O	2.13	0.48
1:B:334:GLN:HE21	1:B:453:ASP:C	2.17	0.47
1:B:23:PCA:HA	7:B:688:HOH:O	2.15	0.46
7:B:871:HOH:O	2:D:2:BGC:C3	2.61	0.45
1:A:205:ALA:HB2	1:A:258:PRO:HG3	1.99	0.45
1:A:434:PHE:N	1:A:435:PRO:CD	2.80	0.44
1:B:392:TRP:HE3	1:B:393:ASP:O	1.99	0.44
1:A:314:THR:HA	1:A:318:GLY:O	2.17	0.43
1:B:182:ASN:HB2	1:B:206:GLY:HA2	2.00	0.43
1:B:427:PRO:O	1:B:431:GLU:HG3	2.18	0.43
1:B:157:LEU:HD11	1:B:439:VAL:HB	1.99	0.43
1:B:227[B]:MET:SD	1:B:260:THR:CG2	3.07	0.43
1:B:160:GLY:HA3	1:B:425:GLY:HA2	2.01	0.42
1:B:204:ILE:O	1:B:227[A]:MET:HE1	2.19	0.42
1:A:158:PRO:HD2	1:A:161:LEU:HD12	2.01	0.42
1:A:51:ILE:HD11	1:A:104:ILE:HD12	2.00	0.42
1:A:63:ASP:HB2	1:A:64:THR:H	1.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:CYS:HB3	1:A:276:CYS:SG	2.59	0.42
1:B:29:THR:HG21	7:B:780:HOH:O	2.20	0.42
1:A:233:GLU:OE2	1:A:235:ASP:OD1	2.38	0.41
1:A:23:PCA:HA	7:A:652:HOH:O	2.20	0.41
1:A:30:GLU:CD	1:A:30:GLU:H	2.25	0.41
1:A:59:HIS:HE1	7:A:747:HOH:O	2.04	0.41
1:B:161:LEU:HD23	1:B:161:LEU:HA	1.79	0.40
1:A:151:ASP:HB2	1:A:442:SER:OG	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/431 (101%)	423 (98%)	9 (2%)	2 (0%)	29	18
1	B	435/431 (101%)	421 (97%)	14 (3%)	0	100	100
All	All	869/862 (101%)	844 (97%)	23 (3%)	2 (0%)	47	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	GLU
1	A	63	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/351 (101%)	344 (97%)	12 (3%)	37	28
1	B	357/351 (102%)	343 (96%)	14 (4%)	32	23
All	All	713/702 (102%)	687 (96%)	26 (4%)	37	26

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	GLU
1	A	44[A]	SER
1	A	44[B]	SER
1	A	54	ASN
1	A	62	GLU
1	A	63	ASP
1	A	64	THR
1	A	91	ASP
1	A	118	GLU
1	A	290	GLN
1	A	292	ASP
1	A	306	LEU
1	B	29	THR
1	B	54	ASN
1	B	64	THR
1	B	91	ASP
1	B	207	LYS
1	B	216	SER
1	B	218	ASN
1	B	262[A]	THR
1	B	262[B]	THR
1	B	290	GLN
1	B	306	LEU
1	B	421	SER
1	B	451	SER
1	B	453	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	156	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	290	GLN
1	A	433	ASN
1	B	156	ASN
1	B	218	ASN
1	B	290	GLN
1	B	334	GLN
1	B	433	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PCA	B	23	1	7,8,9	1.60	2 (28%)	9,10,12	2.26	3 (33%)
1	PCA	A	23	1	7,8,9	0.92	0	9,10,12	2.22	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	B	23	1	-	0/0/11/13	0/1/1/1
1	PCA	A	23	1	-	0/0/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	23	PCA	CA-N	-3.42	1.42	1.46
1	B	23	PCA	OE-CD	-2.24	1.18	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	PCA	CB-CA-C	-5.42	105.25	112.70
1	B	23	PCA	CB-CG-CD	-4.07	97.84	104.40
1	B	23	PCA	CB-CA-C	-3.99	107.21	112.70
1	A	23	PCA	OE-CD-CG	-2.91	121.69	126.76
1	B	23	PCA	O-C-CA	-2.36	118.58	124.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	23	PCA	1	0
1	A	23	PCA	1	0

5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	C	1	2	12,12,12	0.86	0	17,17,17	2.34	8 (47%)
2	BGC	C	2	2	11,11,12	0.91	0	15,15,17	3.45	6 (40%)
2	BGC	D	1	2	12,12,12	0.70	0	17,17,17	2.46	5 (29%)
2	BGC	D	2	2	11,11,12	1.07	0	15,15,17	4.22	8 (53%)
3	GLC	E	1	3	12,12,12	1.27	2 (16%)	17,17,17	2.26	7 (41%)
3	BGC	E	2	3	11,11,12	0.88	0	15,15,17	2.15	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BGC	E	3	3	11,11,12	0.67	0	15,15,17	1.87	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	2/2/22/22	0/1/1/1
2	BGC	C	2	2	-	2/2/19/22	0/1/1/1
2	BGC	D	1	2	-	1/2/22/22	0/1/1/1
2	BGC	D	2	2	-	2/2/19/22	0/1/1/1
3	GLC	E	1	3	-	0/2/22/22	0/1/1/1
3	BGC	E	2	3	-	2/2/19/22	0/1/1/1
3	BGC	E	3	3	-	1/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1	GLC	C1-C2	2.46	1.58	1.52
3	E	1	GLC	C3-C2	2.12	1.57	1.52

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	BGC	C1-O5-C5	-11.50	96.61	112.19
2	C	2	BGC	C1-O5-C5	-8.50	100.67	112.19
2	D	2	BGC	O5-C5-C6	7.34	118.71	107.20
2	D	1	BGC	O4-C4-C5	5.86	123.86	109.30
3	E	2	BGC	C1-C2-C3	5.82	116.82	109.67
2	C	2	BGC	C3-C4-C5	-5.34	100.72	110.24
3	E	3	BGC	C1-O5-C5	5.26	119.31	112.19
2	C	2	BGC	O5-C5-C6	5.20	115.36	107.20
2	D	1	BGC	C4-C3-C2	-4.94	102.20	110.82
3	E	1	GLC	O3-C3-C2	4.82	121.50	110.35
2	C	2	BGC	C2-C3-C4	-4.37	103.34	110.89
2	C	1	BGC	C1-O5-C5	-4.05	106.03	113.66
2	D	2	BGC	C3-C4-C5	-4.03	103.05	110.24
2	D	2	BGC	O4-C4-C5	-3.99	99.39	109.30
2	D	2	BGC	O5-C1-C2	-3.72	105.03	110.77
2	C	1	BGC	O4-C4-C5	-3.64	100.25	109.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	2	BGC	C3-C4-C5	-3.61	103.80	110.24
2	C	1	BGC	O1-C1-C2	3.43	118.69	109.03
2	D	2	BGC	O6-C6-C5	-3.42	99.55	111.29
2	C	1	BGC	O5-C5-C6	3.35	114.78	106.44
2	C	1	BGC	O6-C6-C5	-3.34	99.82	111.29
2	C	2	BGC	O6-C6-C5	-3.33	99.87	111.29
2	D	2	BGC	C2-C3-C4	-3.26	105.25	110.89
3	E	1	GLC	O1-C1-C2	3.26	118.20	109.03
3	E	1	GLC	O4-C4-C3	-3.24	102.85	110.35
2	D	1	BGC	O3-C3-C2	3.18	117.70	110.35
3	E	1	GLC	C3-C4-C5	-2.94	104.99	110.24
3	E	3	BGC	C2-C3-C4	-2.73	106.17	110.89
2	C	2	BGC	O4-C4-C5	2.66	115.90	109.30
2	C	1	BGC	O3-C3-C4	2.65	116.48	110.35
3	E	1	GLC	O5-C5-C4	2.63	114.47	109.69
3	E	2	BGC	C6-C5-C4	2.61	119.11	113.00
2	D	1	BGC	O2-C2-C3	2.47	116.06	110.35
3	E	3	BGC	O2-C2-C1	2.41	114.08	109.15
2	C	1	BGC	O3-C3-C2	2.40	115.90	110.35
3	E	1	GLC	O6-C6-C5	-2.37	103.15	111.29
2	D	1	BGC	C1-O5-C5	-2.36	109.20	113.66
3	E	1	GLC	C1-O5-C5	2.30	118.00	113.66
3	E	2	BGC	C2-C3-C4	2.14	114.59	110.89
3	E	3	BGC	O4-C4-C5	2.10	114.51	109.30
2	C	1	BGC	O5-C1-C2	-2.08	106.57	110.28
2	D	2	BGC	O3-C3-C4	2.01	115.00	110.35

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	BGC	O5-C5-C6-O6
2	C	2	BGC	O5-C5-C6-O6
2	D	2	BGC	O5-C5-C6-O6
2	D	2	BGC	C4-C5-C6-O6
2	C	2	BGC	C4-C5-C6-O6
2	C	1	BGC	C4-C5-C6-O6
3	E	2	BGC	C4-C5-C6-O6
3	E	3	BGC	O5-C5-C6-O6
2	D	1	BGC	O5-C5-C6-O6
3	E	2	BGC	O5-C5-C6-O6


There are no ring outliers.


5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2	BGC	5	0
3	E	1	GLC	1	0
2	D	1	BGC	3	0
3	E	2	BGC	2	0
3	E	3	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	TRS	A	503	-	7,7,7	0.56	0	9,9,9	4.67	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TRS	A	503	-	-	8/9/9/9	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	503	TRS	C2-C-N	-8.18	83.56	107.98
5	A	503	TRS	C3-C-N	-7.97	84.17	107.98
5	A	503	TRS	C1-C-N	-5.88	90.44	107.98
5	A	503	TRS	C3-C-C2	3.95	123.07	110.81
5	A	503	TRS	C3-C-C1	3.05	120.25	110.81

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	TRS	N-C-C1-O1
5	A	503	TRS	N-C-C2-O2
5	A	503	TRS	N-C-C3-O3
5	A	503	TRS	C1-C-C2-O2
5	A	503	TRS	C2-C-C1-O1
5	A	503	TRS	C1-C-C3-O3
5	A	503	TRS	C2-C-C3-O3
5	A	503	TRS	C3-C-C1-O1

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	503	TRS	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/431 (99%)	-0.28	3 (0%) 87 88	4, 12, 25, 63	0
1	B	430/431 (99%)	-0.33	3 (0%) 87 88	3, 11, 23, 65	0
All	All	860/862 (99%)	-0.31	6 (0%) 87 88	3, 11, 24, 65	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	453	ASP	6.5
1	A	120	GLU	3.3
1	A	121	TYR	3.2
1	B	120	GLU	2.8
1	B	121	TYR	2.5
1	A	453	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	B	23	8/9	0.97	0.07	8,9,10,10	0
1	PCA	A	23	8/9	0.97	0.07	11,11,12,12	0


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	BGC	E	2	11/12	0.69	0.22	34,52,56,59	0
3	GLC	E	1	12/12	0.70	0.27	34,49,55,55	0
3	BGC	E	3	11/12	0.72	0.21	58,60,63,64	0
2	BGC	C	1	12/12	0.85	0.17	30,33,35,36	0
2	BGC	C	2	11/12	0.88	0.13	21,29,32,36	0
2	BGC	D	1	12/12	0.90	0.15	23,29,35,45	0
2	BGC	D	2	11/12	0.90	0.13	19,25,28,31	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	TRS	A	503	8/8	0.82	0.20	26,28,31,32	0
6	CA	B	504	1/1	0.91	0.17	54,54,54,54	0
6	CA	B	503	1/1	0.97	0.41	92,92,92,92	0
4	MG	A	502	1/1	0.98	0.10	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.