



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 01:07 AM BST

PDB ID : 3HDL
Title : Crystal Structure of Highly Glycosylated Peroxidase from Royal Palm Tree
Authors : Watanabe, L.; Moura, P.R.; Bleicher, L.; Nascimento, A.S.; Zamorano, L.S.; Calvete, J.J.; Bursakov, S.; Roig, M.G.; Shnyrov, V.L.; Polikarpov, I.
Deposited on : 2009-05-07
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	
2	B	7	
3	C	5	
4	D	3	
4	E	3	
5	F	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	G	6	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
12	EDO	A	332	-	-	X	-
13	NAG	A	1100	-	-	-	X
13	NAG	A	1200	X	-	-	-
2	MAN	B	5	-	-	-	X
3	MAN	C	3	X	-	-	X
5	NAG	F	2	X	-	-	-
6	MAN	G	5	X	-	-	-

2 Entry composition [i](#)


There are 14 unique types of molecules in this entry. The entry contains 3394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Royal Palm Tree Peroxidase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	304	2288	1425	392	454	17	0	12	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)]beta-D-xylofuranose-(1-6)alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	7	80	45	2	33	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	5	60	34	2	24	0	0	0

- Molecule 4 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			38	22	2	14			
4	E	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	G	6	Total	C	N	O	0	0	0
			71	40	2	29			

- Molecule 7 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
7	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	2	Total	Ca	0	0
			2	2		

- Molecule 9 is HYDROGEN PEROXIDE (three-letter code: PEO) (formula: H₂O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total O 2 2	0	0

- Molecule 10 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total O S 5 4 1	0	0
10	A	1	Total O S 5 4 1	0	0
10	A	1	Total O S 5 4 1	0	0
10	A	1	Total O S 5 4 1	0	0
10	A	1	Total O S 5 4 1	0	0
10	A	1	Total O S 5 4 1	0	0
10	A	1	Total O S 5 4 1	0	0
10	A	1	Total O S 5 4 1	0	0

- Molecule 11 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
11	A	1	12	6	1	4	1	0	0

- Molecule 12 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
12	A	1	4	2	2	0	0
12	A	1	4	2	2	0	0
12	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0
12	A	1	Total 4	C 2	O 2	0	0

- Molecule 13 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
13	A	1	14	8	1	5	0	0
13	A	1	14	8	1	5	0	0
13	A	1	14	8	1	5	0	0


- Molecule 14 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
14	A	586	586	586	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Royal Palm Tree Peroxidase




- Molecule 2: alpha-D-mannopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)]beta-D-xylofuranose-(1-6)alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  67% 33%

MAG1
FUC2
MAG3

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  50% 50%

MAG1
MAG2

- Molecule 6: alpha-D-mannopyranose-(1-2)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  33% 67%

MAG1
MAG2
MAG3
MAG4
MAG5
FUC6

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.82Å 117.82Å 93.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.78 – 1.85 44.78 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.78-1.85) 99.8 (44.78-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 1.86Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.176 , 0.187 0.168 , 0.179	Depositor DCC
R_{free} test set	3241 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtrriage
Anisotropy	0.075	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3394	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA, EDO, PEO, FUC, MES, HEM, MAN, XYZ, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2357	0.44	0/3215

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2288	0	2270	40	0
2	B	80	0	67	1	0
3	C	60	0	52	2	0
4	D	38	0	34	0	0
4	E	38	0	34	0	0
5	F	28	0	25	0	0
6	G	71	0	61	0	0
7	A	43	0	30	4	0
8	A	2	0	0	0	0
9	A	2	0	0	1	0
10	A	40	0	0	0	0
11	A	12	0	12	2	0
12	A	64	0	96	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
13	A	42	0	39	4	0
14	A	586	0	0	20	4
All	All	3394	0	2720	54	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
14:A:570:HOH:O	2:B:6:XYZ:H51	1.63	0.98
1:A:116:THR:OG1	14:A:367:HOH:O	1.90	0.88
1:A:47:ARG:HH12	12:A:326:EDO:H21	1.38	0.88
1:A:232:ASN:HD21	12:A:322:EDO:H12	1.40	0.87
13:A:1100:NAG:H2	14:A:554:HOH:O	1.80	0.82
12:A:335:EDO:O1	14:A:923:HOH:O	2.01	0.78
13:A:1100:NAG:N2	14:A:554:HOH:O	2.25	0.69
1:A:281:LYS:HB2	12:A:321:EDO:H21	1.76	0.67
1:A:185:ASN:HB2	12:A:324:EDO:H22	1.77	0.66
1:A:187:ASN:O	14:A:545:HOH:O	2.15	0.65
13:A:1100:NAG:C2	14:A:554:HOH:O	2.41	0.65
1:A:254[B]:GLU:HG2	14:A:462:HOH:O	1.98	0.64
1:A:77:PHE:HB2	12:A:332:EDO:H21	1.80	0.64
1:A:154:ASN:OD1	14:A:548:HOH:O	2.15	0.63
1:A:1:ASP:CG	14:A:452:HOH:O	2.36	0.63
1:A:205[A]:ARG:NH1	12:A:330:EDO:H22	2.14	0.63
1:A:42:HIS:NE2	9:A:309:PEO:O2	2.31	0.62
1:A:233[B]:MET:HA	1:A:233[B]:MET:HE2	1.81	0.62
1:A:233[A]:MET:HA	1:A:236[A]:THR:HG22	1.81	0.62
1:A:1:ASP:CB	14:A:452:HOH:O	2.47	0.61
1:A:205[B]:ARG:HH11	12:A:323:EDO:H21	1.65	0.61
12:A:332:EDO:H12	14:A:708:HOH:O	2.03	0.58
12:A:327:EDO:H21	14:A:549:HOH:O	2.04	0.58
1:A:2:LEU:HD12	1:A:284:GLN:HG2	1.85	0.58
1:A:165:LEU:HB3	7:A:305:HEM:HMC3	1.86	0.57
1:A:280:VAL:HG12	12:A:321:EDO:H22	1.87	0.55
1:A:60:ASN:ND2	14:A:498:HOH:O	2.32	0.55
1:A:225:ILE:CD1	1:A:236[A]:THR:HG23	2.39	0.52
1:A:38:ARG:HD2	7:A:305:HEM:C4D	2.45	0.52
7:A:305:HEM:HHC	7:A:305:HEM:HBB2	1.91	0.52
1:A:205[A]:ARG:HH12	12:A:330:EDO:H22	1.75	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:A:319:MES:H71	14:A:591:HOH:O	2.09	0.51
1:A:40:HIS:HE1	14:A:509:HOH:O	1.93	0.51
1:A:87:VAL:HG11	1:A:95:VAL:HB	1.93	0.51
1:A:233[A]:MET:SD	1:A:236[A]:THR:HG21	2.51	0.50
1:A:239:GLN:NE2	1:A:262:LYS:HA	2.27	0.49
1:A:280:VAL:CG1	12:A:321:EDO:H22	2.42	0.49
12:A:336:EDO:C2	14:A:754:HOH:O	2.62	0.47
1:A:77:PHE:CB	12:A:332:EDO:H21	2.44	0.46
1:A:47:ARG:NH1	12:A:326:EDO:H21	2.20	0.45
14:A:561:HOH:O	3:C:4:MAN:H62	2.16	0.45
1:A:220:VAL:HG13	12:A:320:EDO:H11	1.98	0.45
11:A:319:MES:H32	14:A:823:HOH:O	2.18	0.44
1:A:262:LYS:O	1:A:266[A]:MET:HG3	2.18	0.43
12:A:332:EDO:H22	3:C:4:MAN:H5	2.00	0.43
1:A:221:SER:HB2	12:A:320:EDO:H22	2.01	0.41
1:A:160:ASP:HB3	12:A:336:EDO:H22	2.02	0.41
1:A:233[A]:MET:CA	1:A:236[A]:THR:HG22	2.47	0.41
1:A:65:LYS:HE2	1:A:77:PHE:CE1	2.55	0.41
7:A:305:HEM:CBB	7:A:305:HEM:HHC	2.50	0.41
1:A:154:ASN:ND2	13:A:700:NAG:H61	2.35	0.41
1:A:188:SER:HB3	14:A:545:HOH:O	2.20	0.40
1:A:185:ASN:HB2	12:A:324:EDO:C2	2.48	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
14:A:575:HOH:O	14:A:580:HOH:O 3_564	1.71	0.49
14:A:518:HOH:O	14:A:582:HOH:O 6_555	1.81	0.39
14:A:913:HOH:O	14:A:915:HOH:O 2_665	1.89	0.31
14:A:488:HOH:O	14:A:492:HOH:O 3_564	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/304 (103%)	311 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/244 (105%)	252 (98%)	4 (2%)	62	49

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	ASP
1	A	101[A]	LEU
1	A	101[B]	LEU
1	A	110	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	HIS
1	A	110	ASN
1	A	118	GLN
1	A	137	GLN
1	A	187	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

26 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.52	0	17,19,21	0.74	0
2	NAG	B	2	2	14,14,15	0.48	0	17,19,21	0.77	0
2	MAN	B	3	2	11,11,12	0.71	0	15,15,17	0.62	0
2	MAN	B	4	2	11,11,12	0.57	0	15,15,17	0.68	0
2	MAN	B	5	2	11,11,12	0.63	0	15,15,17	0.85	0
2	XYZ	B	6	2	9,9,10	0.45	0	10,12,14	1.65	2 (20%)
2	FUC	B	7	2	10,10,11	0.67	0	14,14,16	0.57	0
3	NAG	C	1	1,3	14,14,15	0.45	0	17,19,21	1.02	1 (5%)
3	NAG	C	2	3	14,14,15	0.49	0	17,19,21	0.79	0
3	MAN	C	3	3	11,11,12	0.62	0	15,15,17	0.57	0
3	MAN	C	4	3	11,11,12	0.64	0	15,15,17	1.27	3 (20%)
3	FUC	C	5	3	10,10,11	0.65	0	14,14,16	0.53	0
4	NAG	D	1	1,4	14,14,15	0.48	0	17,19,21	1.11	1 (5%)
4	FUC	D	2	4	10,10,11	0.66	0	14,14,16	0.66	0
4	NAG	D	3	4	14,14,15	0.50	0	17,19,21	0.76	0
4	NAG	E	1	1,4	14,14,15	0.54	0	17,19,21	0.63	0
4	FUC	E	2	4	10,10,11	0.65	0	14,14,16	0.60	0
4	NAG	E	3	4	14,14,15	0.47	0	17,19,21	0.86	1 (5%)
5	NAG	F	1	1,5	14,14,15	0.53	0	17,19,21	0.68	0
5	NAG	F	2	5	14,14,15	0.46	0	17,19,21	0.80	1 (5%)
6	NAG	G	1	1,6	14,14,15	0.47	0	17,19,21	0.93	1 (5%)
6	NAG	G	2	6	14,14,15	0.49	0	17,19,21	0.82	1 (5%)
6	MAN	G	3	6	11,11,12	0.75	0	15,15,17	1.40	3 (20%)
6	MAN	G	4	6	11,11,12	0.58	0	15,15,17	0.60	0
6	MAN	G	5	6	11,11,12	0.69	0	15,15,17	1.25	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FUC	G	6	6	10,10,11	0.69	0	14,14,16	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	MAN	B	3	2	-	1/2/19/22	0/1/1/1
2	MAN	B	4	2	-	1/2/19/22	1/1/1/1
2	MAN	B	5	2	-	2/2/19/22	0/1/1/1
2	XYZ	B	6	2	-	0/2/15/18	0/1/1/1
2	FUC	B	7	2	-	-	0/1/1/1
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	MAN	C	3	3	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
3	FUC	C	5	3	-	-	0/1/1/1
4	NAG	D	1	1,4	-	2/6/23/26	0/1/1/1
4	FUC	D	2	4	-	-	0/1/1/1
4	NAG	D	3	4	-	3/6/23/26	0/1/1/1
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	FUC	E	2	4	-	-	0/1/1/1
4	NAG	E	3	4	-	0/6/23/26	0/1/1/1
5	NAG	F	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	F	2	5	1/1/5/7	4/6/23/26	0/1/1/1
6	NAG	G	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	G	2	6	-	0/6/23/26	0/1/1/1
6	MAN	G	3	6	-	1/2/19/22	0/1/1/1
6	MAN	G	4	6	-	1/2/19/22	1/1/1/1
6	MAN	G	5	6	1/1/4/5	2/2/19/22	0/1/1/1
6	FUC	G	6	6	-	-	0/1/1/1

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	6	XYZ	O4-C4-C3	3.60	107.89	104.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	5	MAN	C3-C4-C5	3.44	116.38	110.24
3	C	1	NAG	C1-O5-C5	3.12	116.42	112.19
2	B	6	XYZ	C1-C2-C3	2.89	106.03	101.63
6	G	3	MAN	C1-C2-C3	2.88	113.20	109.67
3	C	4	MAN	C3-C4-C5	2.77	115.19	110.24
4	D	1	NAG	C1-O5-C5	2.73	115.89	112.19
6	G	3	MAN	C2-C3-C4	2.60	115.40	110.89
6	G	5	MAN	O5-C1-C2	-2.55	106.83	110.77
4	E	3	NAG	C1-O5-C5	2.38	115.41	112.19
6	G	3	MAN	C3-C4-C5	2.31	114.36	110.24
3	C	4	MAN	C1-O5-C5	2.23	115.22	112.19
3	C	4	MAN	O5-C1-C2	-2.17	107.42	110.77
6	G	2	NAG	C1-O5-C5	2.17	115.13	112.19
6	G	1	NAG	C1-O5-C5	2.17	115.13	112.19
5	F	2	NAG	C1-O5-C5	2.01	114.91	112.19

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	C	3	MAN	C1
5	F	2	NAG	C1
6	G	5	MAN	C1

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
4	D	3	NAG	C8-C7-N2-C2
4	D	3	NAG	O7-C7-N2-C2
5	F	2	NAG	C8-C7-N2-C2
4	D	1	NAG	C4-C5-C6-O6
5	F	2	NAG	O7-C7-N2-C2
6	G	5	MAN	C4-C5-C6-O6
3	C	4	MAN	C4-C5-C6-O6
6	G	5	MAN	O5-C5-C6-O6
5	F	2	NAG	C1-C2-N2-C7
6	G	4	MAN	O5-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6
5	F	2	NAG	O5-C5-C6-O6
2	B	5	MAN	O5-C5-C6-O6
2	B	5	MAN	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	4	MAN	O5-C5-C6-O6
2	B	3	MAN	O5-C5-C6-O6
4	D	3	NAG	O5-C5-C6-O6
6	G	3	MAN	O5-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6


All (2) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	4	MAN	C1-C2-C3-C4-C5-O5
6	G	4	MAN	C1-C2-C3-C4-C5-O5


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	6	XYZ	1	0
3	C	4	MAN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 2 are monoatomic - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	EDO	A	327	-	3,3,3	0.44	0	2,2,2	0.44	0
10	SO4	A	312	-	4,4,4	0.11	0	6,6,6	0.13	0
10	SO4	A	318	-	4,4,4	0.14	0	6,6,6	0.14	0
10	SO4	A	315	-	4,4,4	0.13	0	6,6,6	0.11	0
12	EDO	A	330	-	3,3,3	0.47	0	2,2,2	0.30	0
10	SO4	A	311	-	4,4,4	0.15	0	6,6,6	0.09	0
12	EDO	A	332	-	3,3,3	0.42	0	2,2,2	0.41	0
12	EDO	A	331	-	3,3,3	0.48	0	2,2,2	0.28	0
12	EDO	A	328	-	3,3,3	0.46	0	2,2,2	0.37	0
12	EDO	A	323	-	3,3,3	0.46	0	2,2,2	0.39	0
13	NAG	A	700	1	14,14,15	0.48	0	17,19,21	1.12	1 (5%)
12	EDO	A	329	-	3,3,3	0.44	0	2,2,2	0.42	0
12	EDO	A	333	-	3,3,3	0.50	0	2,2,2	0.24	0
12	EDO	A	326	-	3,3,3	0.46	0	2,2,2	0.34	0
12	EDO	A	322	-	3,3,3	0.45	0	2,2,2	0.33	0
11	MES	A	319	-	12,12,12	2.13	1 (8%)	14,16,16	2.50	6 (42%)
13	NAG	A	1100	1	14,14,15	0.46	0	17,19,21	1.50	4 (23%)
12	EDO	A	335	-	3,3,3	0.46	0	2,2,2	0.33	0
9	PEO	A	309	7	1,1,1	0.13	0	-	-	-
12	EDO	A	324	-	3,3,3	0.44	0	2,2,2	0.42	0
12	EDO	A	320	-	3,3,3	0.30	0	2,2,2	0.85	0
12	EDO	A	334	-	3,3,3	0.46	0	2,2,2	0.31	0
12	EDO	A	321	-	3,3,3	0.39	0	2,2,2	0.44	0
7	HEM	A	305	1,9	27,50,50	2.14	5 (18%)	17,82,82	1.93	3 (17%)
10	SO4	A	316	-	4,4,4	0.14	0	6,6,6	0.06	0
13	NAG	A	1200	1	14,14,15	0.53	0	17,19,21	0.76	0
10	SO4	A	317	-	4,4,4	0.13	0	6,6,6	0.11	0
12	EDO	A	336	-	3,3,3	0.44	0	2,2,2	0.31	0
10	SO4	A	313	-	4,4,4	0.14	0	6,6,6	0.06	0
10	SO4	A	314	-	4,4,4	0.14	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	NAG	A	1100	1	-	3/6/23/26	0/1/1/1
12	EDO	A	327	-	-	0/1/1/1	-
12	EDO	A	335	-	-	0/1/1/1	-
12	EDO	A	328	-	-	0/1/1/1	-
12	EDO	A	323	-	-	1/1/1/1	-
13	NAG	A	700	1	-	3/6/23/26	0/1/1/1
12	EDO	A	329	-	-	0/1/1/1	-
12	EDO	A	324	-	-	0/1/1/1	-
12	EDO	A	330	-	-	0/1/1/1	-
12	EDO	A	320	-	-	0/1/1/1	-
12	EDO	A	334	-	-	0/1/1/1	-
12	EDO	A	321	-	-	0/1/1/1	-
12	EDO	A	333	-	-	0/1/1/1	-
13	NAG	A	1200	1	1/1/5/7	5/6/23/26	0/1/1/1
12	EDO	A	332	-	-	1/1/1/1	-
12	EDO	A	326	-	-	0/1/1/1	-
12	EDO	A	336	-	-	0/1/1/1	-
12	EDO	A	322	-	-	1/1/1/1	-
11	MES	A	319	-	-	5/6/14/14	0/1/1/1
7	HEM	A	305	1,9	-	0/6/54/54	-
12	EDO	A	331	-	-	0/1/1/1	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	A	319	MES	C8-S	-7.15	1.67	1.77
7	A	305	HEM	C3D-C2D	5.21	1.53	1.37
7	A	305	HEM	C3B-C2B	-4.53	1.34	1.40
7	A	305	HEM	C3C-C2C	-4.10	1.34	1.40
7	A	305	HEM	C3C-CAC	3.75	1.55	1.47
7	A	305	HEM	C3B-CAB	3.44	1.54	1.47

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	305	HEM	CBD-CAD-C3D	-5.75	101.89	112.48
11	A	319	MES	C5-N4-C3	5.58	121.39	108.83
13	A	1100	NAG	C1-O5-C5	4.01	117.62	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	A	319	MES	C7-N4-C3	3.54	120.30	111.23
11	A	319	MES	O2S-S-C8	3.51	111.15	106.92
11	A	319	MES	O1S-S-C8	3.29	110.87	106.92
13	A	700	NAG	O5-C5-C6	3.09	112.05	107.20
7	A	305	HEM	CAA-CBA-CGA	-2.91	107.80	112.67
11	A	319	MES	C7-N4-C5	2.76	118.28	111.23
13	A	1100	NAG	C1-C2-N2	2.69	115.08	110.49
7	A	305	HEM	C1D-C2D-C3D	-2.62	105.17	107.00
13	A	1100	NAG	C2-N2-C7	2.09	125.89	122.90
11	A	319	MES	O2S-S-O1S	-2.07	106.79	113.95
13	A	1100	NAG	C4-C3-C2	-2.03	108.05	111.02

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
13	A	1200	NAG	C1

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
13	A	700	NAG	C1-C2-N2-C7
13	A	700	NAG	C8-C7-N2-C2
13	A	700	NAG	O7-C7-N2-C2
11	A	319	MES	C7-C8-S-O2S
11	A	319	MES	C7-C8-S-O3S
13	A	1100	NAG	C1-C2-N2-C7
13	A	1100	NAG	C8-C7-N2-C2
13	A	1100	NAG	O7-C7-N2-C2
13	A	1200	NAG	C3-C2-N2-C7
13	A	1200	NAG	C8-C7-N2-C2
13	A	1200	NAG	O7-C7-N2-C2
13	A	1200	NAG	C4-C5-C6-O6
13	A	1200	NAG	O5-C5-C6-O6
11	A	319	MES	C8-C7-N4-C5
12	A	332	EDO	O1-C1-C2-O2
11	A	319	MES	C7-C8-S-O1S
12	A	322	EDO	O1-C1-C2-O2
12	A	323	EDO	O1-C1-C2-O2
11	A	319	MES	C8-C7-N4-C3

There are no ring outliers.

16 monomers are involved in 32 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	A	327	EDO	1	0
12	A	330	EDO	2	0
12	A	332	EDO	4	0
12	A	323	EDO	1	0
13	A	700	NAG	1	0
12	A	326	EDO	2	0
12	A	322	EDO	1	0
11	A	319	MES	2	0
13	A	1100	NAG	3	0
12	A	335	EDO	1	0
9	A	309	PEO	1	0
12	A	324	EDO	2	0
12	A	320	EDO	2	0
12	A	321	EDO	3	0
7	A	305	HEM	4	0
12	A	336	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/304 (100%)	-0.23	2 (0%) 87 88	21, 28, 42, 63	2 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	GLY	2.8
1	A	59	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	B	5	11/12	0.65	0.49	103,109,116,117	0
3	MAN	C	3	11/12	0.78	0.45	68,78,88,89	0
3	MAN	C	4	11/12	0.81	0.32	54,66,92,97	0
4	NAG	D	3	14/15	0.83	0.25	75,80,87,92	0
6	MAN	G	4	11/12	0.84	0.33	72,84,95,105	0
5	NAG	F	1	14/15	0.84	0.34	54,65,76,80	0
5	NAG	F	2	14/15	0.84	0.38	91,96,102,103	0
2	MAN	B	3	11/12	0.85	0.34	59,82,94,99	0
4	FUC	D	2	10/11	0.85	0.33	62,79,83,85	0
3	FUC	C	5	10/11	0.87	0.23	74,83,86,94	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	MAN	G	3	11/12	0.88	0.37	85,101,107,109	0
2	MAN	B	4	11/12	0.88	0.47	78,98,108,108	0
2	XYZ	B	6	9/10	0.88	0.38	94,98,103,108	0
4	NAG	D	1	14/15	0.88	0.18	37,60,71,73	0
4	NAG	E	3	14/15	0.88	0.47	102,108,117,120	0
2	NAG	B	2	14/15	0.89	0.24	37,46,53,57	0
6	NAG	G	2	14/15	0.89	0.23	44,51,65,66	0
4	FUC	E	2	10/11	0.89	0.36	89,94,99,101	0
3	NAG	C	1	14/15	0.90	0.20	52,59,74,91	0
6	MAN	G	5	11/12	0.90	0.48	109,111,115,116	0
4	NAG	E	1	14/15	0.92	0.27	49,70,85,86	0
6	FUC	G	6	10/11	0.94	0.22	49,58,63,65	0
6	NAG	G	1	14/15	0.94	0.12	33,38,48,53	0
3	NAG	C	2	14/15	0.95	0.21	51,60,69,85	0
2	NAG	B	1	14/15	0.96	0.11	30,37,47,48	0
2	FUC	B	7	10/11	0.96	0.16	34,47,49,57	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
12	EDO	A	323	4/4	0.49	0.26	78,83,84,86	0
13	NAG	A	1100	14/15	0.65	0.40	81,102,111,119	0
13	NAG	A	1200	14/15	0.73	0.38	91,108,116,124	0
12	EDO	A	330	4/4	0.75	0.25	45,66,71,73	0
12	EDO	A	324	4/4	0.78	0.24	52,57,57,58	0
13	NAG	A	700	14/15	0.78	0.19	64,95,107,110	0
12	EDO	A	326	4/4	0.81	0.12	59,67,71,73	0
10	SO4	A	317	5/5	0.84	0.39	116,120,122,126	0
12	EDO	A	327	4/4	0.85	0.16	62,63,67,68	0
10	SO4	A	311	5/5	0.87	0.30	116,117,118,123	0
12	EDO	A	335	4/4	0.90	0.14	37,45,54,60	0
10	SO4	A	316	5/5	0.90	0.32	96,106,109,112	0
12	EDO	A	333	4/4	0.91	0.10	29,40,48,49	0
12	EDO	A	334	4/4	0.92	0.10	45,52,55,67	0
12	EDO	A	329	4/4	0.92	0.17	49,51,62,65	0
12	EDO	A	322	4/4	0.93	0.14	45,47,50,67	0
10	SO4	A	318	5/5	0.94	0.15	113,114,116,117	0
12	EDO	A	320	4/4	0.94	0.23	30,32,34,49	0
12	EDO	A	336	4/4	0.94	0.22	38,50,57,60	0
12	EDO	A	328	4/4	0.95	0.14	37,41,53,65	0
12	EDO	A	331	4/4	0.96	0.08	30,37,43,49	0
12	EDO	A	332	4/4	0.96	0.14	55,57,66,66	0
11	MES	A	319	12/12	0.96	0.16	33,36,46,48	0
10	SO4	A	313	5/5	0.96	0.28	74,87,96,102	0
12	EDO	A	321	4/4	0.97	0.29	35,43,46,54	0
10	SO4	A	315	5/5	0.97	0.11	51,55,60,81	0
7	HEM	A	305	43/43	0.98	0.13	18,23,26,29	0
9	PEO	A	309	2/2	0.98	0.13	27,27,27,31	1
10	SO4	A	312	5/5	0.99	0.08	34,40,44,53	0
8	CA	A	307	1/1	0.99	0.12	27,27,27,27	0
10	SO4	A	314	5/5	0.99	0.11	30,42,49,65	0
8	CA	A	306	1/1	1.00	0.10	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.