



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 03:39 am BST

PDB ID : 5HGG  
Title : Crystal structure of uPA in complex with a camelid-derived antibody fragment  
Authors : Yung, K.W.Y.; Kromann-Hansen, T.; Andreasen, P.A.; Ngo, J.C.K.  
Deposited on : 2016-01-08  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	
1	B	246	
2	S	128	
2	T	128	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	301	-	-	X	-
3	GOL	T	202	-	-	X	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6216 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urokinase-type plasminogen activator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	246	1950	1231	343	361	15	0	1	0
1	B	246	1950	1231	343	361	15	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	ALA	CYS	engineered mutation	UNP P00749
A	145	GLN	ASN	engineered mutation	UNP P00749
B	122	ALA	CYS	engineered mutation	UNP P00749
B	145	GLN	ASN	engineered mutation	UNP P00749

- Molecule 2 is a protein called Camelid Derived Antibody Fragment, Nb4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	S	128	952	582	172	193	5	0	0	0
2	T	127	950	581	171	193	5	0	1	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	T	1	Total C O 6 3 3	0	0
3	T	1	Total C O 6 3 3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total O S 5 4 1	0	0
4	S	1	Total O S 5 4 1	0	0

- Molecule 5 is (3S)-3-[(2S,3S,4R)-3,4-DIMETHYLTETRAHYDROFURAN-2-YL]BUTYL LAURATE (three-letter code: TWN) (formula: C<sub>22</sub>H<sub>42</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	S	1	Total C O 25 22 3	0	0
5	S	1	Total C O 25 22 3	0	0
5	T	1	Total C O 25 22 3	0	0

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
6	S	1	12	6	1	4	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	100	Total	O	0	0
			100	100		
7	B	79	Total	O	0	0
			79	79		
7	S	56	Total	O	0	0
			56	56		
7	T	52	Total	O	0	0
			52	52		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Urokinase-type plasminogen activator




- Molecule 1: Urokinase-type plasminogen activator



- Molecule 2: Camelid Derived Antibody Fragment, Nb4



- Molecule 2: Camelid Derived Antibody Fragment, Nb4





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	186.99Å 77.75Å 61.45Å 90.00° 102.40° 90.00°	Depositor
Resolution (Å)	30.00 – 1.97 25.66 – 1.97	Depositor EDS
% Data completeness (in resolution range)	97.4 (30.00-1.97) 97.4 (25.66-1.97)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.99 (at 1.96Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.167 , 0.196 0.177 , 0.207	Depositor DCC
$R_{free}$ test set	2999 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.6	Xtrriage
Anisotropy	0.033	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 43.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6216	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TWN, SO4, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	2/2002 (0.1%)	0.79	3/2711 (0.1%)
1	B	0.74	2/2002 (0.1%)	0.75	1/2711 (0.0%)
2	S	0.73	0/967	0.82	0/1305
2	T	0.70	0/968	0.83	2/1306 (0.2%)
All	All	0.74	4/5939 (0.1%)	0.79	6/8033 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	217[A]	ARG	C-N	13.98	1.58	1.33
1	B	217[B]	ARG	C-N	13.98	1.58	1.33
1	A	217[A]	ARG	C-N	12.94	1.56	1.33
1	A	217[B]	ARG	C-N	12.94	1.56	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	239	ARG	NE-CZ-NH1	6.24	123.42	120.30
1	A	116	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	A	70	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	B	70	ARG	NE-CZ-NH2	-5.49	117.56	120.30
2	T	112	ARG	NE-CZ-NH1	5.06	122.83	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	83	MET	CG-SD-CE	5.05	108.28	100.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	203	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1950	0	1908	3	0
1	B	1950	0	1908	5	0
2	S	952	0	911	5	0
2	T	950	0	911	4	0
3	A	12	0	16	4	0
3	B	6	0	8	0	0
3	T	12	0	16	6	0
4	B	5	0	0	0	0
4	S	5	0	0	0	0
5	S	50	0	84	2	0
5	T	25	0	42	0	0
6	S	12	0	12	0	0
7	A	100	0	0	0	0
7	B	79	0	0	0	0
7	S	56	0	0	0	0
7	T	52	0	0	0	0
All	All	6216	0	5816	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:122:GLN:HG3	3:T:202:GOL:H11	1.68	0.75
3:A:301:GOL:H11	2:S:122:GLN:HG3	1.74	0.69
1:B:223(A):ASP:HB2	3:T:202:GOL:H2	1.81	0.61
3:A:301:GOL:H32	2:S:10:GLY:HA2	1.82	0.60
2:T:122:GLN:CG	3:T:202:GOL:H11	2.35	0.55
1:A:185:ASP:HB2	1:A:188:THR:HG22	1.89	0.55
1:B:163:ILE:HD11	1:B:184:ALA:HB3	1.91	0.53
1:B:185:ASP:HB2	1:B:188:THR:HG22	1.92	0.51
3:A:301:GOL:C3	2:S:10:GLY:HA2	2.41	0.51
5:S:202:TWN:HAO3	5:S:202:TWN:HAK2	1.93	0.49
3:A:301:GOL:H11	2:S:122:GLN:CG	2.45	0.45
2:T:11:LEU:N	3:T:202:GOL:O2	2.49	0.45
1:A:109:ARG:HD2	1:A:110(C):GLY:HA2	2.00	0.44
1:B:223(A):ASP:OD2	3:T:202:GOL:C2	2.66	0.44
2:S:30:ASP:OD1	2:S:30:ASP:O	2.36	0.44
1:B:223(A):ASP:OD2	3:T:202:GOL:H2	2.19	0.43
5:S:201:TWN:HAY2	2:T:45:ARG:HB2	2.02	0.41
1:A:53:ILE:HD11	1:A:103:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/246 (100%)	239 (98%)	6 (2%)	0	100	100
1	B	245/246 (100%)	240 (98%)	5 (2%)	0	100	100
2	S	126/128 (98%)	125 (99%)	1 (1%)	0	100	100
2	T	126/128 (98%)	125 (99%)	1 (1%)	0	100	100
All	All	742/748 (99%)	729 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/214 (100%)	214 (100%)	1 (0%)	88	87
1	B	215/214 (100%)	213 (99%)	2 (1%)	78	77
2	S	100/100 (100%)	99 (99%)	1 (1%)	76	73
2	T	101/100 (101%)	99 (98%)	2 (2%)	55	48
All	All	631/628 (100%)	625 (99%)	6 (1%)	76	73

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	HIS
1	B	57	HIS
1	B	243	LYS
2	S	29	LEU
2	T	29	LEU
2	T	30	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	128	ASN
1	A	204	GLN
1	B	37	HIS
2	T	5	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	T	202	-	5,5,5	0.24	0	5,5,5	0.96	0
6	MES	S	204	-	12,12,12	1.88	1 (8%)	14,16,16	2.61	6 (42%)
3	GOL	B	301	-	5,5,5	0.32	0	5,5,5	0.29	0
3	GOL	A	301	-	5,5,5	0.28	0	5,5,5	0.93	0
3	GOL	T	203	-	5,5,5	0.46	0	5,5,5	0.60	0
3	GOL	A	302	-	5,5,5	0.34	0	5,5,5	0.34	0
4	SO4	B	302	-	4,4,4	0.34	0	6,6,6	0.12	0
5	TWN	T	201	-	25,25,25	1.04	1 (4%)	25,30,30	1.40	4 (16%)
5	TWN	S	202	-	25,25,25	0.98	1 (4%)	25,30,30	1.26	4 (16%)
4	SO4	S	203	-	4,4,4	0.24	0	6,6,6	0.18	0
5	TWN	S	201	-	25,25,25	1.09	1 (4%)	25,30,30	1.46	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	T	202	-	-	2/4/4/4	-
6	MES	S	204	-	-	2/6/14/14	0/1/1/1
3	GOL	B	301	-	-	4/4/4/4	-
3	GOL	A	301	-	-	2/4/4/4	-
3	GOL	T	203	-	-	4/4/4/4	-
3	GOL	A	302	-	-	0/4/4/4	-

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TWN	T	201	-	-	10/21/34/34	0/1/1/1
5	TWN	S	202	-	-	10/21/34/34	0/1/1/1
5	TWN	S	201	-	-	9/21/34/34	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	S	204	MES	C8-S	-6.00	1.69	1.77
5	S	201	TWN	OAI-CAE	5.04	1.48	1.33
5	T	201	TWN	OAI-CAE	4.50	1.46	1.33
5	S	202	TWN	OAI-CAE	4.38	1.46	1.33

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	S	204	MES	C5-N4-C3	4.79	119.61	108.83
6	S	204	MES	O2S-S-C8	4.69	112.56	106.92
6	S	204	MES	C7-N4-C3	4.61	123.02	111.23
5	S	201	TWN	OAI-CAE-CAL	3.62	123.27	111.91
5	T	201	TWN	CAK-CAC-CAB	3.46	121.05	114.16
5	S	201	TWN	CAJ-OAI-CAE	3.32	127.66	116.92
5	S	202	TWN	OAI-CAE-CAL	3.30	122.25	111.91
6	S	204	MES	O1S-S-C8	3.03	110.56	106.92
5	T	201	TWN	OAI-CAE-CAL	2.96	121.19	111.91
5	S	202	TWN	CAK-CAC-CAB	2.52	119.18	114.16
5	S	201	TWN	OAI-CAJ-CAM	2.51	113.79	108.16
5	T	201	TWN	CAO-CAH-CAB	2.34	114.97	111.15
6	S	204	MES	C7-N4-C5	2.31	117.14	111.23
5	S	202	TWN	OAI-CAE-OAG	-2.22	118.00	123.59
5	T	201	TWN	OAI-CAE-OAG	-2.16	118.14	123.59
5	S	201	TWN	OAI-CAE-OAG	-2.15	118.15	123.59
6	S	204	MES	O3S-S-C8	2.13	109.22	105.77
5	S	202	TWN	CAD-OAA-CAB	2.05	112.02	107.84

There are no chirality outliers.

All (43) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	T	202	GOL	O1-C1-C2-C3
6	S	204	MES	N4-C7-C8-S

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	301	GOL	C1-C2-C3-O3
3	B	301	GOL	O2-C2-C3-O3
3	A	301	GOL	O1-C1-C2-C3
3	T	203	GOL	O1-C1-C2-C3
3	T	203	GOL	C1-C2-C3-O3
5	S	201	TWN	OAG-CAE-OAI-CAJ
5	S	201	TWN	CAL-CAE-OAI-CAJ
5	T	201	TWN	CAL-CAE-OAI-CAJ
5	S	201	TWN	CAO-CAH-CAM-CAJ
5	T	201	TWN	OAG-CAE-OAI-CAJ
5	T	201	TWN	CAO-CAH-CAM-CAJ
5	S	201	TWN	OAI-CAJ-CAM-CAH
5	S	201	TWN	CAU-CAT-CAX-CAS
5	S	202	TWN	CAL-CAE-OAI-CAJ
5	S	202	TWN	CAX-CAT-CAU-CAV
5	S	201	TWN	CAW-CAS-CAX-CAT
3	T	202	GOL	C1-C2-C3-O3
3	B	301	GOL	O1-C1-C2-C3
5	T	201	TWN	CAX-CAT-CAU-CAV
5	S	202	TWN	OAG-CAE-OAI-CAJ
3	T	203	GOL	O2-C2-C3-O3
5	S	202	TWN	OAI-CAJ-CAM-CAH
5	T	201	TWN	CAL-CAP-CAW-CAS
5	S	201	TWN	CAB-CAH-CAM-CAJ
5	S	202	TWN	CAX-CAS-CAW-CAP
3	A	301	GOL	O1-C1-C2-O2
3	T	203	GOL	O1-C1-C2-O2
5	S	202	TWN	CAL-CAP-CAW-CAS
6	S	204	MES	C8-C7-N4-C3
5	S	202	TWN	CAY-CAQ-CAR-CAV
5	S	202	TWN	OAA-CAB-CAH-CAM
5	S	202	TWN	CAW-CAS-CAX-CAT
5	T	201	TWN	CAT-CAU-CAV-CAR
5	S	201	TWN	CAL-CAP-CAW-CAS
5	T	201	TWN	OAI-CAJ-CAM-CAH
3	B	301	GOL	O1-C1-C2-O2
5	S	202	TWN	CAT-CAU-CAV-CAR
5	T	201	TWN	OAI-CAE-CAL-CAP
5	T	201	TWN	OAG-CAE-CAL-CAP
5	S	201	TWN	CAC-CAB-CAH-CAM
5	T	201	TWN	CAY-CAQ-CAR-CAV


There are no ring outliers.




4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	T	202	GOL	6	0
3	A	301	GOL	4	0
5	S	202	TWN	1	0
5	S	201	TWN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	246/246 (100%)	0.34	28 (11%) 5 5	15, 26, 57, 71	0
1	B	246/246 (100%)	0.39	21 (8%) 10 12	18, 32, 59, 89	0
2	S	128/128 (100%)	-0.10	3 (2%) 60 62	17, 23, 42, 63	0
2	T	127/128 (99%)	-0.12	2 (1%) 72 73	16, 23, 42, 69	0
All	All	747/748 (99%)	0.20	54 (7%) 15 17	15, 26, 55, 89	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	37(A)	ARG	5.3
1	B	37(C)	GLY	5.2
2	T	75	ALA	4.6
1	B	127	TYR	4.4
1	B	204	GLN	4.3
1	B	110(D)	ARG	4.0
1	B	126	MET	4.0
1	A	37(B)	GLY	3.9
1	A	110(A)	LYS	3.7
1	A	243	LYS	3.7
1	A	110(D)	ARG	3.6
1	B	110(A)	LYS	3.6
2	S	128	ALA	3.6
1	B	243	LYS	3.5
1	B	37(B)	GLY	3.5
1	A	44	GLY	3.4
2	T	1	GLN	3.3
1	A	116	ARG	3.3
1	A	37(C)	GLY	3.3
1	B	110(B)	GLU	3.2
1	B	149	TYR	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	53	ILE	3.2
1	A	110(B)	GLU	3.1
1	B	203	LEU	3.1
1	A	213	VAL	3.0
1	B	205	GLY	3.0
1	A	37(A)	ARG	2.9
1	B	212	ILE	2.9
2	S	1	GLN	2.8
1	A	212	ILE	2.8
1	A	78	GLN	2.7
1	A	166	ARG	2.7
1	A	30	PHE	2.7
2	S	75	ALA	2.6
1	B	125	SER	2.6
1	A	43	GLY	2.6
1	B	128	ASN	2.5
1	A	37(D)	SER	2.5
1	A	205	GLY	2.5
1	A	109	ARG	2.5
1	A	45	SER	2.4
1	A	206	ARG	2.4
1	A	33	ILE	2.4
1	A	197	GLY	2.4
1	B	110(C)	GLY	2.4
1	A	204	GLN	2.3
1	B	131	GLN	2.3
1	B	239	ARG	2.2
1	A	196	GLY	2.2
1	A	52	VAL	2.2
1	A	149	TYR	2.1
1	A	239	ARG	2.1
1	B	213	VAL	2.0
1	B	166	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	TWN	S	201	25/25	0.63	0.40	43,66,116,126	0
5	TWN	T	201	25/25	0.64	0.31	49,58,79,82	0
6	MES	S	204	12/12	0.64	0.39	55,71,127,135	0
5	TWN	S	202	25/25	0.72	0.40	50,68,113,115	0
3	GOL	A	302	6/6	0.80	0.15	44,56,65,68	0
3	GOL	T	203	6/6	0.81	0.20	50,61,68,71	0
3	GOL	B	301	6/6	0.85	0.19	48,54,60,60	0
3	GOL	A	301	6/6	0.92	0.24	29,37,46,47	0
3	GOL	T	202	6/6	0.93	0.35	31,40,47,64	0
4	SO4	S	203	5/5	0.95	0.32	52,63,80,83	0
4	SO4	B	302	5/5	0.95	0.31	60,66,73,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around TWN T 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TWN S 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.