



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 09:37 am BST

PDB ID : 6HHC
Title : Allosteric Inhibition as a new mode of Action for BAY 1213790, a Neutralizing Antibody Targeting the Activated form of Coagulation Factor XI
Authors : Schaefer, M.; Buchmueller, A.; Dittmer, F.; Strassburger, J.; Wilmen, A.
Deposited on : 2018-08-27
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	216	 76% 19% ..
2	H	227	 75% 15% . . 6%
3	A	238	 8% 73% 19% . .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FXIA ANTIBODY FAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	1630	1016	271	338	5	0	0	0

- Molecule 2 is a protein called FXIA ANTIBODY FAB HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	214	1591	1006	265	313	7	0	0	0

- Molecule 3 is a protein called Coagulation factor XI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	229	1820	1151	320	339	10	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	500	SER	CYS	engineered mutation	UNP P03951

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total O S 5 4 1	0	0
4	H	1	Total O S 5 4 1	0	0
4	H	1	Total O S 5 4 1	0	0

- Molecule 5 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	1	Total C O S 4 2 1 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FXIA ANTIBODY FAB LIGHT CHAIN

Chain L: 




- Molecule 2: FXIA ANTIBODY FAB HEAVY CHAIN

Chain H: 



- Molecule 3: Coagulation factor XI

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	61.94Å 70.68Å 185.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.95 – 2.70 32.90 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.6 (92.95-2.70) 99.6 (32.90-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.68Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.227 , 0.314 0.230 , 0.310	Depositor DCC
R_{free} test set	1182 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtrriage
Anisotropy	0.363	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 26.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5060	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.96	0/1663	1.07	9/2258 (0.4%)
2	H	1.16	7/1630 (0.4%)	1.16	7/2220 (0.3%)
3	A	0.92	1/1864 (0.1%)	1.01	1/2526 (0.0%)
All	All	1.01	8/5157 (0.2%)	1.08	17/7004 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	2
3	A	0	1
All	All	0	3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	112	GLN	CD-NE2	5.99	1.47	1.32
2	H	46	GLU	CD-OE1	5.90	1.32	1.25
2	H	17	SER	CB-OG	-5.66	1.34	1.42
3	A	413	ARG	CZ-NH1	5.23	1.39	1.33
2	H	111	GLY	N-CA	-5.23	1.38	1.46
2	H	73	ASP	CB-CG	-5.22	1.40	1.51
2	H	1	GLU	CG-CD	5.21	1.59	1.51
2	H	1	GLU	CD-OE1	5.20	1.31	1.25

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	19	ARG	NE-CZ-NH1	10.02	125.31	120.30
2	H	73	ASP	CB-CG-OD1	-9.82	109.46	118.30
1	L	3	ASP	CB-CG-OD1	-8.56	110.60	118.30
1	L	3	ASP	CB-CG-OD2	8.22	125.69	118.30
1	L	30	ASP	CB-CG-OD1	6.52	124.17	118.30
2	H	43	LYS	N-CA-C	-6.38	93.78	111.00
3	A	564	ARG	NE-CZ-NH2	6.32	123.46	120.30
1	L	203	LEU	CA-CB-CG	6.06	129.24	115.30
1	L	20	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	L	110	ARG	NE-CZ-NH1	-5.84	117.38	120.30
1	L	172	ASP	CB-CG-OD1	5.82	123.54	118.30
1	L	52	ASP	CB-CG-OD1	5.34	123.11	118.30
2	H	87	ARG	NE-CZ-NH1	5.29	122.94	120.30
2	H	98	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	L	213	ARG	NE-CZ-NH1	-5.26	117.67	120.30
2	H	67	ARG	NE-CZ-NH2	5.24	122.92	120.30
2	H	19	ARG	NH1-CZ-NH2	-5.04	113.86	119.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	522	ARG	Peptide
2	H	42	GLY	Peptide
2	H	72	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1630	0	1569	23	0
2	H	1591	0	1551	14	0
3	A	1820	0	1776	23	1
4	H	15	0	0	1	0
5	H	4	0	6	0	0
All	All	5060	0	4902	53	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:598:GLY:HA3	3:A:601:GLN:HE21	1.44	0.81
2:H:42:GLY:HA2	2:H:43:LYS:HG2	1.71	0.72
1:L:110:ARG:HD2	1:L:173:SER:HB2	1.71	0.71
3:A:523:LYS:O	3:A:524:LEU:HB2	1.90	0.70
3:A:450:ASN:ND2	3:A:531:THR:OG1	2.25	0.69
1:L:140:ASN:HD21	2:H:171:HIS:HE1	1.44	0.64
3:A:542:ASN:OD1	3:A:559:ILE:HD12	1.98	0.63
1:L:110:ARG:CD	1:L:173:SER:HB2	2.30	0.61
1:L:140:ASN:HD21	2:H:171:HIS:CE1	2.20	0.59
3:A:388:ILE:HD12	3:A:516:VAL:HG13	1.84	0.58
3:A:396:ARG:NH2	3:A:453:GLU:OE2	2.36	0.57
3:A:523:LYS:O	3:A:524:LEU:CB	2.56	0.54
1:L:117:VAL:HG21	1:L:198:VAL:HG21	1.90	0.53
1:L:142:TYR:CG	1:L:143:SER:HA	2.44	0.53
1:L:114:ALA:HB1	1:L:203:LEU:HD13	1.92	0.52
3:A:598:GLY:HA3	3:A:601:GLN:NE2	2.21	0.51
1:L:156:LEU:HD23	1:L:156:LEU:N	2.28	0.49
3:A:404:THR:OG1	3:A:414:HIS:ND1	2.26	0.48
1:L:177:LEU:C	1:L:177:LEU:HD23	2.33	0.48
2:H:103:TYR:CE2	3:A:452:SER:HB3	2.49	0.47
2:H:128:VAL:O	2:H:216:LYS:HE3	2.14	0.47
2:H:51:ILE:HD11	2:H:55:GLY:HA2	1.96	0.47
1:L:85:ILE:HD12	1:L:106:VAL:O	2.13	0.47
3:A:516:VAL:CG1	3:A:537:ILE:HD11	2.44	0.47
1:L:33:ASN:O	1:L:52:ASP:HA	2.15	0.46
3:A:532:LEU:HD12	3:A:533:GLN:N	2.31	0.46
1:L:162:GLN:HE22	2:H:178:GLN:HA	1.81	0.46
3:A:399:TRP:HE3	3:A:401:TRP:CZ2	2.35	0.45
3:A:502:PRO:HB3	3:A:511:TYR:OH	2.17	0.45
1:L:154:ASN:N	1:L:154:ASN:OD1	2.49	0.45
3:A:502:PRO:HB2	3:A:612:VAL:HG23	1.99	0.44
1:L:91:GLN:HE21	1:L:98:VAL:HG13	1.83	0.43
2:H:76:LYS:O	2:H:77:ASN:HB2	2.19	0.43
3:A:473:LYS:NZ	3:A:473:LYS:HB2	2.34	0.43
2:H:6:GLU:HA	2:H:21:SER:O	2.19	0.42
1:L:52:ASP:HB2	1:L:55:ASN:HD22	1.84	0.42
1:L:48:LEU:C	1:L:48:LEU:HD23	2.39	0.42
1:L:89:TYR:CD2	2:H:45:LEU:HD22	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:103:TYR:HA	3:A:528:ILE:O	2.19	0.42
3:A:400:PRO:HB2	3:A:497:ARG:H	1.86	0.41
3:A:516:VAL:HG12	3:A:537:ILE:HD11	2.02	0.41
3:A:567:GLY:N	3:A:600:ALA:O	2.48	0.41
1:L:3:ASP:C	1:L:3:ASP:OD1	2.58	0.41
1:L:38:TYR:O	1:L:88:TYR:HA	2.20	0.41
1:L:149:GLN:OE1	1:L:156:LEU:HD13	2.20	0.41
2:H:5:LEU:HG	2:H:112:GLN:OE1	2.20	0.41
3:A:592:ILE:HB	3:A:609:THR:HB	2.03	0.41
2:H:56:GLY:O	4:H:301:SO4:O3	2.38	0.41
3:A:439:PRO:HB2	3:A:463:VAL:O	2.21	0.41
1:L:166:THR:HG23	2:H:173:PHE:CE2	2.55	0.41
1:L:37:TRP:CE2	1:L:75:PHE:HB2	2.56	0.41
3:A:614:TYR:O	3:A:617:TRP:HB3	2.21	0.40
1:L:108:ILE:HG22	1:L:109:LYS:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:543:GLU:OE1	3:A:543:GLU:OE1 2_455]	1.54	0.66

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	211/216 (98%)	197 (93%)	13 (6%)	1 (0%)	29 54
2	H	210/227 (92%)	197 (94%)	9 (4%)	4 (2%)	8 20
3	A	225/238 (94%)	199 (88%)	20 (9%)	6 (3%)	5 12
All	All	646/681 (95%)	593 (92%)	42 (6%)	11 (2%)	9 23

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	43	LYS
3	A	456	GLU
3	A	524	LEU
2	H	42	GLY
2	H	57	SER
3	A	423	ASN
3	A	502	PRO
2	H	29	PHE
3	A	561	ALA
1	L	86	ALA
3	A	622	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	187/189 (99%)	171 (91%)	16 (9%)	10	24
2	H	176/188 (94%)	155 (88%)	21 (12%)	5	12
3	A	197/204 (97%)	175 (89%)	22 (11%)	6	13
All	All	560/581 (96%)	501 (90%)	59 (10%)	7	16

All (59) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	5	GLN
1	L	14	SER
1	L	35	LEU
1	L	71	THR
1	L	74	THR
1	L	109	LYS
1	L	136	CYS
1	L	144	ARG
1	L	145	GLU
1	L	149	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	152	VAL
1	L	154	ASN
1	L	161	SER
1	L	201	GLN
1	L	203	LEU
1	L	215	GLU
2	H	1	GLU
2	H	5	LEU
2	H	13	GLN
2	H	21	SER
2	H	30	SER
2	H	45	LEU
2	H	59	VAL
2	H	65	LYS
2	H	73	ASP
2	H	77	ASN
2	H	107	MET
2	H	112	GLN
2	H	119	SER
2	H	122	SER
2	H	133	PRO
2	H	142	THR
2	H	166	LEU
2	H	167	THR
2	H	190	THR
2	H	196	LEU
2	H	217	LYS
3	A	424	GLN
3	A	437	GLU
3	A	455	LYS
3	A	459	SER
3	A	473	LYS
3	A	488	THR
3	A	497	ARG
3	A	512	THR
3	A	516	VAL
3	A	522	ARG
3	A	523	LYS
3	A	524	LEU
3	A	527	LYS
3	A	537	ILE
3	A	555	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	557	LYS
3	A	564	ARG
3	A	565	GLU
3	A	571	CYS
3	A	580	SER
3	A	621	LYS
3	A	622	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	55	ASN
1	L	162	GLN
1	L	201	GLN
2	H	77	ASN
2	H	171	HIS
2	H	206	ASN
3	A	424	GLN
3	A	450	ASN
3	A	601	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	H	301	-	4,4,4	0.39	0	6,6,6	0.59	0
5	DMS	H	304	-	3,3,3	0.33	0	3,3,3	1.11	0
4	SO4	H	303	-	4,4,4	0.44	0	6,6,6	0.80	0
4	SO4	H	302	-	4,4,4	0.24	0	6,6,6	0.55	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	301	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	213/216 (98%)	-0.13	1 (0%) 91 92	17, 29, 61, 69	0
2	H	214/227 (94%)	-0.29	1 (0%) 91 92	19, 23, 42, 67	0
3	A	229/238 (96%)	0.58	18 (7%) 12 10	21, 53, 92, 107	0
All	All	656/681 (96%)	0.07	20 (3%) 50 51	17, 33, 82, 107	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	586	VAL	4.8
3	A	548	ARG	4.3
3	A	551	GLY	3.7
3	A	568	LYS	3.7
3	A	623	GLN	3.4
3	A	550	ARG	3.4
3	A	512	THR	3.3
3	A	571	CYS	3.2
3	A	622	THR	3.1
3	A	585	GLU	3.0
3	A	584	ASN	2.7
3	A	564	ARG	2.7
3	A	549	TYR	2.6
3	A	601	GLN	2.6
3	A	620	GLU	2.5
1	L	128	LYS	2.5
2	H	211	ASN	2.3
3	A	598	GLY	2.3
3	A	542	ASN	2.3
3	A	602	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	DMS	H	304	4/4	0.92	0.21	51,51,52,56	0
4	SO4	H	303	5/5	0.96	0.18	45,46,47,48	0
4	SO4	H	301	5/5	0.99	0.14	24,24,27,27	0
4	SO4	H	302	5/5	0.99	0.11	26,28,29,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.