

wwPDB EM Validation Summary Report (i)

Nov 19, 2022 – 10:43 AM EST

PDB ID	:	2HIL
EMDB ID	:	EMD-1236
Title	:	Structure of the Neisseria gonorrhoeae Type IV pilus filament from x-ray crys-
		tallography and electron cryomicroscopy
Authors	:	Craig, L.; Volkmann, N.; Egelman, E.H.; Tainer, J.A.
Deposited on	:	2006-06-29
Resolution	:	12.50 Å(reported)
Based on initial model	:	2HI2

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 12.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM\ structures}\ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Qua	lity of chain	
			80%		-
1	А	158	37%	49%	13% •
			80%		
1	В	158	37%	49%	13% •
			81%		
1	С	158	36%	50%	13% •
			83%		
1	D	158	36%	50%	13% •
			84%	6	
1	E	158	37%	49%	13% •
			87	7%	
1	F	158	39%	48%	11% •
			8	9%	
1	G	158	40%	47%	11% •
				94%	
1	Н	158	41%	46%	11% •



Conti	nued fron	n previous _l	page					
Mol	Chain	Length	Quality of chain					
1	т	150	99%					
1	1	198	44% 44%	11% •				
1	J	158	35% 51%	13% •				
			81%					
1	K	158	36% 50%	13% •				
1	т	150	82%					
1	L	190	<u> </u>	13% •				
1	М	158	36% 50%	13% •				
			84%					
1	N	158	38% 48%	13% •				
1	0	158	38% //0%	12%				
1	0	100	93%	1276 •				
1	Р	158	42% 46%	11% •				
1	0	150	98%	_				
1	Q	158	40% 47%	11% •				
1	R	158	40% 47%	11%				
			100%					
2	S	2	50% 50%					
9	Т	9	100%					
2	1	<u> </u>	<u> </u>					
2	U	2	50% 50%					
		2	100%					
2	V	2	50% 50%					
2	W	2	50%					
		-	100%					
2	Х	2	50% 50%					
0	V	0	100%					
Z	Y	Z	50% 50% 100%					
2	Ζ	2	50% 50%					
			100%					
2	a	2	100%					
9	Ь	9	100%					
2	D	<u> </u>	<u> </u>					
2	с	2	50% 50%					
			100%					
2	d	2	50% 50%					
2	P	2	50% 50%					
4			100%					
2	f	2	50% 50%					
0		0	100%					
2	g	2	50% 50%	,				



Continued from previous page...

Mol	Chain	Length	Quality of chain						
			100%						
2	h	2	100%						
			100%						
2	i	2	100%						
			100%						
2	j	2	100%						

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	MEA	А	1	-	-	Х	-
1	MEA	В	1	-	-	Х	-
1	MEA	С	1	-	-	Х	-
1	MEA	D	1	-	-	Х	-
1	MEA	Е	1	-	-	Х	-
1	MEA	F	1	-	-	Х	-
1	MEA	G	1	-	-	Х	-
1	MEA	Н	1	-	-	Х	-
1	MEA	J	1	-	-	Х	-
1	MEA	Κ	1	-	-	Х	-
1	MEA	L	1	-	-	Х	-
1	MEA	М	1	-	-	Х	-
1	MEA	N	1	-	-	Х	-
1	MEA	0	1	-	-	Х	-
1	MEA	Р	1	-	-	Х	-
1	MEA	Q	1	-	-	Х	-
1	MEA	R	1	-	-	Х	-



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 22374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			AltConf	Trace
1	А	158	Total	С	Ν	0	S	0	0
		100	1207	760	206	237	4		
1	В	158	Total	С	Ν	Ο	\mathbf{S}	0	0
1	D	100	1207	760	206	237	4	0	0
1	С	158	Total	С	Ν	Ο	\mathbf{S}	0	0
1	U	100	1207	760	206	237	4	0	0
1	Л	158	Total	С	Ν	0	\mathbf{S}	0	0
1	D	100	1207	760	206	237	4	0	0
1	F	159	Total	С	Ν	0	\mathbf{S}	0	0
1	Ľ	100	1207	760	206	237	4	0	0
1	Б	159	Total	С	Ν	0	\mathbf{S}	0	0
1	Г	156	1207	760	206	237	4	0	0
1	C	159	Total	С	Ν	0	S	0	0
1	G	156	1207	760	206	237	4	0	U
1	и	150	Total	С	Ν	0	S	0	0
1	п	100	1207	760	206	237	4	0	0
1	т	150	Total	С	Ν	0	S	0	0
1	1	156	1207	760	206	237	4	0	
1	т	150	Total	С	Ν	0	S	0	0
1	1	156	1207	760	206	237	4	0	0
1	V	150	Total	С	Ν	0	S	0	0
1	n	108	1207	760	206	237	4	0	0
1	т	150	Total	С	Ν	0	S	0	0
1	L	108	1207	760	206	237	4	0	0
1	м	150	Total	С	Ν	0	S	0	0
1	IVI	108	1207	760	206	237	4	0	0
1	N	150	Total	С	Ν	0	S	0	0
1	IN	108	1207	760	206	237	4	0	0
1	0	150	Total	С	Ν	0	S	0	0
	0	198	1207	760	206	237	4	0	0
1	р	150	Total	С	Ν	0	S	0	0
	Г	198	1207	760	206	237	4		0
1	0	150	Total	С	Ν	0	S	0	0
	Q	461	1207	760	206	237	4		

• Molecule 1 is a protein called Fimbrial protein.



Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf	Trace	
1	R	158	Total 1207	C 760	N 206	0 237	$\frac{S}{4}$	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	69	SER	PRO	SEE REMARK 999	UNP P02974
А	71	THR	SER	SEE REMARK 999	UNP P02974
В	69	SER	PRO	SEE REMARK 999	UNP P02974
В	71	THR	SER	SEE REMARK 999	UNP P02974
С	69	SER	PRO	SEE REMARK 999	UNP P02974
С	71	THR	SER	SEE REMARK 999	UNP P02974
D	69	SER	PRO	SEE REMARK 999	UNP P02974
D	71	THR	SER	SEE REMARK 999	UNP P02974
Е	69	SER	PRO	SEE REMARK 999	UNP P02974
Е	71	THR	SER	SEE REMARK 999	UNP P02974
F	69	SER	PRO	SEE REMARK 999	UNP P02974
F	71	THR	SER	SEE REMARK 999	UNP P02974
G	69	SER	PRO	SEE REMARK 999	UNP P02974
G	71	THR	SER	SEE REMARK 999	UNP P02974
Н	69	SER	PRO	SEE REMARK 999	UNP P02974
Н	71	THR	SER	SEE REMARK 999	UNP P02974
Ι	69	SER	PRO	SEE REMARK 999	UNP P02974
Ι	71	THR	SER	SEE REMARK 999	UNP P02974
J	69	SER	PRO	SEE REMARK 999	UNP P02974
J	71	THR	SER	SEE REMARK 999	UNP P02974
K	69	SER	PRO	SEE REMARK 999	UNP P02974
K	71	THR	SER	SEE REMARK 999	UNP P02974
L	69	SER	PRO	SEE REMARK 999	UNP P02974
L	71	THR	SER	SEE REMARK 999	UNP P02974
М	69	SER	PRO	SEE REMARK 999	UNP P02974
М	71	THR	SER	SEE REMARK 999	UNP P02974
N	69	SER	PRO	SEE REMARK 999	UNP P02974
N	71	THR	SER	SEE REMARK 999	UNP P02974
0	69	SER	PRO	SEE REMARK 999	UNP P02974
0	71	THR	SER	SEE REMARK 999	UNP P02974
P	69	SER	PRO	SEE REMARK 999	UNP P02974
P	71	THR	SER	SEE REMARK 999	UNP P02974
Q	69	SER	PRO	SEE REMARK 999	UNP P02974
Q	71	THR	SER	SEE REMARK 999	UNP P02974
R	69	SER	PRO	SEE REMARK 999	UNP P02974
R	71	THR	SER	SEE REMARK 999	UNP P02974



• Molecule 2 is an oligosaccharide called alpha-D-galactopyranose-(1-3)-2,4-bisacetamido-2,4-dideoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			AltConf	Trace	
2	S	2	Total	С	Ν	0	0	0
	0	2	28	16	2	10	0	0
2	Т	2	Total	\mathbf{C}	Ν	Ο	0	0
	1	2	28	16	2	10	0	0
2	U	2	Total	\mathbf{C}	Ν	Ο	0	0
		-	28	16	2	10	Ŭ	
2	V	2	Total	С	Ν	Ο	0	0
	•	_	28	16	2	10		
2	W	2	Total	С	Ν	0	0	0
			28	16	2	10	_	
2	Х	2	Total	C	N	0	0	0
			28	<u>16</u>	2	10		
2	Y	2	Total	C	N	0	0	0
			28	$\frac{10}{0}$	2	10		
2	Z	2	Total	C	N	0	0	0
			28	$\frac{10}{C}$	2	$\frac{10}{0}$		
2	a	2		16	N ภ	10	0	0
			Zð Tatal	$\frac{10}{C}$	Z N	10		
2	b	2	10tai	16	IN O	10	0	0
			Z0 Total	$\frac{10}{C}$	$\frac{2}{N}$	$\frac{10}{0}$		
2	с	2	10tai 28	16	1N 9	10	0	0
			 Total	$\frac{10}{C}$	Z N	$\frac{10}{0}$		
2	d	2	28	16	2	10	0	0
			Total	$\frac{10}{C}$	N	$\frac{10}{0}$		
2	е	2	28	16	2	10	0	0
			Total	$\frac{10}{C}$	N	$\frac{10}{0}$		
2	f	2	28	16	2	10	0	0
		_	Total	C	N	0		
2	g	2	28	16	2	10	0	0
	1	2	Total	С	Ν	0	0	0
2	h	2	28	16	2	10	0	0
0	•	0	Total	С	Ν	0	0	0
2	1	2	28	16	2	10	U	
0	<u>.</u>	0	Total	С	Ν	0	0	0
	J	2	28	16	2	10	U	U



• Molecule 3 is PHOSPHORIC ACID MONO-(2-AMINO-ETHYL) ESTER (three-letter code: OPE) (formula: $C_2H_8NO_4P$).



Mol	Chain	Residues		Ate	oms			AltConf
9	٨	1	Total	С	Ν	Ο	Р	0
3	А	L	8	2	1	4	1	0
2	Р	1	Total	С	Ν	Ο	Р	0
່ <u>ບ</u>	D	L	8	2	1	4	1	0
3	С	1	Total	С	Ν	0	Р	0
5	U	T	8	2	1	4	1	0
3	Л	1	Total	С	Ν	Ο	Р	0
5	D	I	8	2	1	4	1	0
3	E	1	Total	С	Ν	Ο	Р	0
0	Ц	T	8	2	1	4	1	0
3	F	1	Total	С	Ν	Ο	Р	0
0	1	T	8	2	1	4	1	0
3	G	1	Total	С	Ν	Ο	Р	0
0		1	8	2	1	4	1	0
3	Н	1	Total	С	Ν	Ο	Р	0
		1	8	2	1	4	1	0
3	T	1	Total	С	Ν	Ο	Р	0
	-	-	8	2	1	4	1	
3	J	1	Total	С	Ν	Ο	Р	0
		-	8	2	1	4	1	
3	K	1	Total	С	Ν	0	Р	0
		-	8	2	1	4	1	
3	L	1	Total	С	Ν	0	Р	0
	-	-	8	2	1	4	1	Ĭ



Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf				
2	М	1	Total	С	Ν	0	Р	0				
່ <u>ບ</u>	111	1	8	2	1	4	1	0				
2	N	1	Total	С	Ν	0	Р	0				
5	11	1	8	2	1	4	1	0				
2	0	1	Total	С	Ν	0	Р	0				
5	U	0	0	0	U	1	8	2	1	4	1	0
2	р	1	Total	С	Ν	Ο	Р	0				
5	1	1	8	2	1	4	1	0				
3	Q	2 0	0 1	1	Total	С	Ν	0	Р	0		
5		1	8	2	1	4	1	0				
2	В	1	Total	С	N	0	Р	0				
3	к	R	R	1	8	2	1	4	1			



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fimbrial protein

F1 172 14 14 15 16 18 V9 V9



G65 A67 A67 A67 A67 A67 A67 A67 A67 S68 S68 S68 S68 S68 S68 S68 S68 S68 S69 C75 V77 V78 V79 V64 V

V125 V126 R127 T126 D131 D131 D131 V133 A134 A134 A134 A134 D138 C131 D138 C140 E141 D147 D148 S149 C151 C151 A156 A156

• Molecule 1: Fimbrial protein



• Molecule 1: Fimbrial protein



• Molecule 1: Fimbrial protein





• Molecule 1: Fimbrial protein





• Molecule 1: Fimbrial protein



• Molecule 1: Fimbrial protein



• Molecule 1: Fimbrial protein





PDB IN DATA BANK

• Molecule 1: Fimbrial protein





• Molecule 1: Fimbrial protein



• Molecule 1: Fimbrial protein







٠	•
DT61	GLA2

• Molecule 2: e	alpha-D-galactopyr	anose-(1-3)-2,4-b	isacetamido-2,4-d	ideoxy-beta-D-gluc	opyranos
Chain V:	50%	100%	50%		
DTG1					
• Molecule 2: e	alpha-D-galactopyr	anose-(1-3)-2,4-b	isacetamido-2,4-d	ideoxy-beta-D-gluc	opyranos
Chain W:	50%	100%	50%		
brei GLA2					
• Molecule 2: e	alpha-D-galactopyr	anose-(1-3)-2,4-b	isacetamido-2,4-d	ideoxy-beta-D-gluc	opyranos
Chain X:	50%	100%	50%	_	
• Molecule 2:	alpha-D-galactopyr	anose-(1-3)-2,4-b	isacetamido-2,4-d	ideoxy-beta-D-gluc	opyranos
е		100%			
Chain Y:	50%		50%		
DT61					
• Molecule 2:	alpha-D-galactopyr	anose- $(1-3)-2,4-b$	isacetamido-2,4-d	ideoxy-beta-D-gluc	opyranos
e		100%			
Chain Z:	50%		50%		
bT61 GLA2 ♦					
• Molecule 2:	alpha-D-galactopyr	anose-(1-3)-2,4-b	isacetamido-2,4-d	ideoxy-beta-D-gluc	opyranos
е		100%			
Chain a:		100%			
		WORLDW PROTEIN DATA			

•		•
	DT61	GLA2

• Molecule e	$ 2: \ alpha-D-galactopyranose-(1-3)-2, 4-bisacetamido-2, 4-dideoxy-beta-D-glucopyranose-(1-3)-2, 4-bisacetamido-2, 4-bisacetamido-$
Chain b:	100% 50% 50%
DT61	
• Molecule e	2: alpha-D-galactopyranose-(1-3)-2, 4-bisacetamido-2, 4-dideoxy-beta-D-glucopyranose-(1-3)-2, 4-bisacetamido-2, 4-bi
Chain c:	100% 50%
DT61 GLA2	
• Molecule e	2: alpha-D-galactopyranose-(1-3)-2,4-bisacetamido-2,4-dideoxy-beta-D-glucopyranos
Chain d:	100% 50% 50%
bT61 GLA2 ♦	
• Molecule e	2: alpha-D-galactopyranose-(1-3)-2,4-bisacetamido-2,4-dideoxy-beta-D-glucopyranos
Chain e:	100% 50% 50%
DT61 GLA2	
• Molecule	$ 2: \ alpha-D-galactopyranose-(1-3)-2, 4-bisacetamido-2, 4-dideoxy-beta-D-glucopyranose-(1-3)-2, 4-bisacetamido-2, 4-bis$
е	100%
Chain f:	50% 50%
DT61	
• Molecule	2: alpha-D-galactopyranose-(1-3)-2,4-bisacetamido-2,4-dideoxy-beta-D-glucopyranos
	100%
Chain g:	50% 50%



• Molecule 2: alpha-D-galactopyranose-(1-3)-2,4-bisacetamido-2,4-dideoxy-beta-D-glucopyranos e

	100%	
Chain h:	100%	
DT61		
• Molecule 2: alpha	a-D-galactopyranose-(1-3)-2,4-bisacetamido-2,4	-dideoxy-beta-D-glucopyranos
e		
	100%	
Chain i:	100%	
DT61 GLA2 ♦		
• Molecule 2: alpha	a-D-galactopyranose-(1-3)-2,4-bisacetamido-2,4	-dideoxy-beta-D-glucopyranos
e		
	100%	
Chain j:	100%	
••		





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=Not provided°, rise=Not	Depositor
	provided Å, axial sym=Not provided	
Number of segments used	25000	Depositor
Resolution determination method	Not provided	
CTF correction method	Wiener filter	Depositor
Microscope	FEI/PHILIPS CM200FEG	Depositor
Voltage (kV)	120	Depositor
Electron dose $(e^-/\text{\AA}^2)$	10	Depositor
Minimum defocus (nm)	1100	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	50000	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	3.946	Depositor
Minimum map value	-0.810	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	2.05	Depositor
Map size (Å)	101.6, 101.6, 152.4	wwPDB
Map dimensions	37, 37, 54	wwPDB
Map angles $(^{\circ})$	90, 90, 90	wwPDB
Pixel spacing (Å)	2.54, 2.54, 2.54	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OPE, GLA, DT6, MEA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	ond lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	В	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	С	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	D	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	Ε	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	F	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	G	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	Н	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	Ι	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	J	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	Κ	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	L	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	М	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	Ν	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	0	0.58	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	Р	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	Q	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
1	R	0.59	1/1215~(0.1%)	0.66	1/1647~(0.1%)	
All	All	0.59	18/21870~(0.1%)	0.66	18/29646~(0.1%)	

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	Е	26	ASP	N-CA	15.11	1.76	1.46
1	R	26	ASP	N-CA	15.11	1.76	1.46
1	Q	26	ASP	N-CA	15.10	1.76	1.46
1	D	26	ASP	N-CA	15.10	1.76	1.46
1	F	26	ASP	N-CA	15.09	1.76	1.46

The worst 5 of 18 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	Ν	25	GLN	C-N-CA	-7.58	102.74	121.70
1	0	25	GLN	C-N-CA	-7.58	102.75	121.70
1	Р	25	GLN	C-N-CA	-7.58	102.75	121.70
1	М	25	GLN	C-N-CA	-7.58	102.76	121.70
1	Q	25	GLN	C-N-CA	-7.58	102.76	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1207	0	1199	497	0
1	В	1207	0	1199	495	0
1	С	1207	0	1199	490	0
1	D	1207	0	1199	491	0
1	Е	1207	0	1199	497	0
1	F	1207	0	1201	409	0
1	G	1207	0	1201	379	0
1	Н	1207	0	1201	380	0
1	Ι	1207	0	1202	293	0
1	J	1207	0	1199	492	0
1	К	1207	0	1199	496	0
1	L	1207	0	1199	501	0
1	М	1207	0	1199	497	0
1	N	1207	0	1199	493	0
1	0	1207	0	1203	415	0
1	Р	1207	0	1203	377	0
1	Q	1207	0	1203	377	0
1	R	1207	0	1203	296	0
2	S	28	0	25	1	0
2	Т	28	0	25	1	0
2	U	28	0	25	1	0
2	V	28	0	25	1	0
2	W	28	0	25	1	0
2	Х	28	0	25	1	0
2	Y	28	0	25	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Ζ	28	0	25	1	0
2	a	28	0	25	0	0
2	b	28	0	25	0	0
2	с	28	0	25	0	0
2	d	28	0	25	0	0
2	е	28	0	25	0	0
2	f	28	0	25	0	0
2	g	28	0	25	0	0
2	h	28	0	25	0	0
2	i	28	0	25	0	0
2	j	28	0	25	0	0
3	А	8	0	6	0	0
3	В	8	0	6	0	0
3	С	8	0	6	0	0
3	D	8	0	6	0	0
3	Е	8	0	6	0	0
3	F	8	0	6	0	0
3	G	8	0	6	0	0
3	Н	8	0	6	0	0
3	Ι	8	0	6	0	0
3	J	8	0	6	0	0
3	K	8	0	6	0	0
3	L	8	0	6	0	0
3	М	8	0	6	0	0
3	Ν	8	0	6	0	0
3	0	8	0	6	0	0
3	Р	8	0	6	0	0
3	Q	8	0	6	0	0
3	R	8	0	6	0	0
All	All	22374	0	22165	4761	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 107.

The worst 5 of 4761 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:8:ILE:HD11	1:L:1:MEA:CE2	1.24	1.68
1:E:1:MEA:CE2	1:F:8:ILE:HD11	1.24	1.66
1:Q:8:ILE:HD11	1:R:1:MEA:CE2	1.24	1.64
1:C:1:MEA:CE2	1:D:8:ILE:HD11	1.24	1.64



α \cdot \cdot \cdot	C		
Continued	trom	previous	page
	J	1	I J

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1:MEA:CE2	1:C:8:ILE:HD11	1.24	1.62

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	erce	$\mathbf{entiles}$;
1	А	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	В	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	С	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	D	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	Е	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	F	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	G	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	Н	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	Ι	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	J	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	К	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	L	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	М	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	Ν	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	Ο	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	Р	156/158~(99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	Q	156/158 (99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
1	R	156/158 (99%)	138 (88%)	9 (6%)	9 (6%)		1	18	
All	All	2808/2844 (99%)	2484 (88%)	162 (6%)	162 (6%)		3	18	



5 of 162 Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	93	LEU
1	А	98	ASN
1	В	93	LEU
1	В	98	ASN
1	С	93	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	eric Outliers		Percentiles		
1	А	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	В	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	С	128/129 (99%)	112 (88%)	16 (12%)		4	19	
1	D	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	Е	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	F	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	G	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	Н	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	Ι	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	J	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	К	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	L	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	М	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	Ν	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	Ο	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	Р	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	Q	128/129~(99%)	112 (88%)	16 (12%)		4	19	
1	R	128/129 (99%)	112 (88%)	16 (12%)		4	19	
All	All	2304/2322 (99%)	2016 (88%)	288 (12%)		8	19	



5 of 288 residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	0	69	SER
1	R	137	LYS
1	0	137	LYS
1	Q	39	LEU
1	G	21	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 104 such side chains are listed below:

Mol	Chain	\mathbf{Res}	Type
1	Κ	53	ASN
1	М	99	ASN
1	R	53	ASN
1	Κ	99	ASN
1	L	99	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

18 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	Chain Bog		Dog	Tink	Bo	ond leng	ths	В	ond ang	les
	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2		
1	MEA	Ι	1	1	10,11,13	0.59	0	10,13,16	0.19	0		
1	MEA	R	1	1	10,11,13	0.60	0	10,13,16	0.20	0		
1	MEA	М	1	1	10,11,13	0.59	0	10,13,16	0.20	0		
1	MEA	K	1	1	10,11,13	0.58	0	10,13,16	0.19	0		
1	MEA	Н	1	1	10,11,13	0.60	0	10,13,16	0.19	0		
1	MEA	D	1	1	10,11,13	0.59	0	10,13,16	0.19	0		
1	MEA	А	1	1	10,11,13	0.58	0	10,13,16	0.19	0		
1	MEA	J	1	1	10,11,13	0.58	0	10,13,16	0.19	0		



Mal	Turne	Chain	Pog Link		Bo	ond leng	$_{\rm ths}$	В	ond ang	les										
MOI	туре	Ullalli	ries	ries	nes	nes	nes	nes	ries	nes	nes	ries	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MEA	N	1	1	$10,\!11,\!13$	0.60	0	10, 13, 16	0.20	0										
1	MEA	Q	1	1	$10,\!11,\!13$	0.60	0	10,13,16	0.21	0										
1	MEA	L	1	1	10,11,13	0.59	0	10,13,16	0.19	0										
1	MEA	0	1	1	10,11,13	0.60	0	10,13,16	0.21	0										
1	MEA	F	1	1	$10,\!11,\!13$	0.59	0	$10,\!13,\!16$	0.19	0										
1	MEA	Р	1	1	10,11,13	0.60	0	10,13,16	0.21	0										
1	MEA	В	1	1	10,11,13	0.58	0	10,13,16	0.18	0										
1	MEA	G	1	1	10,11,13	0.59	0	10,13,16	0.19	0										
1	MEA	С	1	1	10,11,13	0.59	0	10,13,16	0.18	0										
1	MEA	Е	1	1	10,11,13	0.59	0	10,13,16	0.19	0										

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MEA	Ι	1	1	-	0/5/6/10	0/1/1/1
1	MEA	R	1	1	-	0/5/6/10	0/1/1/1
1	MEA	М	1	1	-	0/5/6/10	0/1/1/1
1	MEA	K	1	1	-	0/5/6/10	0/1/1/1
1	MEA	Н	1	1	-	0/5/6/10	0/1/1/1
1	MEA	D	1	1	-	0/5/6/10	0/1/1/1
1	MEA	А	1	1	-	0/5/6/10	0/1/1/1
1	MEA	J	1	1	-	0/5/6/10	0/1/1/1
1	MEA	N	1	1	-	0/5/6/10	0/1/1/1
1	MEA	Q	1	1	-	0/5/6/10	0/1/1/1
1	MEA	L	1	1	-	0/5/6/10	0/1/1/1
1	MEA	0	1	1	-	0/5/6/10	0/1/1/1
1	MEA	F	1	1	-	0/5/6/10	0/1/1/1
1	MEA	Р	1	1	-	0/5/6/10	0/1/1/1
1	MEA	В	1	1	-	0/5/6/10	0/1/1/1
1	MEA	G	1	1	-	0/5/6/10	0/1/1/1
1	MEA	С	1	1	-	0/5/6/10	0/1/1/1
1	MEA	Е	1	1	-	0/5/6/10	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	R	1	MEA	17	0
1	М	1	MEA	17	0
1	Κ	1	MEA	15	0
1	Н	1	MEA	17	0
1	D	1	MEA	17	0
1	А	1	MEA	17	0
1	J	1	MEA	16	0
1	N	1	MEA	16	0
1	Q	1	MEA	16	0
1	L	1	MEA	17	0
1	0	1	MEA	17	0
1	F	1	MEA	17	0
1	Р	1	MEA	17	0
1	В	1	MEA	17	0
1	G	1	MEA	16	0
1	С	1	MEA	15	0
1	Е	1	MEA	15	0

17 monomers are involved in 279 short contacts:

5.5 Carbohydrates (i)

36 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	Dec	Tiple	Bo	Bond lengths			Bond angles		
IVIOI	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	DT6	S	1	2,1	17,17,18	0.61	0	$17,\!23,\!25$	0.69	1 (5%)	
2	GLA	S	2	2	11,11,12	0.39	0	$15,\!15,\!17$	0.74	0	
2	DT6	Т	1	2,1	17,17,18	0.61	0	$17,\!23,\!25$	0.68	1 (5%)	
2	GLA	Т	2	2	11,11,12	0.39	0	$15,\!15,\!17$	0.74	0	
2	DT6	U	1	2,1	17,17,18	0.62	0	$17,\!23,\!25$	0.69	1 (5%)	
2	GLA	U	2	2	11,11,12	0.39	0	$15,\!15,\!17$	0.74	0	
2	DT6	V	1	2,1	17,17,18	0.61	0	$17,\!23,\!25$	0.68	1 (5%)	
2	GLA	V	2	2	11,11,12	0.39	0	$15,\!15,\!17$	0.74	0	
2	DT6	W	1	2,1	17,17,18	0.61	0	17,23,25	0.68	1 (5%)	



Mal	Tuno	Chain	Dog	Tink	Bo	ond lengths		Bond angles		
IVIOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLA	W	2	2	11,11,12	0.38	0	$15,\!15,\!17$	0.74	0
2	DT6	Х	1	2,1	17,17,18	0.61	0	17,23,25	0.68	1 (5%)
2	GLA	Х	2	2	11,11,12	0.37	0	15,15,17	0.75	0
2	DT6	Y	1	2,1	17,17,18	0.62	0	17,23,25	0.69	1 (5%)
2	GLA	Y	2	2	11,11,12	0.38	0	15,15,17	0.74	0
2	DT6	Z	1	2,1	17,17,18	0.62	0	17,23,25	0.69	1 (5%)
2	GLA	Z	2	2	11,11,12	0.39	0	15,15,17	0.74	0
2	DT6	a	1	2,1	17,17,18	0.61	0	17,23,25	0.68	0
2	GLA	a	2	2	11,11,12	0.38	0	$15,\!15,\!17$	0.74	0
2	DT6	b	1	2,1	17,17,18	0.62	0	17,23,25	0.70	1 (5%)
2	GLA	b	2	2	11,11,12	0.39	0	$15,\!15,\!17$	0.75	0
2	DT6	с	1	2,1	17,17,18	0.62	0	17,23,25	0.69	1 (5%)
2	GLA	с	2	2	11,11,12	0.38	0	15,15,17	0.74	0
2	DT6	d	1	2,1	17,17,18	0.62	0	17,23,25	0.70	1 (5%)
2	GLA	d	2	2	11,11,12	0.38	0	15,15,17	0.74	0
2	DT6	е	1	2,1	17,17,18	0.62	0	17,23,25	0.69	1 (5%)
2	GLA	е	2	2	11,11,12	0.38	0	15,15,17	0.74	0
2	DT6	f	1	2,1	17,17,18	0.62	0	17,23,25	0.69	1 (5%)
2	GLA	f	2	2	11,11,12	0.38	0	15,15,17	0.74	0
2	DT6	g	1	2,1	17,17,18	0.62	0	17,23,25	0.68	1 (5%)
2	GLA	g	2	2	11,11,12	0.38	0	$15,\!15,\!17$	0.74	0
2	DT6	h	1	2,1	17,17,18	0.62	0	17,23,25	0.67	0
2	GLA	h	2	2	11,11,12	0.38	0	15,15,17	0.75	0
2	DT6	i	1	2,1	17,17,18	0.63	0	17,23,25	0.67	0
2	GLA	i	2	2	11,11,12	0.37	0	15,15,17	0.75	0
2	DT6	j	1	2,1	17,17,18	0.63	0	17,23,25	0.67	0
2	GLA	j	2	2	11,11,12	0.37	0	15,15,17	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DT6	S	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	S	2	2	-	1/2/19/22	0/1/1/1
2	DT6	Т	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	Т	2	2	-	1/2/19/22	0/1/1/1
2	DT6	U	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	U	2	2	-	1/2/19/22	0/1/1/1
2	DT6	V	1	2,1	-	1/10/27/30	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLA	V	2	2	-	1/2/19/22	0/1/1/1
2	DT6	W	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	W	2	2	-	1/2/19/22	0/1/1/1
2	DT6	Х	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	Х	2	2	-	1/2/19/22	0/1/1/1
2	DT6	Y	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	Y	2	2	-	1/2/19/22	0/1/1/1
2	DT6	Ζ	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	Ζ	2	2	-	1/2/19/22	0/1/1/1
2	DT6	a	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	a	2	2	-	1/2/19/22	0/1/1/1
2	DT6	b	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	b	2	2	-	1/2/19/22	0/1/1/1
2	DT6	с	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	с	2	2	-	1/2/19/22	0/1/1/1
2	DT6	d	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	d	2	2	-	1/2/19/22	0/1/1/1
2	DT6	е	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	е	2	2	-	1/2/19/22	0/1/1/1
2	DT6	f	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	f	2	2	-	1/2/19/22	0/1/1/1
2	DT6	g	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	g	2	2	-	1/2/19/22	0/1/1/1
2	DT6	h	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	h	2	2	-	1/2/19/22	0/1/1/1
2	DT6	i	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	i	2	2	-	1/2/19/22	0/1/1/1
2	DT6	j	1	2,1	-	1/10/27/30	0/1/1/1
2	GLA	j	2	2	-	1/2/19/22	0/1/1/1

Continued from previous page...

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	d	1	DT6	C2-N2-C7	-2.07	119.96	122.90
2	b	1	DT6	C2-N2-C7	-2.05	119.98	122.90
2	е	1	DT6	C2-N2-C7	-2.04	120.00	122.90
2	f	1	DT6	C2-N2-C7	-2.04	120.00	122.90
2	с	1	DT6	C2-N2-C7	-2.03	120.01	122.90

There are no chirality outliers.

5 of 36 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	W	2	GLA	O5-C5-C6-O6
2	Х	2	GLA	O5-C5-C6-O6
2	h	2	GLA	O5-C5-C6-O6
2	i	2	GLA	O5-C5-C6-O6
2	S	2	GLA	O5-C5-C6-O6

There are no ring outliers.

16 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	U	2	GLA	1	0
2	S	2	GLA	1	0
2	V	2	GLA	1	0
2	W	1	DT6	1	0
2	Х	1	DT6	1	0
2	V	1	DT6	1	0
2	Х	2	GLA	1	0
2	Y	2	GLA	1	0
2	W	2	GLA	1	0
2	S	1	DT6	1	0
2	Т	1	DT6	1	0
2	Y	1	DT6	1	0
2	Ζ	1	DT6	1	0
2	Т	2	GLA	1	0
2	Ζ	2	GLA	1	0
2	U	1	DT6	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









































































5.6 Ligand geometry (i)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tiple	B	Bond lengths			Bond angles		
	Type	Unain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2	
3	OPE	N	824	1	7,7,7	0.89	0	$9,\!9,\!9$	0.97	0	
3	OPE	G	824	1	7,7,7	0.88	0	9,9,9	0.96	0	
3	OPE	J	824	1	7,7,7	0.88	0	9,9,9	0.97	0	
3	OPE	0	824	1	7,7,7	0.89	0	9,9,9	0.97	0	
3	OPE	Е	824	1	7,7,7	0.88	0	9,9,9	0.97	0	
3	OPE	F	824	1	7,7,7	0.88	0	9,9,9	0.97	0	



Mal	Turne	Chain	Dec	Tink	B	ond leng	gths	E	Bond ang	gles
WIOI	туре	Unann	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
3	OPE	K	824	1	7,7,7	0.89	0	$9,\!9,\!9$	0.97	0
3	OPE	R	824	1	7,7,7	0.88	0	$9,\!9,\!9$	0.97	0
3	OPE	Р	824	1	7,7,7	0.88	0	9,9,9	0.97	0
3	OPE	D	824	1	7,7,7	0.88	0	$9,\!9,\!9$	0.97	0
3	OPE	В	824	1	7,7,7	0.88	0	9,9,9	0.97	0
3	OPE	А	824	1	7,7,7	0.89	0	$9,\!9,\!9$	0.97	0
3	OPE	Q	824	1	7,7,7	0.88	0	$9,\!9,\!9$	0.97	0
3	OPE	Ι	824	1	7,7,7	0.87	0	$9,\!9,\!9$	0.96	0
3	OPE	Н	824	1	7,7,7	0.88	0	9,9,9	0.97	0
3	OPE	М	824	1	7,7,7	0.89	0	$9,\!9,\!9$	0.97	0
3	OPE	L	824	1	7,7,7	0.89	0	9,9,9	0.97	0
3	OPE	С	824	1	7,7,7	0.88	0	$9,\!9,\!9$	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
3	OPE	N	824	1	-	0/5/5/5	-
3	OPE	G	824	1	-	0/5/5/5	-
3	OPE	J	824	1	-	0/5/5/5	-
3	OPE	0	824	1	-	0/5/5/5	-
3	OPE	Е	824	1	-	0/5/5/5	-
3	OPE	F	824	1	-	0/5/5/5	-
3	OPE	K	824	1	-	0/5/5/5	-
3	OPE	R	824	1	-	0/5/5/5	-
3	OPE	Р	824	1	-	0/5/5/5	-
3	OPE	D	824	1	-	0/5/5/5	-
3	OPE	В	824	1	-	0/5/5/5	-
3	OPE	А	824	1	-	0/5/5/5	-
3	OPE	Q	824	1	-	0/5/5/5	-
3	OPE	Ι	824	1	-	0/5/5/5	-
3	OPE	Н	824	1	-	0/5/5/5	-
3	OPE	М	824	1	-	0/5/5/5	-
3	OPE	L	824	1	-	0/5/5/5	-
3	OPE	С	824	1	-	0/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.



There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-1236. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 18



Y Index: 18



Z Index: 27

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 26



Y Index: 26



Z Index: 52

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 60 nm^3 ; this corresponds to an approximate mass of 55 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-1236 and PDB model 2HIL. Per-residue inclusion information can be found in section 3 on page 10.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 2.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2.05).



9.4 Atom inclusion (i)



At the recommended contour level, 14% of all backbone atoms, 12% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (2.05) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	$\mathbf{Q} extsf{-score}$
All	0.1216	0.0350
А	0.1746	0.0520
В	0.1779	0.0520
С	0.1763	0.0470
D	0.1671	0.0560
Е	0.1662	0.0530
F	0.1404	0.0400
G	0.1011	0.0250
Н	0.0643	0.0070
Ι	0.0084	-0.0050
J	0.1763	0.0520
K	0.1738	0.0480
L	0.1713	0.0470
М	0.1796	0.0520
N	0.1679	0.0500
0	0.1028	0.0310
Р	0.0668	0.0190
Q	0.0242	-0.0090
R	0.0008	-0.0050
S	0.0000	0.0700
Т	0.0000	0.0790
U	0.0000	0.0680
V	0.0000	0.0810
W	0.0000	0.0730
X	0.0000	0.0860
Y	0.0000	0.1040
Z	0.0357	0.0940
a	0.0000	0.0140
b	0.0000	0.0590
с	0.0000	0.0710
d	0.0000	0.0460
e	0.0000	0.0390
f	0.0000	0.0410
g	0.0000	0.0000
h	0.0000	0.0000



Continued from previous page...

Chain	Atom inclusion	Q-score
i	0.0000	0.0000
j	0.0000	0.0000

