



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 02:18 pm BST

PDB ID : 6HI1
Title : PI3 Kinase Delta in complex with 3[6(morpholin4yl)pyridin2yl]phenol
Authors : Convery, M.A.; Summers, D.; Peace, S.
Deposited on : 2018-08-29
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	940	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7473 atoms, of which 16 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	863	6944	4438	1191	1261	54	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	105	GLY	GLU	conflict	UNP O35904
A	508	GLN	-	insertion	UNP O35904

- Molecule 2 is 3-(6-morpholin-4-ylpyridin-2-yl)phenol (three-letter code: G5T) (formula: C₁₅H₁₆N₂O₂) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	35	15	16	2	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	494	Total 494	O 494	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	141.57Å 64.36Å 116.30Å 90.00° 103.24° 90.00°	Depositor
Resolution (Å)	30.31 – 2.07 54.02 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.31-2.07) 99.3 (54.02-2.07)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.07Å)	Xtrriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.211 , 0.254 0.224 , 0.271	Depositor DCC
R_{free} test set	3126 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	39.6	Xtrriage
Anisotropy	0.100	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 59.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7473	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G5T

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/7096	0.64	0/9582

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6944	0	6918	34	0
2	A	19	16	0	0	0
3	A	494	0	0	1	0
All	All	7457	16	6918	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:324:SER:HB3	1:A:376:GLU:HG3	1.83	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:859:LEU:HD21	1:A:901:ILE:HD11	1.86	0.57
1:A:209:THR:HB	1:A:257:CYS:HB3	1.87	0.56
1:A:752:MET:HB2	1:A:758:PRO:HG2	1.89	0.53
1:A:371:TRP:HB3	1:A:373:GLN:HG2	1.92	0.51
1:A:169:LEU:HD22	1:A:259:PHE:HE1	1.75	0.50
1:A:715:MET:O	1:A:719:MET:HG3	2.11	0.50
1:A:154:ARG:HD2	1:A:165:TYR:CE2	2.47	0.49
1:A:971:GLY:HA3	1:A:1004:LEU:HD21	1.96	0.48
1:A:834:ILE:HG23	1:A:855:LEU:HD21	1.96	0.48
1:A:419:ALA:HB1	1:A:441:MET:HB3	1.96	0.47
1:A:356:LYS:HG3	3:A:1375:HOH:O	2.13	0.47
1:A:342:VAL:HG22	1:A:362:GLU:HG2	1.96	0.47
1:A:917:GLY:HA2	1:A:930:VAL:HG23	1.97	0.47
1:A:503:ILE:HG22	1:A:504:THR:H	1.80	0.46
1:A:936:TYR:HD1	1:A:1030:LYS:HE3	1.81	0.46
1:A:154:ARG:HA	1:A:157:LEU:HG	1.99	0.45
1:A:614:GLN:HG3	1:A:981:MET:HG2	1.99	0.45
1:A:808:LEU:HD11	1:A:963:ALA:HB2	1.99	0.44
1:A:213:PRO:O	1:A:217:MET:HG3	2.17	0.44
1:A:385:PRO:HD2	1:A:388:ALA:HB2	1.98	0.44
1:A:395:TYR:HA	1:A:418:ILE:HG12	2.00	0.44
1:A:205:PHE:HB3	1:A:223:LYS:HE3	2.00	0.44
1:A:191:LEU:O	1:A:272:PRO:HD2	2.18	0.43
1:A:691:LYS:HG3	1:A:727:ALA:HB3	2.01	0.43
1:A:859:LEU:HD21	1:A:905:GLY:HA2	2.01	0.43
1:A:507:GLU:HB2	1:A:530:LEU:HD11	2.02	0.42
1:A:702:SER:HB3	1:A:711:THR:HG23	2.01	0.42
1:A:1003:ALA:HB1	1:A:1006:LYS:HD2	2.02	0.41
1:A:1027:TRP:CD1	1:A:1030:LYS:HD2	2.55	0.41
1:A:349:HIS:HB2	1:A:354:LEU:HD21	2.01	0.41
1:A:637:ALA:HB1	1:A:644:GLY:HA2	2.02	0.41
1:A:581:GLU:HB2	1:A:976:HIS:NE2	2.35	0.40
1:A:341:LEU:HG	1:A:365:VAL:HG22	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	849/940 (90%)	825 (97%)	21 (2%)	3 (0%)	34 25

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	229	ARG
1	A	911	ASP
1	A	328	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	760/827 (92%)	745 (98%)	15 (2%)	55 51

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	ARG
1	A	115	SER
1	A	145	GLN
1	A	270	LEU
1	A	325	ILE
1	A	368	GLU
1	A	394	LEU
1	A	416	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	560	ASP
1	A	684	LYS
1	A	696	ASN
1	A	787	ASP
1	A	895	HIS
1	A	915	PHE
1	A	995	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	349	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	G5T	A	1101	-	21,21,21	0.87	1 (4%)	28,28,28	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G5T	A	1101	-	-	0/8/16/16	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	G5T	C12-N13	2.13	1.37	1.34

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	863/940 (91%)	0.45	51 (5%) 22 23	26, 49, 85, 123	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	GLY	8.1
1	A	509	LEU	7.7
1	A	934	LEU	7.4
1	A	109	VAL	5.8
1	A	341	LEU	5.2
1	A	233	VAL	5.1
1	A	1017	VAL	4.9
1	A	294	PRO	4.8
1	A	395	TYR	4.1
1	A	365	VAL	4.0
1	A	370	VAL	3.7
1	A	232	LEU	3.6
1	A	1027	TRP	3.5
1	A	363	VAL	3.5
1	A	1032	ASN	3.3
1	A	330	GLY	3.2
1	A	1024	ARG	3.2
1	A	933	ILE	3.2
1	A	1013	LYS	3.0
1	A	342	VAL	2.9
1	A	398	VAL	2.9
1	A	393	ALA	2.8
1	A	445	VAL	2.8
1	A	210	LYS	2.8
1	A	492	LEU	2.7
1	A	847	THR	2.7
1	A	225	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	710	GLN	2.7
1	A	371	TRP	2.6
1	A	190	LEU	2.6
1	A	230	GLN	2.6
1	A	506	GLU	2.5
1	A	1021	GLU	2.5
1	A	108	ARG	2.4
1	A	525	GLU	2.4
1	A	176	ARG	2.3
1	A	503	ILE	2.3
1	A	992	SER	2.3
1	A	337	GLU	2.3
1	A	372	LYS	2.2
1	A	227	VAL	2.2
1	A	231	PRO	2.2
1	A	228	PHE	2.2
1	A	470	ALA	2.2
1	A	474	TYR	2.2
1	A	526	HIS	2.1
1	A	113	ILE	2.1
1	A	1028	LYS	2.1
1	A	204	THR	2.0
1	A	483	VAL	2.0
1	A	521	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	G5T	A	1101	19/19	0.91	0.14	29,37,41,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.