



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:26 pm BST

PDB ID : 2HJG
Title : The crystal structure of the *B. subtilis* YphC GTPase in complex with GDP
Authors : Muench, S.P.; Xu, L.; Sedelnikova, S.E.; Rice, D.W.
Deposited on : 2006-06-30
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3235 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding protein engA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	400	3147	2018	527	591	1	10	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	80	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	89	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	126	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	195	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	233	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	306	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	328	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	340	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	364	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	369	MSE	MET	MODIFIED RESIDUE	UNP P50743
A	405	MSE	MET	MODIFIED RESIDUE	UNP P50743

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GTP-binding protein engA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.71Å 65.05Å 110.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.96 – 2.50 19.90 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.96-2.50) 100.0 (19.90-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.15 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.274 0.250 , 0.289	Depositor DCC
R_{free} test set	814 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.032 for k,h,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3235	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/3196	0.69	3/4305 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	131	TYR	O-C-N	7.49	134.68	122.70
1	A	131	TYR	CA-C-N	-5.79	104.45	117.20
1	A	402	PRO	O-C-N	-5.40	114.06	122.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	122	ASP	Peptide
1	A	435	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3147	0	3124	50	0
2	A	1	0	0	0	0
3	A	56	0	24	2	0
4	A	31	0	0	0	0
All	All	3235	0	3148	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:TYR:CD1	1:A:131:TYR:N	2.16	1.06
1:A:130:ILE:HD11	1:A:141:PRO:CB	1.87	1.04
1:A:131:TYR:N	1:A:131:TYR:HD1	1.53	1.01
1:A:130:ILE:HD11	1:A:141:PRO:HB2	1.01	0.99
1:A:130:ILE:CD1	1:A:141:PRO:HB2	1.92	0.98
1:A:131:TYR:H	1:A:131:TYR:HD1	0.94	0.92
1:A:328:MSE:HE1	1:A:339:LEU:HD22	1.54	0.88
1:A:195:MSE:CE	1:A:339:LEU:HD23	2.08	0.82
1:A:191:LEU:HD22	1:A:328:MSE:HE3	1.66	0.77
1:A:191:LEU:CD2	1:A:328:MSE:HE3	2.15	0.75
1:A:292:VAL:CG1	1:A:328:MSE:HE2	2.20	0.72
1:A:292:VAL:HG11	1:A:328:MSE:HE2	1.72	0.71
1:A:435:ARG:HH11	1:A:435:ARG:HG2	1.58	0.67
1:A:294:ASN:HD21	1:A:330:ALA:H	1.43	0.67
1:A:195:MSE:HE2	1:A:339:LEU:HD23	1.80	0.63
1:A:360:ASN:HD21	1:A:385:ALA:H	1.48	0.60
1:A:375:HIS:CD2	1:A:376:ASN:ND2	2.74	0.56
1:A:180:LEU:HD13	1:A:262:ALA:HB3	1.87	0.56
1:A:231:ALA:O	1:A:235:LYS:HD3	2.06	0.55
1:A:397:VAL:HG21	1:A:405:MSE:HE1	1.87	0.55
1:A:3:LYS:HB3	1:A:4:PRO:HD3	1.87	0.55
1:A:195:MSE:CE	1:A:339:LEU:CD2	2.83	0.54
1:A:360:ASN:O	1:A:364:MSE:HG2	2.08	0.54
1:A:360:ASN:HD21	1:A:385:ALA:N	2.07	0.53
1:A:295:LYS:NZ	3:A:500:GDP:O4'	2.42	0.52
1:A:44:SER:HA	1:A:54:ASN:HD22	1.74	0.52
1:A:292:VAL:HG13	1:A:328:MSE:HE2	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:ASP:OD1	1:A:303:GLU:N	2.44	0.49
1:A:330:ALA:HA	1:A:336:ILE:HD11	1.95	0.49
1:A:122:ASP:O	1:A:123:ASN:HB2	2.15	0.46
1:A:340:MSE:HB2	1:A:341:PRO:HD3	1.98	0.46
1:A:131:TYR:HB2	1:A:132:ASP:H	1.29	0.46
1:A:146:GLY:N	3:A:800:GDP:O6	2.49	0.46
1:A:435:ARG:CG	1:A:435:ARG:HH11	2.28	0.45
1:A:235:LYS:HD2	1:A:238:LYS:HZ2	1.82	0.45
1:A:401:ASP:HB3	1:A:404:LEU:HG	1.99	0.44
1:A:191:LEU:HD21	1:A:328:MSE:HE3	1.97	0.44
1:A:49:LEU:HD21	1:A:156:LEU:HD13	2.00	0.44
1:A:354:VAL:HG13	1:A:421:PHE:CE2	2.53	0.44
1:A:130:ILE:HG23	1:A:130:ILE:O	2.18	0.43
1:A:49:LEU:CD2	1:A:156:LEU:HD13	2.48	0.43
1:A:328:MSE:HE1	1:A:339:LEU:CD2	2.38	0.42
1:A:280:GLY:O	1:A:284:GLU:HG2	2.19	0.42
1:A:195:MSE:HE2	1:A:339:LEU:CD2	2.48	0.42
1:A:276:LYS:HE2	1:A:316:HIS:HB3	2.02	0.42
1:A:381:LYS:HB2	1:A:400:ASN:HB2	2.03	0.41
1:A:6:VAL:HB	1:A:55:LEU:HD13	2.02	0.41
1:A:186:VAL:HG21	1:A:264:VAL:CG2	2.51	0.40
1:A:384:TYR:HD2	1:A:398:PHE:CD1	2.39	0.40
1:A:414:GLU:HG2	1:A:428:ILE:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	390/436 (89%)	376 (96%)	13 (3%)	1 (0%)	41 61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	135	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	336/359 (94%)	323 (96%)	13 (4%)	32 57

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	ARG
1	A	93	ARG
1	A	130	ILE
1	A	131	TYR
1	A	148	HIS
1	A	173	GLU
1	A	180	LEU
1	A	214	VAL
1	A	235	LYS
1	A	301	LYS
1	A	333	LYS
1	A	353	ARG
1	A	436	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	ASN
1	A	75	GLN
1	A	172	ASN
1	A	294	ASN
1	A	360	ASN
1	A	376	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GDP	A	800	-	24,30,30	1.13	2 (8%)	31,47,47	1.84	7 (22%)
3	GDP	A	500	-	24,30,30	1.15	2 (8%)	31,47,47	1.89	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	800	-	-	5/12/32/32	0/3/3/3
3	GDP	A	500	-	-	5/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	800	GDP	C6-C5	4.03	1.48	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	GDP	C6-C5	3.96	1.48	1.41
3	A	500	GDP	C5-C4	2.36	1.47	1.40
3	A	800	GDP	C5-C4	2.21	1.46	1.40

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	800	GDP	C2-N3-C4	4.89	120.94	115.36
3	A	500	GDP	C6-C5-C4	-4.56	116.44	120.80
3	A	500	GDP	C6-N1-C2	4.49	123.07	115.93
3	A	800	GDP	C6-C5-C4	-4.06	116.92	120.80
3	A	500	GDP	C5-C6-N1	-3.90	118.10	123.43
3	A	800	GDP	C6-N1-C2	3.86	122.07	115.93
3	A	500	GDP	C2-N3-C4	3.76	119.66	115.36
3	A	800	GDP	C5-C6-N1	-3.68	118.39	123.43
3	A	500	GDP	N3-C2-N1	-3.50	122.55	127.22
3	A	500	GDP	PA-O3A-PB	-3.39	121.19	132.83
3	A	800	GDP	N3-C2-N1	-3.37	122.73	127.22
3	A	800	GDP	PA-O3A-PB	-2.76	123.34	132.83
3	A	800	GDP	C4-C5-N7	-2.51	106.79	109.40

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	800	GDP	C5'-O5'-PA-O3A
3	A	800	GDP	C5'-O5'-PA-O2A
3	A	500	GDP	PA-O3A-PB-O3B
3	A	500	GDP	C5'-O5'-PA-O3A
3	A	800	GDP	O4'-C4'-C5'-O5'
3	A	800	GDP	C3'-C4'-C5'-O5'
3	A	500	GDP	PA-O3A-PB-O2B
3	A	800	GDP	C5'-O5'-PA-O1A
3	A	500	GDP	PA-O3A-PB-O1B
3	A	500	GDP	O4'-C4'-C5'-O5'

There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	800	GDP	1	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	500	GDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/436 (89%)	0.84	54 (13%) 2 2	31, 42, 51, 61	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	123	ASN	6.3
1	A	68	PHE	6.0
1	A	29	SER	5.9
1	A	130	ILE	5.3
1	A	28	ILE	4.5
1	A	122	ASP	4.5
1	A	148	HIS	4.3
1	A	264	VAL	4.2
1	A	71	GLN	4.2
1	A	291	ILE	3.8
1	A	265	LEU	3.6
1	A	300	ASP	3.5
1	A	67	PRO	3.4
1	A	436	LYS	3.4
1	A	234	ARG	3.3
1	A	302	ASP	3.2
1	A	93	ARG	3.2
1	A	129	ASN	3.2
1	A	292	VAL	3.1
1	A	117	ALA	3.1
1	A	131	TYR	3.1
1	A	293	VAL	3.1
1	A	168	GLU	3.0
1	A	237	GLY	3.0
1	A	199	GLU	3.0
1	A	96	VAL	2.7
1	A	303	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	13	ASN	2.6
1	A	135	SER	2.6
1	A	81	ASP	2.6
1	A	75	GLN	2.5
1	A	69	LEU	2.5
1	A	111	LYS	2.4
1	A	435	ARG	2.4
1	A	242	THR	2.4
1	A	204	SER	2.4
1	A	315	ASP	2.3
1	A	262	ALA	2.3
1	A	244	GLU	2.3
1	A	263	VAL	2.3
1	A	241	GLU	2.3
1	A	146	GLY	2.3
1	A	363	ILE	2.3
1	A	272	ILE	2.2
1	A	417	ILE	2.2
1	A	42	ILE	2.2
1	A	378	SER	2.2
1	A	27	ARG	2.2
1	A	90	VAL	2.2
1	A	98	ALA	2.1
1	A	376	ASN	2.1
1	A	147	THR	2.1
1	A	178	PHE	2.1
1	A	235	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GDP	A	800	28/28	0.88	0.21	41,48,51,52	0
3	GDP	A	500	28/28	0.95	0.13	32,34,42,42	0
2	ZN	A	900	1/1	0.96	0.06	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.