



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 9, 2023 – 10:33 PM EDT

PDB ID : 4HK5  
Title : Crystal structure of Cordyceps militaris IDCase in apo form  
Authors : Xu, S.; Zhu, J.; Ding, J.  
Deposited on : 2012-10-15  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	 7% 85% 6% • 7%
1	B	387	 7% 88% 5% • 6%
1	C	387	 5% 91% 5% •
1	D	387	 2% 93% 5% •

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 12209 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uracil-5-carboxylate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	358	2742	1769	469	490	14	0	0	0
1	B	362	2781	1790	480	497	14	0	0	0
1	C	370	2826	1819	483	510	14	0	0	0
1	D	380	2924	1879	507	524	14	0	1	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP G3J531
A	-1	GLY	-	expression tag	UNP G3J531
A	0	SER	-	expression tag	UNP G3J531
A	377	LEU	-	expression tag	UNP G3J531
A	378	GLU	-	expression tag	UNP G3J531
A	379	HIS	-	expression tag	UNP G3J531
A	380	HIS	-	expression tag	UNP G3J531
A	381	HIS	-	expression tag	UNP G3J531
A	382	HIS	-	expression tag	UNP G3J531
A	383	HIS	-	expression tag	UNP G3J531
A	384	HIS	-	expression tag	UNP G3J531
B	-2	MET	-	expression tag	UNP G3J531
B	-1	GLY	-	expression tag	UNP G3J531
B	0	SER	-	expression tag	UNP G3J531
B	377	LEU	-	expression tag	UNP G3J531
B	378	GLU	-	expression tag	UNP G3J531
B	379	HIS	-	expression tag	UNP G3J531
B	380	HIS	-	expression tag	UNP G3J531
B	381	HIS	-	expression tag	UNP G3J531
B	382	HIS	-	expression tag	UNP G3J531
B	383	HIS	-	expression tag	UNP G3J531

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	384	HIS	-	expression tag	UNP G3J531
C	-2	MET	-	expression tag	UNP G3J531
C	-1	GLY	-	expression tag	UNP G3J531
C	0	SER	-	expression tag	UNP G3J531
C	377	LEU	-	expression tag	UNP G3J531
C	378	GLU	-	expression tag	UNP G3J531
C	379	HIS	-	expression tag	UNP G3J531
C	380	HIS	-	expression tag	UNP G3J531
C	381	HIS	-	expression tag	UNP G3J531
C	382	HIS	-	expression tag	UNP G3J531
C	383	HIS	-	expression tag	UNP G3J531
C	384	HIS	-	expression tag	UNP G3J531
D	-2	MET	-	expression tag	UNP G3J531
D	-1	GLY	-	expression tag	UNP G3J531
D	0	SER	-	expression tag	UNP G3J531
D	377	LEU	-	expression tag	UNP G3J531
D	378	GLU	-	expression tag	UNP G3J531
D	379	HIS	-	expression tag	UNP G3J531
D	380	HIS	-	expression tag	UNP G3J531
D	381	HIS	-	expression tag	UNP G3J531
D	382	HIS	-	expression tag	UNP G3J531
D	383	HIS	-	expression tag	UNP G3J531
D	384	HIS	-	expression tag	UNP G3J531

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	2	Total Zn 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	213	Total O 213 213	0	0

Continued on next page...


*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	193	Total 193	O 193	0	0
3	C	234	Total 234	O 234	0	0
3	D	291	Total 291	O 291	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uracil-5-carboxylate decarboxylase

Chain A: 





- Molecule 1: Uracil-5-carboxylate decarboxylase

Chain B: 




- Molecule 1: Uracil-5-carboxylate decarboxylase

Chain C: 



- Molecule 1: Uracil-5-carboxylate decarboxylase

Chain D: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.65Å 146.16Å 77.77Å 90.00° 96.52° 90.00°	Depositor
Resolution (Å)	77.27 – 1.90 48.77 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.3 (77.27-1.90) 99.3 (48.77-1.90)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.77 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.157 , 0.197 0.157 , 0.197	Depositor DCC
$R_{free}$ test set	6731 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtrriage
Anisotropy	0.366	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 48.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.015 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12209	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	2/2810 (0.1%)	0.53	0/3821
1	B	0.46	0/2853	0.50	0/3880
1	C	0.45	1/2898 (0.0%)	0.52	0/3948
1	D	0.48	2/3006 (0.1%)	0.53	0/4095
All	All	0.46	5/11567 (0.0%)	0.52	0/15744

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	338	TRP	CD2-CE2	5.28	1.47	1.41
1	D	287	TRP	CD2-CE2	5.17	1.47	1.41
1	A	338	TRP	CD2-CE2	5.08	1.47	1.41
1	C	287	TRP	CD2-CE2	5.08	1.47	1.41
1	A	287	TRP	CD2-CE2	5.08	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2742	0	2768	16	0
1	B	2781	0	2786	12	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2826	0	2844	6	0
1	D	2924	0	2921	8	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
3	A	213	0	0	1	0
3	B	193	0	0	2	0
3	C	234	0	0	0	0
3	D	291	0	0	1	0
All	All	12209	0	11319	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:LEU:HD13	1:A:34:ARG:HE	1.37	0.87
1:B:31:PRO:HG3	1:B:99:PRO:HG3	1.62	0.78
1:A:73:HIS:HD2	3:A:661:HOH:O	1.78	0.66
1:A:32:LEU:HD13	1:A:34:ARG:NE	2.13	0.62
1:D:380[A]:HIS:NE2	3:D:741:HOH:O	2.31	0.60
1:B:281:LYS:H	1:B:281:LYS:NZ	1.99	0.59
1:A:139:LEU:HD21	1:A:164:LEU:HD12	1.84	0.58
1:B:73:HIS:HD2	3:B:627:HOH:O	1.88	0.56
1:B:281:LYS:H	1:B:281:LYS:HZ2	1.56	0.54
1:D:114:ASP:HB2	1:D:152:ARG:NH2	2.23	0.54
1:A:139:LEU:HD21	1:A:164:LEU:CD1	2.37	0.54
1:B:209:SER:HB2	3:B:644:HOH:O	2.10	0.52
1:A:166:THR:OG1	1:A:225:GLU:OE1	2.25	0.51
1:D:45:ILE:HD13	1:D:54:LEU:HD13	1.92	0.51
1:D:31:PRO:HG3	1:D:99:PRO:HG3	1.93	0.50
1:D:128:VAL:HG12	1:D:380[B]:HIS:CE1	2.46	0.50
1:B:264:GLU:HG2	1:B:286:ILE:HG22	1.93	0.50
1:A:45:ILE:HD13	1:A:54:LEU:HD13	1.95	0.48
1:A:16:TYR:OH	1:A:46:LEU:HD21	2.14	0.48
1:A:101:PHE:HB3	1:A:104:LEU:HD12	1.95	0.47
1:A:23:MET:SD	1:A:104:LEU:HD21	2.55	0.47
1:C:264:GLU:HG2	1:C:286:ILE:HG22	1.95	0.47
1:D:5:THR:HB	1:D:6:PRO:HD3	1.95	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:VAL:HG23	1:A:90:ARG:HB3	1.97	0.46
1:B:16:TYR:CE1	1:B:97:ALA:HB3	2.52	0.45
1:B:23:MET:HE1	1:B:104:LEU:HD21	1.97	0.45
1:B:297:ASP:OD1	1:B:297:ASP:C	2.55	0.45
1:A:14:HIS:HB3	1:A:97:ALA:HB2	1.99	0.44
1:C:101:PHE:HB3	1:C:104:LEU:HD12	1.98	0.44
1:C:45:ILE:HD13	1:C:54:LEU:HD13	2.01	0.43
1:A:99:PRO:HB3	1:A:197:HIS:NE2	2.33	0.43
1:A:16:TYR:CE1	1:A:97:ALA:HB3	2.54	0.42
1:C:10:ASP:OD1	1:C:321:GLY:HA2	2.20	0.42
1:C:5:THR:HA	1:C:6:PRO:HD3	1.89	0.42
1:A:139:LEU:CD2	1:A:164:LEU:HD12	2.49	0.41
1:A:240:ASP:OD1	1:A:284:ARG:HD3	2.20	0.41
1:C:16:TYR:CE1	1:C:97:ALA:HB3	2.55	0.41
1:D:264:GLU:OE1	1:D:283:ARG:NH2	2.52	0.41
1:D:297:ASP:C	1:D:297:ASP:OD1	2.59	0.41
1:B:72:THR:O	1:B:76:SER:HB3	2.21	0.40
1:B:21:ILE:HD12	1:B:70:LEU:HD21	2.02	0.40
1:B:185:VAL:HG13	1:B:190:LEU:HB2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	352/387 (91%)	343 (97%)	7 (2%)	2 (1%)	25 15
1	B	356/387 (92%)	346 (97%)	8 (2%)	2 (1%)	25 15
1	C	368/387 (95%)	359 (98%)	7 (2%)	2 (0%)	29 18
1	D	379/387 (98%)	371 (98%)	6 (2%)	2 (0%)	29 18
All	All	1455/1548 (94%)	1419 (98%)	28 (2%)	8 (0%)	25 18

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	197	HIS
1	D	197	HIS
1	A	197	HIS
1	B	197	HIS
1	A	364	GLY
1	B	364	GLY
1	D	364	GLY
1	C	364	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/313 (93%)	285 (98%)	7 (2%)	49	43
1	B	296/313 (95%)	294 (99%)	2 (1%)	84	84
1	C	300/313 (96%)	297 (99%)	3 (1%)	76	76
1	D	310/313 (99%)	308 (99%)	2 (1%)	86	87
All	All	1198/1252 (96%)	1184 (99%)	14 (1%)	71	70

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LYS
1	A	46	LEU
1	A	47	LEU
1	A	90	ARG
1	A	139	LEU
1	A	189	LYS
1	A	247	MET
1	B	21	ILE
1	B	281	LYS
1	C	40	ASP
1	C	243	ARG
1	C	374	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	154	LYS
1	D	281	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	B	127	HIS
1	B	382	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	358/387 (92%)	0.09	27 (7%) 14 15	9, 19, 46, 69	0
1	B	362/387 (93%)	0.08	26 (7%) 15 17	10, 21, 55, 79	0
1	C	370/387 (95%)	-0.24	18 (4%) 29 33	8, 16, 37, 83	0
1	D	380/387 (98%)	-0.20	8 (2%) 63 66	9, 16, 35, 58	0
All	All	1470/1548 (94%)	-0.07	79 (5%) 25 29	8, 18, 46, 83	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	21	ILE	8.8
1	A	35	THR	6.4
1	A	34	ARG	6.4
1	A	44	LEU	6.3
1	A	57	ALA	6.1
1	B	28	GLN	6.0
1	A	56	ALA	5.8
1	C	39	ALA	5.5
1	B	56	ALA	5.4
1	A	52	ALA	5.3
1	A	28	GLN	5.2
1	B	51	LEU	5.1
1	B	22	ALA	4.9
1	B	24	LEU	4.6
1	D	5	THR	4.6
1	B	66	PRO	4.4
1	B	52	ALA	4.4
1	A	36	PHE	4.2
1	B	55	ASP	4.2
1	B	42	PRO	4.0
1	C	62	ALA	4.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	25	GLU	4.0
1	B	383	HIS	4.0
1	B	65	LEU	3.9
1	C	40	ASP	3.8
1	C	38	GLN	3.8
1	A	53	ALA	3.8
1	A	55	ASP	3.7
1	C	333	ASP	3.7
1	D	62	ALA	3.6
1	B	29	THR	3.4
1	A	65	LEU	3.4
1	B	46	LEU	3.3
1	B	31	PRO	3.3
1	B	6	PRO	3.3
1	A	58	LEU	3.3
1	A	374	LYS	3.1
1	C	334	VAL	3.1
1	C	374	LYS	3.1
1	A	51	LEU	3.0
1	C	60	ASP	3.0
1	B	50	GLU	2.9
1	B	26	LYS	2.9
1	A	48	SER	2.9
1	B	53	ALA	2.9
1	B	44	LEU	2.8
1	C	59	ALA	2.8
1	C	5	THR	2.7
1	B	69	PRO	2.7
1	A	54	LEU	2.7
1	B	47	LEU	2.7
1	C	332	GLU	2.7
1	C	336	GLY	2.7
1	C	37	PRO	2.7
1	A	30	ILE	2.6
1	A	31	PRO	2.6
1	B	70	LEU	2.5
1	D	377	LEU	2.5
1	A	69	PRO	2.5
1	D	381	HIS	2.5
1	B	33	VAL	2.5
1	C	58	LEU	2.5
1	A	29	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	382	HIS	2.4
1	B	107	ASP	2.3
1	A	24	LEU	2.3
1	C	63	ALA	2.3
1	C	36	PHE	2.3
1	C	335	GLN	2.3
1	A	6	PRO	2.3
1	A	70	LEU	2.3
1	D	39	ALA	2.2
1	D	378	GLU	2.2
1	C	331	GLU	2.2
1	A	49	SER	2.2
1	A	332	GLU	2.2
1	B	45	ILE	2.1
1	A	281	LYS	2.1
1	D	59	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	401	1/1	1.00	0.05	17,17,17,17	0
2	ZN	B	401	1/1	1.00	0.04	21,21,21,21	0
2	ZN	C	401	1/1	1.00	0.04	15,15,15,15	0
2	ZN	D	401	1/1	1.00	0.04	16,16,16,16	0
2	ZN	D	402	1/1	1.00	0.03	25,25,25,25	0



## 6.5 Other polymers [i](#)

There are no such residues in this entry.