



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2023 – 11:50 AM EDT

PDB ID : 3HN1
Title : Crystal structure of HCR/T complexed with GT2 and lactose
Authors : Chen, C.; Fu, Z.; Kim, J.-J.P.; Barbieri, J.T.; Baldwin, M.R.
Deposited on : 2009-05-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	 6% 80% 16% ..
2	B	2	 100%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3834 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetanus toxin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	442	3575	2289	597	679	10	0	0	0

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-8)-N-acetyl-alpha-neuraminic acid.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	40	22	2	16	0	0	0

- Molecule 3 is alpha-D-galactopyranose (three-letter code: GLA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	192	Total	O	0	0
			192	192		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.97Å 72.68Å 119.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.53 – 2.10 25.53 – 2.10	Depositor EDS
% Data completeness (in resolution range)	93.5 (25.53-2.10) 93.5 (25.53-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 2.10Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.216 , 0.257 0.203 , 0.244	Depositor DCC
R_{free} test set	3245 reflections (9.98%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtrriage
Anisotropy	0.525	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3834	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, SO4, GLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/3655	0.62	0/4958

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3575	0	3512	60	0
2	B	40	0	33	0	0
3	A	12	0	12	1	0
4	A	15	0	0	0	0
5	A	192	0	0	3	0
All	All	3834	0	3557	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1097:ASN:HD21	1:A:1100:GLU:HG3	1.35	0.91
1:A:1097:ASN:ND2	1:A:1100:GLU:HG3	1.97	0.78
1:A:1178:LYS:HE2	5:A:1488:HOH:O	1.84	0.75
1:A:949:ASN:ND2	1:A:1040:ARG:H	1.86	0.73
1:A:1141:GLN:HE21	1:A:1142:LEU:H	1.36	0.72
1:A:972:ASN:ND2	1:A:1078:ASN:H	1.92	0.68
1:A:1183:ASN:HD21	1:A:1186:ILE:HA	1.58	0.68
1:A:972:ASN:HD21	1:A:1078:ASN:H	1.40	0.68
1:A:1097:ASN:ND2	1:A:1100:GLU:H	1.91	0.68
1:A:986:SER:O	1:A:987:ILE:HG22	1.94	0.67
1:A:1297:LYS:HG3	1:A:1298:ILE:HG12	1.76	0.67
1:A:1141:GLN:NE2	1:A:1142:LEU:H	1.95	0.65
1:A:1171:ASN:HB2	5:A:1532:HOH:O	1.98	0.62
1:A:1211:TYR:CE2	1:A:1223:ARG:HG2	2.36	0.60
1:A:949:ASN:HD21	1:A:1040:ARG:H	1.49	0.60
1:A:1241:GLU:HG2	1:A:1243:VAL:HG13	1.83	0.59
1:A:1112:THR:HG1	1:A:1113:PHE:HD1	1.50	0.59
1:A:1183:ASN:HD22	1:A:1183:ASN:H	1.50	0.58
1:A:1184:ASN:ND2	1:A:1185:GLU:HG3	2.20	0.57
1:A:1097:ASN:HD22	1:A:1097:ASN:C	2.09	0.56
1:A:1224:ILE:HA	1:A:1285:ILE:HG22	1.87	0.56
1:A:867:LEU:HD12	1:A:1040:ARG:HB2	1.88	0.56
1:A:869:CYS:O	1:A:870:TRP:HB3	2.06	0.55
1:A:1241:GLU:HG2	1:A:1243:VAL:CG1	2.36	0.55
1:A:1170:TYR:CE1	1:A:1308:THR:HA	2.42	0.54
1:A:876:ASP:HB3	1:A:879:VAL:HG23	1.90	0.54
1:A:1083:TYR:OH	1:A:1310:GLU:HG3	2.07	0.54
1:A:1213:LYS:HB2	1:A:1235:PRO:HG2	1.89	0.54
1:A:1022:LYS:HB2	1:A:1166:TYR:HB3	1.90	0.54
1:A:1183:ASN:HD22	1:A:1183:ASN:N	2.05	0.53
1:A:1297:LYS:HG3	1:A:1298:ILE:N	2.25	0.52
1:A:1183:ASN:O	1:A:1184:ASN:C	2.49	0.50
1:A:1002:TRP:CE3	1:A:1036:ILE:HD13	2.47	0.49
1:A:1023:PHE:CE2	1:A:1136:SER:HB2	2.47	0.49
1:A:1178:LYS:HD2	1:A:1199:TYR:CE1	2.47	0.49
1:A:1111:ILE:HG12	1:A:1248:LEU:HD13	1.94	0.49
1:A:1018:ASP:OD2	1:A:1025:ALA:HB1	2.13	0.49
1:A:1183:ASN:ND2	1:A:1186:ILE:HA	2.27	0.48
1:A:1270:THR:HG22	1:A:1284:LEU:HD22	1.95	0.48
1:A:1111:ILE:HG12	1:A:1248:LEU:CD1	2.44	0.46
1:A:991:TRP:HB3	1:A:1004:LEU:HD23	1.97	0.46
1:A:1183:ASN:H	1:A:1183:ASN:ND2	2.13	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1290:TYR:OH	3:A:7:GLA:H3	2.16	0.45
1:A:957:ARG:HB3	1:A:1085:SER:HB2	1.98	0.45
1:A:1097:ASN:HD22	1:A:1100:GLU:H	1.59	0.45
1:A:1111:ILE:O	1:A:1111:ILE:HG22	2.17	0.45
1:A:1260:ASP:OD1	1:A:1261:LYS:HD3	2.16	0.45
1:A:1293:HIS:HE1	5:A:1547:HOH:O	2.01	0.44
1:A:909:TYR:HB3	1:A:910:PRO:HD2	2.00	0.43
1:A:949:ASN:HD22	1:A:1040:ARG:HG2	1.83	0.43
1:A:972:ASN:HD21	1:A:1077:CYS:HA	1.83	0.43
1:A:1179:ARG:NH1	1:A:1183:ASN:OD1	2.52	0.43
1:A:1261:LYS:HE3	1:A:1261:LYS:HB2	1.90	0.43
1:A:937:LYS:HD3	1:A:1068:ASN:O	2.19	0.43
1:A:947:PHE:CE1	1:A:1063:ALA:HB2	2.54	0.42
1:A:1181:THR:HB	1:A:1182:PRO:HD2	2.01	0.42
1:A:1312:TRP:CD1	1:A:1312:TRP:C	2.93	0.42
1:A:1046:LEU:C	1:A:1046:LEU:HD23	2.40	0.42
1:A:1240:MET:HA	1:A:1256:LYS:O	2.20	0.41
1:A:1253:VAL:HG12	1:A:1254:GLN:N	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	436/450 (97%)	401 (92%)	33 (8%)	2 (0%)	29 26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1184	ASN
1	A	1111	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	398/406 (98%)	385 (97%)	13 (3%)	38 40

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	944	ASN
1	A	1022	LYS
1	A	1029	ASN
1	A	1097	ASN
1	A	1104	LEU
1	A	1122	LEU
1	A	1167	ARG
1	A	1183	ASN
1	A	1184	ASN
1	A	1204	ASN
1	A	1245	LEU
1	A	1261	LYS
1	A	1284	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	903	ASN
1	A	949	ASN
1	A	972	ASN
1	A	998	ASN
1	A	1029	ASN
1	A	1068	ASN
1	A	1080	ASN
1	A	1097	ASN
1	A	1141	GLN
1	A	1183	ASN
1	A	1184	ASN
1	A	1204	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1293	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SIA	B	1	2	20,20,21	1.83	6 (30%)	24,28,31	2.08	8 (33%)
2	SIA	B	2	2	20,20,21	2.13	7 (35%)	24,28,31	2.05	7 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SIA	B	1	2	-	11/18/34/38	0/1/1/1
2	SIA	B	2	2	-	4/18/34/38	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	SIA	C5-N5	4.70	1.53	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	SIA	C2-C1	4.15	1.56	1.52
2	B	1	SIA	C4-C5	3.70	1.56	1.53
2	B	1	SIA	C5-N5	3.68	1.51	1.45
2	B	2	SIA	O6-C2	3.50	1.48	1.43
2	B	2	SIA	O1B-C1	-3.04	1.20	1.30
2	B	2	SIA	C4-C5	3.02	1.55	1.53
2	B	1	SIA	O1B-C1	-2.92	1.21	1.30
2	B	1	SIA	O6-C2	2.86	1.47	1.43
2	B	1	SIA	C2-C1	2.43	1.54	1.52
2	B	2	SIA	C3-C2	2.34	1.56	1.52
2	B	2	SIA	C8-C7	2.08	1.57	1.53
2	B	1	SIA	C3-C2	2.00	1.55	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	SIA	C6-O6-C2	4.74	121.48	111.34
2	B	1	SIA	C6-O6-C2	4.59	121.15	111.34
2	B	1	SIA	C11-C10-N5	4.53	123.77	116.10
2	B	2	SIA	C11-C10-N5	4.48	123.68	116.10
2	B	2	SIA	C9-C8-C7	3.32	119.61	112.41
2	B	1	SIA	O8-C8-C9	-3.25	101.52	109.14
2	B	1	SIA	O4-C4-C3	-2.65	103.36	109.94
2	B	1	SIA	C3-C4-C5	2.51	114.49	111.46
2	B	2	SIA	O4-C4-C3	-2.47	103.82	109.94
2	B	2	SIA	O8-C8-C9	-2.45	103.40	109.14
2	B	1	SIA	O10-C10-C11	-2.24	117.89	122.06
2	B	2	SIA	O10-C10-N5	-2.23	117.86	121.95
2	B	2	SIA	C3-C4-C5	2.22	114.14	111.46
2	B	1	SIA	C8-C7-C6	-2.13	109.00	113.03
2	B	1	SIA	O4-C4-C5	2.07	114.53	109.77

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	SIA	C5-C6-C7-C8
2	B	1	SIA	C5-C6-C7-O7
2	B	1	SIA	O6-C6-C7-C8
2	B	1	SIA	O6-C6-C7-O7
2	B	1	SIA	C11-C10-N5-C5
2	B	1	SIA	O10-C10-N5-C5

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	2	SIA	C11-C10-N5-C5
2	B	2	SIA	O10-C10-N5-C5
2	B	2	SIA	O8-C8-C9-O9
2	B	2	SIA	C7-C8-C9-O9
2	B	1	SIA	O8-C8-C9-O9
2	B	1	SIA	O1A-C1-C2-C3
2	B	1	SIA	O1A-C1-C2-O6
2	B	1	SIA	C7-C8-C9-O9
2	B	1	SIA	O1B-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	SO4	A	1403	-	4,4,4	0.19	0	6,6,6	0.19	0
4	SO4	A	1402	-	4,4,4	0.15	0	6,6,6	0.13	0
3	GLA	A	7	-	12,12,12	1.59	4 (33%)	17,17,17	1.02	1 (5%)
4	SO4	A	1401	-	4,4,4	0.16	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLA	A	7	-	-	1/2/22/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	7	GLA	C1-C2	2.88	1.59	1.52
3	A	7	GLA	C4-C3	2.41	1.58	1.52
3	A	7	GLA	O5-C1	2.07	1.48	1.42
3	A	7	GLA	O5-C5	2.02	1.49	1.44

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	7	GLA	C4-C3-C2	-2.31	106.78	110.82

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	7	GLA	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	7	GLA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/450 (98%)	0.17	29 (6%) 18 23	16, 26, 44, 56	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	870	TRP	8.7
1	A	986	SER	5.8
1	A	866	ASN	5.7
1	A	1111	ILE	5.3
1	A	1113	PHE	4.6
1	A	1184	ASN	4.4
1	A	943	TYR	4.4
1	A	1183	ASN	4.1
1	A	939	MET	3.9
1	A	987	ILE	3.7
1	A	869	CYS	3.2
1	A	1185	GLU	3.1
1	A	1112	THR	3.0
1	A	940	ASP	2.8
1	A	981	LYS	2.7
1	A	1110	SER	2.6
1	A	944	ASN	2.6
1	A	1076	ARG	2.5
1	A	1161	LYS	2.3
1	A	1233	GLY	2.2
1	A	1091	ILE	2.2
1	A	1068	ASN	2.2
1	A	875	GLU	2.2
1	A	1232	PRO	2.2
1	A	867	LEU	2.1
1	A	1202	TYR	2.1
1	A	952	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1297	LYS	2.1
1	A	876	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SIA	B	1	20/21	0.61	0.34	48,54,57,58	0
2	SIA	B	2	20/21	0.78	0.29	44,49,52,54	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GLA	A	7	12/12	0.59	0.32	40,43,44,45	0
4	SO4	A	1403	5/5	0.81	0.29	73,73,74,74	0
4	SO4	A	1401	5/5	0.91	0.21	63,64,65,65	0
4	SO4	A	1402	5/5	0.98	0.08	36,37,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.