

Full wwPDB X-ray Structure Validation Report (i)

Aug 10, 2020 – 10:39 AM BST

PDB ID	:	3HN3
Title	:	Human beta-glucuronidase at 1.7 A resolution
Authors	:	Klei, H.E.; Ghosh, K.; Anumula, R.
Deposited on	:	2009-05-29
Resolution	:	1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	4695(1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	613	4% 91%	8% •
1	В	613	^{3%} 93%	6% ·
1	D	613	4% 92%	7% •
1	Е	613	3% 89%	9% ••
2	С	10	40% 60%	
3	F	10	60% 4	10%
4	G	10	40% 60%	



Mol	Chain	Length	Q	uality of chain	
5	Н	10	40%	50%	10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	NAG	А	650	X	-	-	-



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 23390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1 1	608	Total	С	Ν	Ο	\mathbf{S}	7	10	0
	A	008	5014	3227	855	917	15	1	10	0
1	1 B 609	600	Total	С	Ν	Ο	S	10	8	0
		009	5007	3225	847	920	15	10		0
1	р	607	Total	С	Ν	Ο	S	4	10	0
		007	5013	3223	856	918	16	4	10	U
1	1 F	C0C	Total	С	Ν	Ο	S	10	7	0
	000	4977	3208	844	910	15	10			

• Molecule 1 is a protein called Beta-glucuronidase.

• Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran ose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-L-gulopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyrano se-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	С	10	Total 116	С 64	N 2	O 50	0	0	0

• Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran ose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyran ose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	F	10	Total C N O 116 64 2 50	0	0	0

• Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran ose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyra nose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyra nose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyran



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	10	Total 116	С 64	N 2	O 50	0	0	0

• Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran ose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyra nose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyra nose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyran



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
5	Н	10	Total C N O 116 64 2 50	0	0	0

• Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf	
6	Δ	1	Total	С	Ν	Ο	0	0	
0 11	T	14	8	1	5	0	0		
6	В	1	Total	С	Ν	Ο	0	0	
0	0 D	T	14	8	1	5	0		
6	р	1	Total	С	Ν	Ο	0	0	
0	6 D	L	14	8	1	5	0		
6	F	1	Total	С	Ν	0	0	0	
			14	8	1	5		U	

• Molecule 7 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: $C_6H_{14}O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 8 6 2 \end{array}$	0	0
7	D	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 8 6 2 \end{array}$	0	0
7	Е	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 8 6 2 \end{array}$	0	0

• Molecule 8 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 8 & 6 & 2 \end{array}$	0	0
8	Е	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 8 6 2 \end{array}$	0	0

• Molecule 9 is beta-D-mannopyranose (three-letter code: BMA) (formula: $C_6H_{12}O_6$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	Е	1	Total 12	$\begin{array}{c} \mathrm{C} \\ \mathrm{6} \end{array}$	O 6	0	0

• Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	691	Total O 691 691	0	0
10	В	706	Total O 706 706	0	0
10	D	646	Total O 646 646	0	0
10	Е	764	Total O 764 764	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Beta-glucuronidase





 $\label{eq:mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-L-gulopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluco$

Chain C:	40%	60%
NAG 1 NAG 2 NAG 2 MAA 3 MAN 5 MAN 5 MAN 9 GUP 10 GUP 10		

• Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[beta-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

Chain F:	60%	40%
NAG1 NAG2 BMA3 MAN5 MAN5 MAN5 MAN5 MAN5 MAN5 BMA10 BMA10		

 $\label{eq:mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glu copyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glu copyranose$

Chain G:	40%	60%
NAG1 NAG2 BMA3 BMA3 MAN6 MAN6 MAN6 MAN7 MAN9 MAN10 MAN10		

 $\label{eq:stability} \bullet \ Molecule \ 5: \ alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)] \\ alpha-D-mannopyranose-(1-6)] \\ beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranoy-2-acetamido-2-deoxy-beta-D-glucopyranoy-$



Chain H:	40%	50%	10%
NAG1 NAG2 BMA3 MAN4 MAN5 MAN5 MAN7 MAN7 MAN3 MAN9 MAN10			



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	93.58Å 123.11 Å 266.13 Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
$\mathbf{Bosolution} \left(\overset{\circ}{\mathbf{A}} \right)$	19.90 - 1.70	Depositor
Resolution (A)	19.92 - 1.66	EDS
% Data completeness	90.7 (19.90-1.70)	Depositor
(in resolution range)	84.5(19.92 - 1.66)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.17 (at 1.67 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
D D .	0.203 , 0.240	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.211 , (Not available)	DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	11.7	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	$0.35\;,59.4$	EDS
L-test for $twinning^2$	$ \langle L \rangle = 0.43, \langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	23390	wwPDB-VP
Average B, all atoms $(Å^2)$	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 32.11 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.9123e-04.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, BMA, NAG, GUP, MRD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	ond lengths	B	ond angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.48	2/5203~(0.0%)	0.63	1/7080~(0.0%)
1	В	0.49	4/5188~(0.1%)	0.65	3/7063~(0.0%)
1	D	0.44	0/5199	0.63	1/7076~(0.0%)
1	Е	0.70	5/5150~(0.1%)	0.66	5/7012~(0.1%)
All	All	0.54	11/20740~(0.1%)	0.64	10/28231~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Е	0	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	Е	266	GLU	CD-OE1	27.08	1.55	1.25
1	Е	266	GLU	CD-OE2	-23.65	0.99	1.25
1	Е	194	LYS	CD-CE	-12.30	1.20	1.51
1	А	424	HIS	CB-CG	-11.31	1.29	1.50
1	В	239	GLN	CD-OE1	-9.06	1.04	1.24
1	В	110	ARG	CG-CD	-8.85	1.29	1.51
1	Ε	295	GLU	CD-OE2	8.59	1.35	1.25
1	В	281	LYS	CD-CE	-6.79	1.34	1.51
1	Ε	295	GLU	CD-OE1	6.75	1.33	1.25
1	А	38	CYS	CB-SG	-5.45	1.73	1.81
1	B	38	CYS	CB-SG	-5.01	1.73	1.81

All (10) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	110	ARG	CB-CG-CD	10.21	138.13	111.60
1	Е	295	GLU	OE1-CD-OE2	-10.02	111.28	123.30
1	Е	194	LYS	CG-CD-CE	8.36	136.97	111.90
1	Е	266	GLU	CG-CD-OE2	7.32	132.94	118.30
1	А	424	HIS	CA-CB-CG	5.85	123.54	113.60
1	D	295	GLU	CA-CB-CG	5.59	125.70	113.40
1	Е	48	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	E	266	GLU	CG-CD-OE1	-5.05	108.19	118.30
1	В	139[A]	THR	CA-CB-CG2	5.01	119.41	112.40
1	В	139[B]	THR	CA-CB-CG2	5.01	119.41	112.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	Ε	295	GLU	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	5014	0	4876	35	0
1	В	5007	0	4864	28	0
1	D	5013	0	4860	34	0
1	Е	4977	0	4833	37	0
2	С	116	0	97	1	0
3	F	116	0	97	0	0
4	G	116	0	97	1	0
5	Н	116	0	97	3	0
6	А	14	0	13	1	0
6	В	14	0	13	0	0
6	D	14	0	13	1	0
6	Е	14	0	13	0	0
7	А	8	0	14	1	0
7	D	8	0	14	3	0
7	Е	8	0	14	0	0
8	В	8	0	14	1	0
8	E	8	0	14	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	Е	12	0	12	3	0
10	А	691	0	0	7	0
10	В	706	0	0	5	0
10	D	646	0	0	7	0
10	Е	764	0	0	5	0
All	All	23390	0	19955	137	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (137) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
	Atom-2	distance (Å)	overlap (Å)	
1:D:465:MET:HA	1:D:465:MET:HE2	1.52	0.90	
1:D:390[B]:GLU:OE1	10:D:2481:HOH:O	1.91	0.86	
1:E:183:PRO:HG3	1:E:412:LEU:HD23	1.57	0.86	
1:B:143[B]:GLU:OE1	1:B:425:HIS:NE2	2.10	0.85	
1:E:134:VAL:HG13	1:E:166:LEU:HD11	1.65	0.78	
1:D:585:LEU:HD13	10:D:2217:HOH:O	1.82	0.77	
1:E:243:LEU:HD22	10:E:2067:HOH:O	1.84	0.76	
1:A:390[B]:GLU:OE2	10:A:2523:HOH:O	2.03	0.75	
1:B:42:ASP:OD1	10:B:2467:HOH:O	2.03	0.75	
1:A:602:LEU:HD12	1:E:549:PHE:HD1	1.51	0.74	
1:A:431:GLU:HG2	1:A:473:LEU:HD11	1.70	0.74	
1:D:465:MET:HA	1:D:465:MET:CE	2.17	0.74	
1:B:424:HIS:HA	1:B:427:MET:HE2	1.70	0.73	
1:A:143[B]:GLU:OE1	1:A:425:HIS:NE2	2.22	0.71	
1:D:42:ASP:OD1	10:D:2498:HOH:O	2.10	0.70	
1:D:522:ALA:O	1:D:526:GLU:HG2	1.92	0.70	
1:D:134:VAL:HG13	1:D:166:LEU:HD11	1.75	0.69	
1:D:106:ILE:HG23	10:D:2390:HOH:O	1.92	0.69	
7:D:3:MRD:O2	7:D:3:MRD:C5	2.41	0.68	
1:D:260:VAL:CG1	1:D:303:LEU:HD11	2.25	0.66	
6:D:650:NAG:H81	10:D:2146:HOH:O	1.95	0.65	
1:A:524:GLN:HG3	7:A:1:MRD:H1C3	1.77	0.65	
1:B:374:ARG:CZ	10:B:702:HOH:O	2.46	0.63	
1:B:134:VAL:HG13	1:B:166:LEU:HD11	1.81	0.63	
9:E:661:BMA:H1	5:H:10:MAN:O2	1.99	0.62	
1:E:464:LYS:NZ	10:E:1464:HOH:O	2.32	0.62	
1:A:465:MET:SD	10:A:1818:HOH:O	2.56	0.61	
1:D:29:TYR:CD2	1:D:394[A]:GLN:NE2	2.68	0.61	



	h - 0	Interatomic	Clash	
Atom-1	Atom-2	$distance (m \AA)$	overlap (Å)	
1:D:431:GLU:HG2	1:D:473:LEU:HD11	1.83	0.60	
1:D:123:ILE:HD13	1:D:218:VAL:HG13	1.83	0.60	
1:A:308[A]:THR:HG22	10:A:2075:HOH:O	2.02	0.60	
1:A:412:LEU:HD21	10:A:856:HOH:O	2.02	0.59	
9:E:661:BMA:C1	5:H:10:MAN:O2	2.50	0.59	
1:A:602:LEU:HD12	1:E:549:PHE:CD1	2.36	0.59	
1:B:123:ILE:HD13	1:B:218:VAL:HG13	1.85	0.59	
1:B:177:THR:HG22	10:B:2270:HOH:O	2.03	0.58	
1:E:137:VAL:O	1:E:139:THR:HG23	2.03	0.58	
1:A:257:LYS:HG3	1:A:308[A]:THR:OG1	2.04	0.56	
1:A:48[B]:ARG:HD3	1:A:73:PRO:O	2.04	0.56	
1:D:48[B]:ARG:NH2	1:D:68:LEU:O	2.39	0.56	
1:A:567:GLU:HG3	1:A:621[B]:LEU:HD21	1.87	0.56	
1:E:134:VAL:HG13	1:E:166:LEU:CD1	2.34	0.56	
1:E:501:LEU:HD22	8:E:4:MPD:HM2	1.88	0.55	
1:B:123:ILE:CD1	1:B:218:VAL:HG13	2.37	0.55	
1:E:424:HIS:HA	1:E:427:MET:CE	2.37	0.55	
1:D:134:VAL:HG13	1:D:166:LEU:CD1	2.36	0.55	
1:E:260:VAL:CG1	1:E:303:LEU:HD11	2.36	0.55	
1:B:139[A]:THR:HG23	1:B:156:LEU:HD11	1.88	0.54	
1:B:156:LEU:HD22	10:B:1892:HOH:O	2.08	0.53	
1:B:134:VAL:HG13	1:B:166:LEU:CD1	2.39	0.53	
7:D:3:MRD:O2	7:D:3:MRD:H5C3	2.07	0.53	
1:D:260:VAL:HG11	1:D:303:LEU:HD11	1.91	0.53	
1:A:29:TYR:CD2	1:A:394[A]:GLN:NE2	2.77	0.52	
1:E:48:ARG:HD3	1:E:73:PRO:O	2.09	0.52	
1:E:424:HIS:HA	1:E:427:MET:HE3	1.92	0.52	
1:A:107:LEU:HD11	1:A:166:LEU:HD22	1.92	0.52	
1:E:243:LEU:CD2	10:E:2067:HOH:O	2.49	0.52	
1:D:177:THR:HG22	4:G:2:NAG:O6	2.09	0.52	
1:D:567:GLU:HG3	1:D:621:LEU:HD11	1.92	0.51	
1:A:424:HIS:HA	1:A:427:MET:CE	2.39	0.51	
1:D:350:LYS:NZ	10:D:973:HOH:O	2.42	0.51	
1:B:109:GLU:O	1:B:113:GLN:HG2	2.11	0.51	
1:D:238:GLU:HG3	1:D:243:LEU:HD11	1.93	0.51	
1:D:238:GLU:CG	1:D:243:LEU:HD11	2.42	0.50	
1:B:501:LEU:HD22	8:B:2:MPD:HM2	1.93	0.49	
1:A:431:GLU:HG2	1:A:473:LEU:CD1	2.42	0.49	
1:D:431:GLU:HG2	1:D:473:LEU:CD1	2.43	0.49	
1:E:134:VAL:CG1	1:E:166:LEU:HD11	2.38	0.48	
1:A:419:ASN:OD1	1:A:422:SER:N	2.45	0.48	



	A L D	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:106:ILE:HD13	1:D:33:SER:HA	1.96	0.48	
1:B:521:LEU:HD23	1:B:572:GLY:HA3	1.95	0.47	
1:A:310:GLN:OE1	1:A:310:GLN:N	2.48	0.47	
1:E:181:LEU:HD22	1:E:409:GLY:HA2	1.97	0.46	
1:B:58:GLY:HA2	1:B:63:TRP:CE2	2.51	0.46	
1:E:285:VAL:HG13	1:E:287:LEU:HD13	1.97	0.45	
1:E:374[B]:ARG:HD3	10:E:1026:HOH:O	2.16	0.45	
1:E:143:GLU:OE2	1:E:425:HIS:NE2	2.45	0.45	
1:D:285:VAL:HG13	1:D:287:LEU:HD13	1.98	0.45	
1:E:183:PRO:CG	1:E:412:LEU:HD23	2.37	0.45	
1:E:412:LEU:HD22	10:E:806:HOH:O	2.17	0.45	
1:E:263:LEU:CD2	1:E:269:VAL:HG22	2.46	0.45	
1:E:331[A]:VAL:HG13	1:E:535:PRO:HG2	1.99	0.44	
1:B:567:GLU:HG3	1:B:621[B]:LEU:HD11	2.00	0.44	
1:E:493:ALA:N	1:E:494:PRO:CD	2.80	0.44	
1:E:363:TRP:HB2	1:E:364:PRO:HD3	1.98	0.44	
1:B:139[A]:THR:CG2	1:B:156:LEU:HD11	2.46	0.44	
1:A:412:LEU:CD2	10:A:856:HOH:O	2.64	0.44	
1:D:181:LEU:HD22	1:D:409:GLY:HA2	2.00	0.44	
1:D:112:THR:HG22	1:D:157:VAL:HG13	2.00	0.43	
1:A:532:TYR:OH	10:A:2494:HOH:O	2.18	0.43	
1:E:189:LEU:HD13	1:E:195:TYR:CZ	2.53	0.43	
7:D:3:MRD:O2	7:D:3:MRD:H5C2	2.16	0.43	
1:A:424:HIS:HA	1:A:427:MET:HE2	2.00	0.43	
1:A:23:LEU:HD12	1:A:23:LEU:H	1.83	0.43	
1:A:521:LEU:HD23	1:A:572:GLY:HA3	2.01	0.43	
1:E:260:VAL:HG11	1:E:303:LEU:HD11	2.00	0.43	
1:A:274:THR:HG23	6:A:650:NAG:C8	2.49	0.42	
1:B:374:ARG:NH1	10:B:2089:HOH:O	2.51	0.42	
9:E:661:BMA:O1	5:H:1:NAG:H3	2.18	0.42	
1:A:493:ALA:N	1:A:494:PRO:CD	2.82	0.42	
1:E:345:PHE:HB2	1:E:583:GLY:HA3	2.01	0.42	
1:A:268:LYS:NZ	1:A:270:VAL:HG12	2.34	0.42	
1:E:384:SER:HA	1:E:385:HIS:HA	1.85	0.42	
1:B:181:LEU:HD11	1:B:426:HIS:CG	2.54	0.42	
1:E:155:ASN:ND2	1:E:156:LEU:HD13	2.34	0.42	
1:E:503[B]:SER:OG	1:E:505:TYR:CZ	2.69	0.42	
1:E:586:ILE:HD13	1:E:608:ILE:HG21	2.02	0.42	
1:E:58:GLY:HA2	1:E:63:TRP:CD2	2.54	0.42	
1:A:507:TRP:CZ2	1:A:544:GLU:HB2	2.55	0.42	
1:D:139:THR:O	1:D:139:THR:HG22	2.20	0.42	



Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:A:198:GLY:HA2	2:C:8:MAN:C6	2.50	0.42	
1:B:549:PHE:CD2	1:D:602:LEU:HD12	2.54	0.41	
1:B:58:GLY:HA2	1:B:63:TRP:CD2	2.56	0.41	
1:D:621:LEU:HD23	1:D:621:LEU:C	2.40	0.41	
1:A:181:LEU:HD11	1:A:426:HIS:CG	2.54	0.41	
1:B:139[A]:THR:HG21	1:B:156:LEU:CD1	2.50	0.41	
1:A:464:LYS:HD2	1:A:495:TYR:CE1	2.56	0.41	
1:E:556:MET:O	1:E:557:PHE:HB2	2.19	0.41	
1:D:245:ASN:ND2	10:D:1806:HOH:O	2.53	0.41	
1:D:58:GLY:HA2	1:D:63:TRP:CE2	2.55	0.41	
1:B:139[A]:THR:HG21	1:B:156:LEU:HD12	2.01	0.41	
1:B:263:LEU:CD1	1:B:269:VAL:HG22	2.51	0.41	
1:A:621[A]:LEU:HD22	10:A:717:HOH:O	2.20	0.41	
1:D:263:LEU:HB3	1:D:267:ASN:HA	2.03	0.41	
1:E:147:LEU:HD13	1:E:432:GLU:HB3	2.03	0.40	
1:A:134:VAL:HG13	1:A:166:LEU:HD11	2.03	0.40	
1:A:77:MET:SD	1:A:78:PRO:HD2	2.62	0.40	
1:E:183:PRO:HG3	1:E:412:LEU:CD2	2.41	0.40	
1:E:424:HIS:HA	1:E:427:MET:HE2	2.03	0.40	
1:B:504:TYR:HB3	1:B:507:TRP:HB3	2.03	0.40	
1:D:29:TYR:CE2	1:D:394[A]:GLN:NE2	2.85	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	614/613~(100%)	592~(96%)	22~(4%)	0	100	100
1	В	614/613~(100%)	593~(97%)	21 (3%)	0	100	100
1	D	613/613~(100%)	594 (97%)	19 (3%)	0	100	100
1	Е	609/613~(99%)	586~(96%)	23 (4%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2450/2452~(100%)	2365~(96%)	85~(4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	547/542~(101%)	538~(98%)	9(2%)	62 48
1	В	546/542~(101%)	540~(99%)	6 (1%)	73 63
1	D	548/542~(101%)	542~(99%)	6 (1%)	73 63
1	Е	542/542~(100%)	533~(98%)	9 (2%)	60 46
All	All	2183/2168~(101%)	2153~(99%)	30 (1%)	67 53

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	71	SER
1	А	138[A]	ASP
1	А	138[B]	ASP
1	А	165	ARG
1	А	217	SER
1	А	240	ASP
1	А	254	ASN
1	А	327	ARG
1	А	509	HIS
1	В	110	ARG
1	В	187	GLN
1	В	327	ARG
1	В	420	ASN
1	В	509	HIS
1	В	534	LYS
1	D	109	GLU
1	D	110	ARG



Mol	Chain	$\overline{\mathbf{Res}}$	Type
1	D	287	LEU
1	D	312	SER
1	D	327	ARG
1	D	509	HIS
1	Е	156	LEU
1	Е	159	VAL
1	Е	217	SER
1	Е	219	LEU
1	Е	267	ASN
1	Е	287	LEU
1	E	327	ARG
1	Е	509	HIS
1	Е	578	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	245	ASN
1	А	279	GLN
1	В	575	GLN
1	D	245	ASN
1	D	279	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

40 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



3.4-1	m		Der	T • 1	Bo	ond leng	ths	B	ond ang	les
	Type	Chain	Res		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	С	1	1,2	14, 14, 15	0.39	0	$17,\!19,\!21$	0.82	0
2	GUP	С	10	2	11,11,12	0.58	0	$15,\!15,\!17$	0.59	0
2	NAG	С	2	2	14, 14, 15	0.56	0	17,19,21	0.99	0
2	BMA	С	3	2	11, 11, 12	0.24	0	$15,\!15,\!17$	0.87	1(6%)
2	MAN	С	4	2	11,11,12	0.68	0	$15,\!15,\!17$	0.97	0
2	MAN	С	5	2	11,11,12	0.55	0	$15,\!15,\!17$	1.30	2(13%)
2	MAN	С	6	2	11,11,12	0.65	0	$15,\!15,\!17$	0.80	1(6%)
2	MAN	С	7	2	11,11,12	0.59	0	$15,\!15,\!17$	0.99	1(6%)
2	MAN	С	8	2	11,11,12	0.57	0	$15,\!15,\!17$	0.67	0
2	MAN	С	9	2	11,11,12	0.53	0	$15,\!15,\!17$	1.38	3 (20%)
3	NAG	F	1	1,3	14,14,15	0.45	0	17,19,21	0.86	0
3	BMA	F	10	3	11,11,12	0.78	0	$15,\!15,\!17$	2.23	4 (26%)
3	NAG	F	2	3	14,14,15	0.62	0	17,19,21	1.18	0
3	BMA	F	3	3	11, 11, 12	0.33	0	$15,\!15,\!17$	1.03	0
3	MAN	F	4	3	11,11,12	0.56	0	$15,\!15,\!17$	0.91	0
3	MAN	F	5	3	11,11,12	0.60	0	$15,\!15,\!17$	1.19	1 (6%)
3	MAN	F	6	3	11,11,12	0.67	0	$15,\!15,\!17$	0.56	0
3	MAN	F	7	3	11, 11, 12	0.43	0	$15,\!15,\!17$	0.93	1(6%)
3	MAN	F	8	3	11, 11, 12	0.59	0	$15,\!15,\!17$	0.81	0
3	MAN	F	9	3	11,11,12	0.69	0	$15,\!15,\!17$	1.08	1(6%)
4	NAG	G	1	1,4	14,14,15	0.52	0	$17,\!19,\!21$	0.97	0
4	MAN	G	10	4	11,11,12	0.77	0	$15,\!15,\!17$	1.37	3 (20%)
4	NAG	G	2	4	14,14,15	0.56	0	17,19,21	0.93	0
4	BMA	G	3	4	11,11,12	0.24	0	$15,\!15,\!17$	1.11	1(6%)
4	MAN	G	4	4	11,11,12	0.49	0	$15,\!15,\!17$	1.06	1(6%)
4	MAN	G	5	4	11,11,12	0.55	0	$15,\!15,\!17$	0.97	2 (13%)
4	MAN	G	6	4	11,11,12	0.71	0	15,15,17	0.95	0
4	MAN	G	7	4	11,11,12	0.56	0	$15,\!15,\!17$	1.13	1(6%)
4	MAN	G	8	4	11,11,12	0.61	0	$15,\!15,\!17$	0.64	0
4	MAN	G	9	4	11,11,12	0.77	0	$15,\!15,\!17$	0.82	0
5	NAG	Н	1	1,5	14,14,15	0.79	1 (7%)	17,19,21	0.80	0
5	MAN	Н	10	5	11,11,12	0.60	0	15,15,17	0.97	0
5	NAG	Н	2	5	14,14,15	0.46	0	17,19,21	1.11	0
5	BMA	Н	3	5	11,11,12	0.44	0	$15,\!15,\!17$	1.01	0
5	MAN	Н	4	5	11,11,12	0.59	0	$15,\!15,\!17$	1.23	2 (13%)
5	MAN	Н	5	5	11,11,12	0.60	0	$15,\!15,\!17$	1.21	2 (13%)

expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



	True	Chain	Dog	Tink	Bo	Bond lengths			Bond angles		
	n Type Cham Res	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2		
5	MAN	Н	6	5	11,11,12	0.55	0	$15,\!15,\!17$	0.78	0	
5	MAN	Н	7	5	11,11,12	0.65	0	15,15,17	1.28	1 (6%)	
5	MAN	Н	8	5	11,11,12	0.53	0	15,15,17	0.98	1 (6%)	
5	MAN	Н	9	5	11,11,12	0.63	0	15,15,17	0.74	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	GUP	С	10	2	-	0/2/19/22	0/1/1/1
2	NAG	С	2	2	-	0/6/23/26	0/1/1/1
2	BMA	С	3	2	-	0/2/19/22	0/1/1/1
2	MAN	С	4	2	-	0/2/19/22	0/1/1/1
2	MAN	С	5	2	-	0/2/19/22	0/1/1/1
2	MAN	С	6	2	-	2/2/19/22	0/1/1/1
2	MAN	С	7	2	_	0/2/19/22	0/1/1/1
2	MAN	С	8	2	-	0/2/19/22	0/1/1/1
2	MAN	С	9	2	-	0/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	BMA	F	10	3	-	0/2/19/22	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
3	MAN	F	5	3	-	2/2/19/22	0/1/1/1
3	MAN	F	6	3	-	2/2/19/22	0/1/1/1
3	MAN	F	7	3	-	0/2/19/22	0/1/1/1
3	MAN	F	8	3	_	0/2/19/22	0/1/1/1
3	MAN	F	9	3	-	0/2/19/22	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	MAN	G	10	4	-	2/2/19/22	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	MAN	G	4	4	-	0/2/19/22	0/1/1/1
4	MAN	G	5	4	-	1/2/19/22	0/1/1/1
4	MAN	G	6	4	-	2/2/19/22	0/1/1/1
4	MAN	G	7	4	-	0/2/19/22	0/1/1/1
4	MAN	G	8	4	-	1/2/19/22	0/1/1/1
4	MAN	G	9	4	-	0/2/19/22	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	Н	1	1,5	-	0/6/23/26	0/1/1/1
5	MAN	Н	10	5	-	1/2/19/22	0/1/1/1
5	NAG	Н	2	5	-	0/6/23/26	0/1/1/1
5	BMA	Н	3	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	4	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	5	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	6	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	7	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	8	5	-	0/2/19/22	0/1/1/1
5	MAN	Н	9	5	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
5	Н	1	NAG	O5-C1	-2.36	1.39	1.43

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	F	10	BMA	C1-O5-C5	4.65	118.50	112.19
3	F	10	BMA	O5-C1-C2	4.56	117.81	110.77
3	F	10	BMA	C1-C2-C3	4.05	114.64	109.67
2	С	9	MAN	O2-C2-C1	-3.68	101.61	109.15
5	Н	7	MAN	C1-O5-C5	3.67	117.16	112.19
2	С	5	MAN	C1-O5-C5	3.33	116.70	112.19
3	F	5	MAN	C1-O5-C5	2.90	116.12	112.19
4	G	10	MAN	C3-C4-C5	2.81	115.25	110.24
3	F	10	BMA	C3-C4-C5	2.72	115.10	110.24
3	F	9	MAN	C1-C2-C3	2.72	113.01	109.67
5	Н	4	MAN	O2-C2-C3	-2.69	104.75	110.14
4	G	10	MAN	C2-C3-C4	2.51	115.24	110.89
5	Н	5	MAN	C1-O5-C5	2.39	115.43	112.19
5	Н	5	MAN	O2-C2-C1	2.37	114.01	109.15
5	Н	4	MAN	C2-C3-C4	-2.32	106.88	110.89
2	С	7	MAN	O2-C2-C1	2.31	113.88	109.15
4	G	3	BMA	O3-C3-C2	-2.30	105.60	109.99
4	G	7	MAN	O2-C2-C3	-2.23	105.67	110.14
4	G	10	MAN	O5-C1-C2	-2.22	107.34	110.77
2	С	5	MAN	O2-C2-C3	-2.21	105.72	110.14
2	С	9	MAN	O2-C2-C3	2.18	114.52	110.14
4	G	5	MAN	O2-C2-C3	-2.14	105.85	110.14
4	G	5	MAN	C1-O5-C5	2.08	115.00	112.19



Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	С	6	MAN	O5-C1-C2	-2.06	107.59	110.77
2	С	9	MAN	C1-O5-C5	2.02	114.93	112.19
3	F	7	MAN	C1-O5-C5	2.01	114.92	112.19
2	С	3	BMA	O6-C6-C5	-2.01	104.39	111.29
5	Н	8	MAN	O5-C1-C2	-2.01	107.67	110.77
4	G	4	MAN	C1-O5-C5	2.00	114.90	112.19

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
4	G	10	MAN	O5-C5-C6-O6
4	G	10	MAN	C4-C5-C6-O6
2	С	6	MAN	C4-C5-C6-O6
3	F	6	MAN	C4-C5-C6-O6
4	G	6	MAN	C4-C5-C6-O6
3	F	5	MAN	C4-C5-C6-O6
3	F	6	MAN	O5-C5-C6-O6
2	С	6	MAN	O5-C5-C6-O6
4	G	6	MAN	O5-C5-C6-O6
4	G	5	MAN	C4-C5-C6-O6
3	F	5	MAN	O5-C5-C6-O6
4	G	8	MAN	C4-C5-C6-O6
5	Н	10	MAN	C4-C5-C6-O6

All (13) torsion outliers are listed below:

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	\mathbf{Res}	Type	Clashes	Symm-Clashes
2	С	8	MAN	1	0
5	Н	10	MAN	2	0
4	G	2	NAG	1	0
5	Н	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.













5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Tune	Chain	Dog	Tink	Bo	ond leng	$_{\rm ths}$	В	ond ang	les
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	MRD	E	5	-	7,7,7	0.25	0	$9,\!10,\!10$	0.18	0
8	MPD	Е	4	-	7,7,7	0.24	0	$9,\!10,\!10$	0.54	0
6	NAG	D	650	1	14,14,15	0.52	0	$17,\!19,\!21$	0.75	0
7	MRD	D	3	-	7,7,7	0.31	0	$9,\!10,\!10$	0.72	0
6	NAG	А	650	1	14,14,15	0.50	0	$17,\!19,\!21$	1.37	3 (17%)
8	MPD	В	2	-	7,7,7	0.35	0	$9,\!10,\!10$	0.70	0
7	MRD	А	1	-	7,7,7	0.28	0	$9,\!10,\!10$	0.54	0
9	BMA	E	661	-	12,12,12	0.58	0	$17,\!17,\!17$	1.86	4 (23%)
6	NAG	В	650	1	14,14,15	0.54	0	$17,\!19,\!21$	0.83	0
6	NAG	Е	650	1	14,14,15	0.50	0	$17,\!19,\!21$	0.91	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MRD	Е	5	-	-	0/5/5/5	-
8	MPD	Е	4	-	-	3/5/5/5	-
6	NAG	D	650	1	-	2/6/23/26	0/1/1/1
7	MRD	D	3	-	-	5/5/5/5	-
6	NAG	А	650	1	1/1/5/7	4/6/23/26	0/1/1/1
8	MPD	В	2	-	-	1/5/5/5	-
7	MRD	А	1	-	-	2/5/5/5	-
9	BMA	Е	661	-	-	0/2/22/22	0/1/1/1
6	NAG	В	650	1	-	2/6/23/26	0/1/1/1
6	NAG	Е	650	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
9	Ε	661	BMA	O5-C1-C2	4.20	117.78	110.28
9	Е	661	BMA	C1-O5-C5	4.02	121.26	113.66
9	Е	661	BMA	C3-C4-C5	-2.77	105.30	110.24
6	А	650	NAG	C1-O5-C5	2.70	115.85	112.19
6	А	650	NAG	O5-C5-C6	2.57	111.23	107.20
6	А	650	NAG	C8-C7-N2	2.46	120.27	116.10
6	Е	650	NAG	C1-O5-C5	2.08	115.02	112.19
9	Е	661	BMA	C1-C2-C3	2.02	114.51	110.31



All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	А	650	NAG	C1

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	D	3	MRD	O2-C2-C3-C4
7	D	3	MRD	C2-C3-C4-C5
6	А	650	NAG	C8-C7-N2-C2
6	А	650	NAG	O7-C7-N2-C2
6	D	650	NAG	C8-C7-N2-C2
6	D	650	NAG	O7-C7-N2-C2
6	В	650	NAG	C8-C7-N2-C2
6	В	650	NAG	O7-C7-N2-C2
6	А	650	NAG	O5-C5-C6-O6
6	А	650	NAG	C4-C5-C6-O6
8	Е	4	MPD	O2-C2-C3-C4
7	D	3	MRD	C1-C2-C3-C4
7	D	3	MRD	CM-C2-C3-C4
8	Е	4	MPD	C1-C2-C3-C4
8	Е	4	MPD	CM-C2-C3-C4
8	В	2	MPD	CM-C2-C3-C4
7	А	1	MRD	C2-C3-C4-C5
7	D	3	MRD	C2-C3-C4-O4
7	А	1	MRD	C2-C3-C4-O4

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	Е	4	MPD	1	0
6	D	650	NAG	1	0
7	D	3	MRD	3	0
6	А	650	NAG	1	0
8	В	2	MPD	1	0
7	А	1	MRD	1	0
9	Е	661	BMA	3	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< RSRZ >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	608/613~(99%)	0.31	24 (3%) 39 44	6, 13, 27, 36	2(0%)
1	В	609/613~(99%)	0.21	21 (3%) 45 50	5, 12, 24, 34	3(0%)
1	D	607/613~(99%)	0.24	24 (3%) 38 42	6, 13, 27, 33	1 (0%)
1	Е	606/613~(98%)	0.17	19 (3%) 49 53	6, 12, 23, 30	6 (0%)
All	All	2430/2452 (99%)	0.23	88 (3%) 42 47	5, 12, 26, 36	12 (0%)

All (88) RSRZ outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	RSRZ
1	А	159	VAL	9.2
1	Е	23	LEU	8.0
1	D	633	THR	7.7
1	D	23	LEU	7.5
1	А	23	LEU	7.0
1	В	160	GLY	6.9
1	Е	159	VAL	6.9
1	D	159	VAL	6.6
1	А	90	TRP	6.1
1	А	24	GLN	5.7
1	В	633	THR	5.5
1	В	23	LEU	5.5
1	В	164	SER	5.3
1	В	90	TRP	5.0
1	В	159	VAL	5.0
1	A	22	GLY	4.9
1	A	164	SER	4.8
1	A	113	GLN	4.7
1	D	164	SER	4.4
1	Е	90	TRP	3.9
1	В	315	PRO	3.5



Mol	Chain	Res	Type	RSRZ
1	D	165	ARG	3.4
1	В	24	GLN	3.4
1	D	237	VAL	3.4
1	D	239	GLN	3.4
1	В	255	LEU	3.3
1	Е	578	ARG	3.3
1	А	239	GLN	3.2
1	А	165	ARG	3.1
1	В	240[A]	ASP	3.1
1	А	420	ASN	3.1
1	Е	631	ASN	3.1
1	В	424	HIS	3.1
1	Е	164	SER	3.1
1	Е	296	ARG	3.0
1	А	255	LEU	3.0
1	D	90	TRP	3.0
1	В	310	GLN	3.0
1	А	421	VAL	2.9
1	Е	255	LEU	2.9
1	В	69	TRP	2.9
1	Е	113	GLN	2.8
1	В	71	SER	2.8
1	В	157	VAL	2.8
1	В	296	ARG	2.7
1	D	24	GLN	2.7
1	D	578	ARG	2.7
1	А	160	GLY	2.6
1	D	268	LYS	2.6
1	Е	260	VAL	2.6
1	D	255	LEU	2.5
1	A	69	TRP	2.5
1	A	115	LEU	2.5
1	A	73	PRO	2.4
1	Е	424	HIS	2.4
1	В	139[A]	THR	2.4
1	D	241	SER	2.4
1	A	313	LEU	2.4
1	D	113	GLN	2.4
1	В	22	GLY	2.4
1	E	533	GLN	2.4
1	E	267	ASN	2.4
1	D	110	ARG	2.3



Mol	Chain	Res	Type	RSRZ
1	А	252	GLY	2.3
1	А	91	ARG	2.3
1	D	312	SER	2.3
1	D	277	GLN	2.3
1	Е	266	GLU	2.3
1	В	254	ASN	2.3
1	В	309	ALA	2.3
1	А	156	LEU	2.3
1	А	254	ASN	2.2
1	А	267	ASN	2.2
1	D	248	ILE	2.2
1	D	69	TRP	2.2
1	Е	239	GLN	2.2
1	D	296	ARG	2.1
1	D	157	VAL	2.1
1	А	312	SER	2.1
1	Е	24	GLN	2.1
1	D	261	ARG	2.1
1	А	238	GLU	2.1
1	В	113	GLN	2.1
1	Е	315	PRO	2.1
1	D	254	ASN	2.1
1	Е	268	LYS	2.0
1	D	263	LEU	2.0
1	Е	155	ASN	2.0

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} extsf{-}\mathbf{B} extsf{-}\mathbf{factors}(\mathbf{A}^2)$	Q < 0.9
4	MAN	G	10	11/12	0.47	0.34	$40,\!42,\!43,\!44$	0
2	GUP	С	10	11/12	0.62	0.31	$34,\!36,\!38,\!39$	0
3	BMA	F	10	11/12	0.63	0.17	$34,\!36,\!37,\!37$	0



Mol		Chain	\mathbf{Res}	Atoms	BSCC	RSR	B -factors($Å^2$)	Q<0.9
3	MAN	F	6	11/12	0.67	0.23	37 40 41 41	0
2	MAN	C	8	$\frac{11}{12}$	0.69	0.23	26.28.29.31	0
4	MAN	G	8	$\frac{11}{12}$	0.72	0.21	31.32.34.34	0
4	MAN	G	9	$\frac{11}{12}$	0.76	0.14	29.32.34.36	0
3	MAN	F	5	11/12	0.79	0.14	28,31,33,34	0
4	MAN	G	7	11/12	0.80	0.16	27,28,29,29	0
3	MAN	F	9	11/12	0.80	0.13	23,27,30,30	0
4	MAN	G	6	11/12	0.80	0.19	32,34,36,36	0
3	MAN	F	8	11/12	0.82	0.15	26,26,28,29	0
2	MAN	С	9	11/12	0.83	0.18	$22,\!23,\!26,\!29$	0
2	MAN	С	6	11/12	0.83	0.16	$28,\!30,\!32,\!33$	0
2	MAN	С	7	11/12	0.84	0.11	$22,\!23,\!25,\!26$	0
3	NAG	F	2	14/15	0.86	0.12	$11,\!13,\!18,\!19$	0
4	MAN	G	5	11/12	0.87	0.10	$24,\!26,\!29,\!29$	0
3	MAN	F	7	11/12	0.87	0.13	$18,\!20,\!21,\!23$	0
2	MAN	С	5	11/12	0.88	0.17	$20,\!21,\!23,\!26$	0
3	MAN	F	4	11/12	0.89	0.10	$15,\!17,\!20,\!23$	0
5	MAN	Н	9	11/12	0.89	0.12	$15,\!18,\!20,\!20$	0
4	MAN	G	4	11/12	0.89	0.09	$19,\!21,\!23,\!24$	0
2	MAN	С	4	11/12	0.90	0.09	$18,\!19,\!20,\!21$	0
3	NAG	F	1	14/15	0.90	0.11	11,13,19,19	0
4	NAG	G	2	14/15	0.91	0.12	12,14,17,21	0
5	MAN	H	6	11/12	0.91	0.12	15,16,18,19	0
3	BMA	F	3	11/12	0.92	0.09	12,13,15,19	0
4	BMA	G	3	11/12	0.92	0.08	13,16,18,23	0
2	NAG	С	1	14/15	0.93	0.09	10,12,15,15	0
2	NAG	С	2	14/15	0.93	0.10	12,14,18,19	0
5	MAN	Н	8	11/12	0.94	0.08	15,17,21,21	0
5	MAN	Н	4	11/12	0.94	0.08	9,11,14,15	0
2	BMA	С	3	11/12	0.94	0.10	12,14,16,19	0
4	NAG	G	1	14/15	0.95	0.08	9,12,14,15	0
5	MAN	H	10	11/12	0.95	0.07	11,14,16,17	0
5	MAN	H	5	11/12	0.95	0.08	12,13,14,15	0
5	NAG	H	2	14/15	0.95	0.08	7,9,13,13	0
5	BMA	H	3	11/12	0.96	0.07	7,9,10,11	0
5	MAN	H	7	11/12	0.96	0.07	8,11,13,14	0
5	NAG	H	$\mid 1$	14/15	0.97	0.07	7,9,10,10	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.













6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	${f B} ext{-factors}({f A}^2)$	Q < 0.9
6	NAG	А	650	14/15	0.62	0.23	$37,\!40,\!41,\!41$	0
9	BMA	Е	661	12/12	0.62	0.22	$33,\!36,\!36,\!38$	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
7	MRD	Е	5	8/8	0.72	0.20	$36,\!36,\!37,\!38$	0
6	NAG	D	650	14/15	0.75	0.24	$31,\!33,\!35,\!36$	0
6	NAG	Е	650	14/15	0.78	0.19	24,30,34,34	0
6	NAG	В	650	14/15	0.81	0.16	$20,\!22,\!24,\!25$	0
7	MRD	А	1	8/8	0.82	0.17	$23,\!26,\!27,\!28$	0
8	MPD	Е	4	8/8	0.86	0.16	$24,\!26,\!28,\!29$	0
8	MPD	В	2	8/8	0.86	0.13	$20,\!23,\!25,\!27$	0
7	MRD	D	3	8/8	0.87	0.16	$25,\!27,\!29,\!30$	0

Continued from previous page...

6.5 Other polymers (i)

There are no such residues in this entry.

