



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 01:49 PM BST

PDB ID : 4HNP
Title : Crystal structure of yeast 20S proteasome in complex with vinylketone carmaphycin analogue VNK1
Authors : Trivella, D.B.B.; Stein, M.; Groll, M.
Deposited on : 2012-10-20
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 89%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">2% 89% 10% •</p>
1	O	250	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 90%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">2% 90% 10% •</p>
2	B	244	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 84%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">2% 84% 13% •</p>
2	P	244	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">2% 88% 11% •</p>
3	C	241	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 85%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">3% 85% 13% •</p>
3	Q	241	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 89%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 1%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">6% 89% 11% •</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	242	 2% 90% 9%
4	R	242	 3% 91% 9%
5	E	233	 2% 89% 10%
5	S	233	 2% 86% 13%
6	F	244	 % 84% 14%
6	T	244	 2% 86% 13%
7	G	243	 % 88% 11%
7	U	243	 % 89% 11%
8	H	222	 % 91% 9%
8	V	222	 % 90% 10%
9	I	204	 % 91% 9%
9	W	204	 % 88% 12%
10	J	198	 2% 89% 10%
10	X	198	 2% 88% 12%
11	K	212	 % 87% 11%
11	Y	212	 % 92% 7%
12	L	222	 % 89% 9%
12	Z	222	 % 89% 9%
13	M	233	 % 80% 20%
13	a	233	 % 95% 5%
14	N	196	 % 92% 8%
14	b	196	 % 97%

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 50504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome component Y7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	1915	1219	315	377	4	0	0	0
1	O	250	1915	1219	315	377	4	0	0	0

- Molecule 2 is a protein called Proteasome component Y13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	1904	1201	321	379	3	0	0	0
2	P	244	1904	1201	321	379	3	0	0	0

- Molecule 3 is a protein called Proteasome component PRE6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	241	1890	1181	331	374	4	0	0	0
3	Q	241	1890	1181	331	374	4	0	0	0

- Molecule 4 is a protein called Proteasome component PUP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	242	1861	1162	314	378	7	0	0	0
4	R	242	1861	1162	314	378	7	0	0	0

- Molecule 5 is a protein called Proteasome component PRE5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			
5	S	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			

- Molecule 6 is a protein called Proteasome component C1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	244	Total	C	N	O	S	0	0	0
			1896	1205	330	357	4			
6	T	244	Total	C	N	O	S	0	0	0
			1896	1205	330	357	4			

- Molecule 7 is a protein called Proteasome component C7-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			
7	U	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			

- Molecule 8 is a protein called Proteasome component PUP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			
8	V	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			

- Molecule 9 is a protein called Proteasome component PUP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome component C11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	X	198	1585	1005	269	305	6	0	0	0

- Molecule 11 is a protein called Proteasome component PRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	K	212	1644	1045	280	312	7	0	0	0
11	Y	212	1644	1045	280	312	7	0	0	0

- Molecule 12 is a protein called Proteasome component C5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	L	222	1757	1115	303	335	4	0	0	0
12	Z	222	1757	1115	303	335	4	0	0	0


- Molecule 13 is a protein called Proteasome component PRE4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
13	M	233	1824	1154	312	351	7	0	0	0
13	a	233	1824	1154	312	351	7	0	0	0

- Molecule 14 is a protein called Proteasome component PRE3.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
14	N	196	1512	955	250	300	7	0	0	0
14	b	196	1512	955	250	300	7	0	0	0

- Molecule 15 is N-hexanoyl-L-valyl-N 1 -[(3S,4S)-3-hydroxy-2,6-dimethylhept-1-en-4-yl]-N 5 ,N 5 -dimethyl-L-glutamamide (three-letter code: ONK) (formula: C₂₇H₅₀N₄O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
15	H	1	Total	C	N	O	0	0
			36	27	4	5		
15	V	1	Total	C	N	O	0	0
			36	27	4	5		

- Molecule 16 is N-hexanoyl-L-valyl-N 1 -[(3S,4S)-3-hydroxy-2,6-dimethylheptan-4-yl]-N 5 ,N 5 -dimethyl-L-glutamamide (three-letter code: VNK) (formula: C₂₇H₅₂N₄O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
16	K	1	Total	C	N	O	0	0
			36	27	4	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
16	N	1	Total	C	N	O	0	0
			36	27	4	5		
16	Y	1	Total	C	N	O	0	0
			36	27	4	5		
16	b	1	Total	C	N	O	0	0
			36	27	4	5		

- Molecule 17 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	29	Total	O	0	0
			29	29		
17	B	21	Total	O	0	0
			21	21		
17	C	30	Total	O	0	0
			30	30		
17	D	29	Total	O	0	0
			29	29		
17	E	38	Total	O	0	0
			38	38		
17	F	22	Total	O	0	0
			22	22		
17	G	30	Total	O	0	0
			30	30		
17	H	27	Total	O	0	0
			27	27		
17	I	40	Total	O	0	0
			40	40		
17	J	22	Total	O	0	0
			22	22		
17	K	23	Total	O	0	0
			23	23		
17	L	32	Total	O	0	0
			32	32		
17	M	27	Total	O	0	0
			27	27		
17	N	31	Total	O	0	0
			31	31		
17	O	22	Total	O	0	0
			22	22		
17	P	23	Total	O	0	0
			23	23		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	Q	29	Total 29	O 29	0	0
17	R	28	Total 28	O 28	0	0
17	S	26	Total 26	O 26	0	0
17	T	30	Total 30	O 30	0	0
17	U	21	Total 21	O 21	0	0
17	V	37	Total 37	O 37	0	0
17	W	19	Total 19	O 19	0	0
17	X	18	Total 18	O 18	0	0
17	Y	27	Total 27	O 27	0	0
17	Z	16	Total 16	O 16	0	0
17	a	29	Total 29	O 29	0	0
17	b	24	Total 24	O 24	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Proteasome component Y7




- Molecule 1: Proteasome component Y7





- Molecule 2: Proteasome component Y13





- Molecule 2: Proteasome component Y13





- Molecule 3: Proteasome component PRE6





- Molecule 3: Proteasome component PRE6




- Molecule 4: Proteasome component PUP2




- Molecule 4: Proteasome component PUP2



- Molecule 5: Proteasome component PRE5





- Molecule 5: Proteasome component PRE5








- Molecule 6: Proteasome component C1




- Molecule 6: Proteasome component C1





- Molecule 7: Proteasome component C7-alpha




- Molecule 7: Proteasome component C7-alpha



- Molecule 8: Proteasome component PUP1



- Molecule 8: Proteasome component PUP1








- Molecule 9: Proteasome component PUP3




- Molecule 9: Proteasome component PUP3




- Molecule 10: Proteasome component C11



- Molecule 10: Proteasome component C11




- Molecule 11: Proteasome component PRE2



- Molecule 11: Proteasome component PRE2




- Molecule 12: Proteasome component C5

Chain L:  89% 9%





- Molecule 12: Proteasome component C5

Chain Z:  89% 9%



- Molecule 13: Proteasome component PRE4

Chain M:  80% 20%




- Molecule 13: Proteasome component PRE4

Chain a:  95% 5%



- Molecule 14: Proteasome component PRE3

Chain N:  92% 8%



- Molecule 14: Proteasome component PRE3

Chain b:  97%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	135.85Å 301.36Å 144.46Å 90.00° 112.82° 90.00°	Depositor
Resolution (Å)	48.42 – 2.80 48.42 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.3 (48.42-2.80) 98.3 (48.42-2.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.203 , 0.239 0.203 , 0.237	Depositor DCC
R_{free} test set	12815 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	59.7	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	50504	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ONK, VNK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1952	0.50	0/2642
1	O	0.36	0/1952	0.48	0/2642
2	B	0.35	0/1934	0.51	0/2618
2	P	0.33	0/1934	0.50	0/2618
3	C	0.33	0/1919	0.50	0/2598
3	Q	0.33	0/1919	0.50	0/2598
4	D	0.32	0/1886	0.52	0/2541
4	R	0.32	0/1886	0.50	0/2541
5	E	0.33	0/1823	0.48	0/2463
5	S	0.33	0/1823	0.48	0/2463
6	F	0.34	0/1936	0.46	0/2614
6	T	0.33	0/1936	0.48	0/2614
7	G	0.34	0/1959	0.49	0/2652
7	U	0.33	0/1959	0.48	0/2652
8	H	0.30	0/1715	0.49	0/2326
8	V	0.32	0/1715	0.48	0/2326
9	I	0.33	0/1611	0.50	0/2174
9	W	0.33	0/1611	0.50	0/2174
10	J	0.33	0/1613	0.50	0/2173
10	X	0.32	0/1613	0.49	0/2173
11	K	0.33	0/1681	0.52	1/2274 (0.0%)
11	Y	0.32	0/1681	0.51	1/2274 (0.0%)
12	L	0.34	0/1795	0.50	0/2420
12	Z	0.40	0/1795	0.52	0/2420
13	M	0.34	0/1855	0.53	0/2514
13	a	0.33	0/1855	0.51	0/2514
14	N	0.31	0/1541	0.47	0/2087
14	b	0.31	0/1541	0.47	0/2087
All	All	0.33	0/50440	0.50	2/68192 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if

the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
11	K	0	1
11	Y	0	1
All	All	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	Y	4	LEU	CA-CB-CG	5.96	129.00	115.30
11	K	4	LEU	CA-CB-CG	5.71	128.44	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
11	K	1	THR	Peptide
11	Y	1	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1929	21	0
1	O	1915	0	1929	21	0
2	B	1904	0	1904	55	0
2	P	1904	0	1904	16	0
3	C	1890	0	1903	24	0
3	Q	1890	0	1903	15	0
4	D	1861	0	1839	12	0
4	R	1861	0	1839	12	0
5	E	1795	0	1800	15	0
5	S	1795	0	1800	15	0
6	F	1896	0	1889	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	T	1896	0	1889	15	0
7	G	1921	0	1913	15	0
7	U	1921	0	1913	15	0
8	H	1684	0	1687	10	0
8	V	1684	0	1687	12	0
9	I	1581	0	1574	9	0
9	W	1581	0	1574	15	0
10	J	1585	0	1590	11	0
10	X	1585	0	1590	11	0
11	K	1644	0	1592	15	0
11	Y	1644	0	1592	10	0
12	L	1757	0	1711	14	0
12	Z	1757	0	1711	11	0
13	M	1824	0	1832	23	0
13	a	1824	0	1832	0	0
14	N	1512	0	1478	7	0
14	b	1512	0	1478	0	0
15	H	36	0	49	2	0
15	V	36	0	49	0	0
16	K	36	0	50	1	0
16	N	36	0	50	2	0
16	Y	36	0	50	0	0
16	b	36	0	50	0	0
17	A	29	0	0	0	0
17	B	21	0	0	0	0
17	C	30	0	0	1	0
17	D	29	0	0	0	0
17	E	38	0	0	0	0
17	F	22	0	0	0	0
17	G	30	0	0	0	0
17	H	27	0	0	0	0
17	I	40	0	0	0	0
17	J	22	0	0	0	0
17	K	23	0	0	0	0
17	L	32	0	0	0	0
17	M	27	0	0	1	0
17	N	31	0	0	0	0
17	O	22	0	0	0	0
17	P	23	0	0	0	0
17	Q	29	0	0	0	0
17	R	28	0	0	0	0
17	S	26	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
17	T	30	0	0	0	0
17	U	21	0	0	0	0
17	V	37	0	0	0	0
17	W	19	0	0	0	0
17	X	18	0	0	0	0
17	Y	27	0	0	0	0
17	Z	16	0	0	0	0
17	a	29	0	0	0	0
17	b	24	0	0	0	0
All	All	50504	0	49580	363	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 363 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:209:ARG:HH21	2:B:209:ARG:CG	1.61	1.12
2:B:52:THR:CG2	2:B:209:ARG:HD3	1.82	1.08
7:G:92:ALA:HA	7:G:103:MET:HE2	1.48	0.96
2:B:209:ARG:HG3	2:B:209:ARG:NH2	1.51	0.94
2:B:52:THR:HG22	2:B:209:ARG:HD3	1.55	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	248/250 (99%)	245 (99%)	2 (1%)	1 (0%)	34 66
1	O	248/250 (99%)	243 (98%)	5 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	242/244 (99%)	228 (94%)	12 (5%)	2 (1%)	19	49
2	P	242/244 (99%)	230 (95%)	10 (4%)	2 (1%)	19	49
3	C	239/241 (99%)	233 (98%)	5 (2%)	1 (0%)	34	66
3	Q	239/241 (99%)	231 (97%)	8 (3%)	0	100	100
4	D	240/242 (99%)	234 (98%)	5 (2%)	1 (0%)	34	66
4	R	240/242 (99%)	231 (96%)	8 (3%)	1 (0%)	34	66
5	E	231/233 (99%)	224 (97%)	7 (3%)	0	100	100
5	S	231/233 (99%)	225 (97%)	6 (3%)	0	100	100
6	F	242/244 (99%)	232 (96%)	10 (4%)	0	100	100
6	T	242/244 (99%)	229 (95%)	13 (5%)	0	100	100
7	G	241/243 (99%)	233 (97%)	7 (3%)	1 (0%)	34	66
7	U	241/243 (99%)	237 (98%)	3 (1%)	1 (0%)	34	66
8	H	220/222 (99%)	214 (97%)	6 (3%)	0	100	100
8	V	220/222 (99%)	213 (97%)	7 (3%)	0	100	100
9	I	202/204 (99%)	198 (98%)	4 (2%)	0	100	100
9	W	202/204 (99%)	197 (98%)	5 (2%)	0	100	100
10	J	196/198 (99%)	188 (96%)	7 (4%)	1 (0%)	29	61
10	X	196/198 (99%)	188 (96%)	7 (4%)	1 (0%)	29	61
11	K	210/212 (99%)	206 (98%)	4 (2%)	0	100	100
11	Y	210/212 (99%)	204 (97%)	6 (3%)	0	100	100
12	L	220/222 (99%)	216 (98%)	4 (2%)	0	100	100
12	Z	220/222 (99%)	214 (97%)	6 (3%)	0	100	100
13	M	231/233 (99%)	221 (96%)	9 (4%)	1 (0%)	34	66
13	a	231/233 (99%)	221 (96%)	9 (4%)	1 (0%)	34	66
14	N	194/196 (99%)	189 (97%)	5 (3%)	0	100	100
14	b	194/196 (99%)	187 (96%)	6 (3%)	1 (0%)	29	61
All	All	6312/6368 (99%)	6111 (97%)	186 (3%)	15 (0%)	47	78

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	118	GLY
1	A	166	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	203	SER
2	P	203	SER
13	a	229	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	205 (98%)	4 (2%)	57	85
1	O	209/209 (100%)	201 (96%)	8 (4%)	33	67
2	B	203/203 (100%)	190 (94%)	13 (6%)	17	45
2	P	203/203 (100%)	194 (96%)	9 (4%)	28	61
3	C	213/213 (100%)	203 (95%)	10 (5%)	26	59
3	Q	213/213 (100%)	207 (97%)	6 (3%)	43	77
4	D	198/198 (100%)	191 (96%)	7 (4%)	36	70
4	R	198/198 (100%)	193 (98%)	5 (2%)	47	80
5	E	192/192 (100%)	181 (94%)	11 (6%)	20	50
5	S	192/192 (100%)	185 (96%)	7 (4%)	35	69
6	F	201/201 (100%)	191 (95%)	10 (5%)	24	56
6	T	201/201 (100%)	189 (94%)	12 (6%)	19	48
7	G	207/207 (100%)	194 (94%)	13 (6%)	18	46
7	U	207/207 (100%)	198 (96%)	9 (4%)	29	62
8	H	181/181 (100%)	176 (97%)	5 (3%)	43	77
8	V	181/181 (100%)	178 (98%)	3 (2%)	60	87
9	I	172/172 (100%)	169 (98%)	3 (2%)	60	87
9	W	172/172 (100%)	168 (98%)	4 (2%)	50	82
10	J	175/175 (100%)	169 (97%)	6 (3%)	37	71
10	X	175/175 (100%)	170 (97%)	5 (3%)	42	76
11	K	169/169 (100%)	163 (96%)	6 (4%)	35	69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	Y	169/169 (100%)	163 (96%)	6 (4%)	35	69
12	L	185/185 (100%)	179 (97%)	6 (3%)	39	73
12	Z	185/185 (100%)	173 (94%)	12 (6%)	17	44
13	M	199/199 (100%)	191 (96%)	8 (4%)	31	65
13	a	199/199 (100%)	189 (95%)	10 (5%)	24	56
14	N	162/162 (100%)	157 (97%)	5 (3%)	40	74
14	b	162/162 (100%)	157 (97%)	5 (3%)	40	74
All	All	5332/5332 (100%)	5124 (96%)	208 (4%)	32	66

5 of 208 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	M	12	ILE
2	P	62	THR
12	Z	173	LYS
13	M	48	ASN
14	N	126	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 153 such sidechains are listed below:

Mol	Chain	Res	Type
12	L	80	ASN
2	P	95	GLN
12	Z	79	HIS
12	L	165	ASN
13	M	108	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
15	ONK	H	301	8	34,35,35	0.34	0	40,46,46	0.64	0
16	VNK	Y	301	11	35,35,35	0.50	1 (2%)	42,46,46	1.28	3 (7%)
16	VNK	b	301	14	35,35,35	0.47	0	42,46,46	0.90	1 (2%)
16	VNK	N	301	14	35,35,35	0.59	1 (2%)	42,46,46	0.95	2 (4%)
16	VNK	K	301	11	35,35,35	0.46	0	42,46,46	1.19	1 (2%)
15	ONK	V	301	8	34,35,35	0.33	0	40,46,46	0.86	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	ONK	H	301	8	-	11/50/50/50	-
16	VNK	Y	301	11	-	20/50/50/50	-
16	VNK	b	301	14	-	13/50/50/50	-
16	VNK	N	301	14	-	16/50/50/50	-
16	VNK	K	301	11	-	18/50/50/50	-
15	ONK	V	301	8	-	11/50/50/50	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
16	N	301	VNK	C17-C16	2.39	1.56	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
16	Y	301	VNK	C17-C16	2.03	1.56	1.53

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
16	Y	301	VNK	C12-C11-N3	-6.34	101.98	110.18
16	K	301	VNK	C12-C11-N3	-5.63	102.91	110.18
16	b	301	VNK	C12-C11-N3	-4.18	104.78	110.18
16	N	301	VNK	C13-C12-C11	3.75	123.32	115.84
15	V	301	ONK	C8-C7-C6	-2.30	108.86	113.16

There are no chirality outliers.

5 of 89 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
15	H	301	ONK	C8-C9-N5-C24
15	H	301	ONK	C8-C9-N5-C25
15	H	301	ONK	O7-C9-N5-C24
15	H	301	ONK	O7-C9-N5-C25
15	H	301	ONK	N3-C11-C12-C13


There are no ring outliers.


3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	H	301	ONK	2	0
16	N	301	VNK	2	0
16	K	301	VNK	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.34	4 (1%) 72 66	44, 62, 91, 106	0
1	O	250/250 (100%)	-0.36	5 (2%) 65 56	25, 64, 94, 111	0
2	B	244/244 (100%)	-0.24	4 (1%) 72 66	43, 62, 99, 115	0
2	P	244/244 (100%)	-0.14	6 (2%) 57 47	49, 66, 98, 113	0
3	C	241/241 (100%)	-0.21	7 (2%) 51 41	44, 65, 104, 137	0
3	Q	241/241 (100%)	0.10	15 (6%) 20 13	54, 79, 121, 155	0
4	D	242/242 (100%)	-0.28	5 (2%) 63 54	51, 67, 91, 114	0
4	R	242/242 (100%)	-0.16	7 (2%) 51 41	58, 73, 100, 126	0
5	E	233/233 (100%)	-0.20	5 (2%) 63 54	56, 73, 98, 106	0
5	S	233/233 (100%)	-0.08	4 (1%) 70 63	58, 78, 105, 114	0
6	F	244/244 (100%)	-0.29	2 (0%) 86 81	48, 70, 98, 125	0
6	T	244/244 (100%)	-0.19	4 (1%) 72 66	48, 67, 99, 125	0
7	G	243/243 (100%)	-0.31	3 (1%) 79 73	47, 65, 93, 116	0
7	U	243/243 (100%)	-0.38	2 (0%) 86 81	46, 62, 88, 108	0
8	H	222/222 (100%)	-0.37	2 (0%) 84 80	45, 58, 77, 95	0
8	V	222/222 (100%)	-0.41	2 (0%) 84 80	44, 58, 76, 100	0
9	I	204/204 (100%)	-0.40	1 (0%) 91 88	40, 52, 75, 81	0
9	W	204/204 (100%)	-0.40	1 (0%) 91 88	41, 53, 75, 82	0
10	J	198/198 (100%)	-0.35	3 (1%) 73 68	41, 54, 77, 99	0
10	X	198/198 (100%)	-0.41	4 (2%) 65 56	44, 55, 77, 97	0
11	K	212/212 (100%)	-0.48	2 (0%) 84 80	38, 52, 73, 81	0
11	Y	212/212 (100%)	-0.46	0 100 100	41, 55, 77, 85	0
12	L	222/222 (100%)	-0.50	0 100 100	39, 56, 72, 81	0
12	Z	222/222 (100%)	-0.42	0 100 100	40, 57, 78, 89	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	233/233 (100%)	-0.42	2 (0%) 84 80	42, 58, 79, 95	0
13	a	233/233 (100%)	-0.49	1 (0%) 92 91	39, 56, 77, 91	0
14	N	196/196 (100%)	-0.46	1 (0%) 91 88	45, 54, 76, 82	0
14	b	196/196 (100%)	-0.38	0 100 100	44, 53, 75, 82	0
All	All	6368/6368 (100%)	-0.32	92 (1%) 75 70	25, 61, 96, 155	0

The worst 5 of 92 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	219	ALA	7.0
1	A	1	MET	6.5
6	T	1	GLY	6.3
2	P	220	ASN	6.0
4	D	120	SER	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
16	VNK	N	301	36/36	0.89	0.27	56,67,72,73	0
16	VNK	b	301	36/36	0.90	0.29	56,67,72,72	0
15	ONK	H	301	36/36	0.92	0.20	53,55,60,60	0
15	ONK	V	301	36/36	0.93	0.23	53,55,59,59	0
16	VNK	K	301	36/36	0.94	0.19	43,47,53,54	0
16	VNK	Y	301	36/36	0.94	0.19	43,49,54,55	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around VNK b 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ONK H 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ONK V 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around VNK K 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.