

# Full wwPDB X-ray Structure Validation Report (i)

Jun 14, 2023 – 08:08 PM JST

PDB ID : 8HP5

Title : Crystal structure of (S)-2-haloacid dehalogenase Authors : Yang, Q.; Wang, L.; Xu, X.; Xing, X.; Zhou, J.

Deposited on : 2022-12-12

Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.33

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


 $\begin{tabular}{lll} Validation Pipeline (wwPDB-VP) & : & 2.33 \end{tabular}$ 

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	289	80%	·	17%
1	В	289	77%	6%	17%
1	С	289	75%	8%	17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	С	301	-	-	X	-



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called (S)-2-haloacid dehalogenase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	۸	240	Total	С	N	О	S	0	1	0	
1	A	240	1960	1238	342	370	10	0	1		
1	D	240	Total	С	N	О	S	0	0	0 0	0
1	Б	240	1951	1233	341	367	10	U		0	
1	C	240	Total	С	N	О	S	0	2	0	
1		240	1969	1243	344	372	10				

There are 141 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	initiating methionine	UNP A0A0S6WSA3
A	-32	GLY	-	expression tag	UNP A0A0S6WSA3
A	-31	SER	-	expression tag	UNP A0A0S6WSA3
A	-30	SER	-	expression tag	UNP A0A0S6WSA3
A	-29	HIS	-	expression tag	UNP A0A0S6WSA3
A	-28	HIS	-	expression tag	UNP A0A0S6WSA3
A	-27	HIS	-	expression tag	UNP A0A0S6WSA3
A	-26	HIS	-	expression tag	UNP A0A0S6WSA3
A	-25	HIS	_	expression tag	UNP A0A0S6WSA3
A	-24	HIS	-	expression tag	UNP A0A0S6WSA3
A	-23	SER	-	expression tag	UNP A0A0S6WSA3
A	-22	SER	-	expression tag	UNP A0A0S6WSA3
A	-21	GLY	-	expression tag	UNP A0A0S6WSA3
A	-20	LEU	-	expression tag	UNP A0A0S6WSA3
A	-19	VAL	-	expression tag	UNP A0A0S6WSA3
A	-18	PRO	_	expression tag	UNP A0A0S6WSA3
A	-17	ARG	-	expression tag	UNP A0A0S6WSA3
A	-16	GLY	-	expression tag	UNP A0A0S6WSA3
A	-15	SER	-	expression tag	UNP A0A0S6WSA3
A	-14	HIS	-	expression tag	UNP A0A0S6WSA3
A	-13	MET	-	expression tag	UNP A0A0S6WSA3
A	-12	ALA	-	expression tag	UNP A0A0S6WSA3
A	-11	SER	-	expression tag	UNP A0A0S6WSA3



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
A	-10	MET	-	expression tag	UNP A0A0S6WSA3
A	-9	THR	-	expression tag	UNP A0A0S6WSA3
A	-8	GLY	-	expression tag	UNP A0A0S6WSA3
A	-7	GLY	-	expression tag	UNP A0A0S6WSA3
A	-6	GLN	-	expression tag	UNP A0A0S6WSA3
A	-5	GLN	-	expression tag	UNP A0A0S6WSA3
A	-4	MET	-	expression tag	UNP A0A0S6WSA3
A	-3	GLY	-	expression tag	UNP A0A0S6WSA3
A	-2	ARG	-	expression tag	UNP A0A0S6WSA3
A	-1	GLY	-	expression tag	UNP A0A0S6WSA3
A	0	SER	-	expression tag	UNP A0A0S6WSA3
A	243	LYS	-	expression tag	UNP A0A0S6WSA3
A	244	LEU	-	expression tag	UNP A0A0S6WSA3
A	245	ALA	-	expression tag	UNP A0A0S6WSA3
A	246	ALA	-	expression tag	UNP A0A0S6WSA3
A	247	ALA	-	expression tag	UNP A0A0S6WSA3
A	248	LEU	-	expression tag	UNP A0A0S6WSA3
A	249	GLU	-	expression tag	UNP A0A0S6WSA3
A	250	HIS	-	expression tag	UNP A0A0S6WSA3
A	251	HIS	-	expression tag	UNP A0A0S6WSA3
A	252	HIS	-	expression tag	UNP A0A0S6WSA3
A	253	HIS	-	expression tag	UNP A0A0S6WSA3
A	254	HIS	-	expression tag	UNP A0A0S6WSA3
A	255	HIS	-	expression tag	UNP A0A0S6WSA3
В	-33	MET	-	initiating methionine	UNP A0A0S6WSA3
В	-32	GLY	-	expression tag	UNP A0A0S6WSA3
В	-31	SER	-	expression tag	UNP A0A0S6WSA3
В	-30	SER	-	expression tag	UNP A0A0S6WSA3
В	-29	HIS	-	expression tag	UNP A0A0S6WSA3
В	-28	HIS	-	expression tag	UNP A0A0S6WSA3
В	-27	HIS	-	expression tag	UNP A0A0S6WSA3
В	-26	HIS	-	expression tag	UNP A0A0S6WSA3
В	-25	HIS	-	expression tag	UNP A0A0S6WSA3
В	-24	HIS	-	expression tag	UNP A0A0S6WSA3
В	-23	SER	-	expression tag	UNP A0A0S6WSA3
В	-22	SER	-	expression tag	UNP A0A0S6WSA3
В	-21	GLY	-	expression tag	UNP A0A0S6WSA3
В	-20	LEU	-	expression tag	UNP A0A0S6WSA3
В	-19	VAL	-	expression tag	UNP A0A0S6WSA3
В	-18	PRO	-	expression tag	UNP A0A0S6WSA3
В	-17	ARG	-	expression tag	UNP A0A0S6WSA3
В	-16	GLY	-	expression tag	UNP A0A0S6WSA3



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
В	-15	SER	-	expression tag	UNP A0A0S6WSA3
В	-14	HIS	-	expression tag	UNP A0A0S6WSA3
В	-13	MET	-	expression tag	UNP A0A0S6WSA3
В	-12	ALA	-	expression tag	UNP A0A0S6WSA3
В	-11	SER	-	expression tag	UNP A0A0S6WSA3
В	-10	MET	-	expression tag	UNP A0A0S6WSA3
В	-9	THR	-	expression tag	UNP A0A0S6WSA3
В	-8	GLY	-	expression tag	UNP A0A0S6WSA3
В	-7	GLY	-	expression tag	UNP A0A0S6WSA3
В	-6	GLN	-	expression tag	UNP A0A0S6WSA3
В	-5	GLN	-	expression tag	UNP A0A0S6WSA3
В	-4	MET	-	expression tag	UNP A0A0S6WSA3
В	-3	GLY	-	expression tag	UNP A0A0S6WSA3
В	-2	ARG	-	expression tag	UNP A0A0S6WSA3
В	-1	GLY	_	expression tag	UNP A0A0S6WSA3
В	0	SER	-	expression tag	UNP A0A0S6WSA3
В	243	LYS	-	expression tag	UNP A0A0S6WSA3
В	244	LEU	-	expression tag	UNP A0A0S6WSA3
В	245	ALA	-	expression tag	UNP A0A0S6WSA3
В	246	ALA	-	expression tag	UNP A0A0S6WSA3
В	247	ALA	-	expression tag	UNP A0A0S6WSA3
В	248	LEU	-	expression tag	UNP A0A0S6WSA3
В	249	GLU	-	expression tag	UNP A0A0S6WSA3
В	250	HIS	-	expression tag	UNP A0A0S6WSA3
В	251	HIS	-	expression tag	UNP A0A0S6WSA3
В	252	HIS	-	expression tag	UNP A0A0S6WSA3
В	253	HIS	-	expression tag	UNP A0A0S6WSA3
В	254	HIS	-	expression tag	UNP A0A0S6WSA3
В	255	HIS	-	expression tag	UNP A0A0S6WSA3
С	-33	MET	-	initiating methionine	UNP A0A0S6WSA3
С	-32	GLY	-	expression tag	UNP A0A0S6WSA3
С	-31	SER	-	expression tag	UNP A0A0S6WSA3
С	-30	SER	-	expression tag	UNP A0A0S6WSA3
С	-29	HIS	-	expression tag	UNP A0A0S6WSA3
С	-28	HIS	-	expression tag	UNP A0A0S6WSA3
С	-27	HIS	-	expression tag	UNP A0A0S6WSA3
С	-26	HIS	-	expression tag	UNP A0A0S6WSA3
С	-25	HIS	-	expression tag	UNP A0A0S6WSA3
С	-24	HIS	-	expression tag	UNP A0A0S6WSA3
С	-23	SER	-	expression tag	UNP A0A0S6WSA3
С	-22	SER	-	expression tag	UNP A0A0S6WSA3
С	-21	GLY	-	expression tag	UNP A0A0S6WSA3




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	-20	LEU	-	expression tag	UNP A0A0S6WSA3
С	-19	VAL	-	expression tag	UNP A0A0S6WSA3
С	-18	PRO	-	expression tag	UNP A0A0S6WSA3
С	-17	ARG	-	expression tag	UNP A0A0S6WSA3
С	-16	GLY	-	expression tag	UNP A0A0S6WSA3
С	-15	SER	-	expression tag	UNP A0A0S6WSA3
С	-14	HIS	-	expression tag	UNP A0A0S6WSA3
С	-13	MET	-	expression tag	UNP A0A0S6WSA3
С	-12	ALA	-	expression tag	UNP A0A0S6WSA3
С	-11	SER	-	expression tag	UNP A0A0S6WSA3
С	-10	MET	-	expression tag	UNP A0A0S6WSA3
С	-9	THR	-	expression tag	UNP A0A0S6WSA3
С	-8	GLY	-	expression tag	UNP A0A0S6WSA3
С	-7	GLY	-	expression tag	UNP A0A0S6WSA3
С	-6	GLN	-	expression tag	UNP A0A0S6WSA3
С	-5	GLN	-	expression tag	UNP A0A0S6WSA3
С	-4	MET	-	expression tag	UNP A0A0S6WSA3
С	-3	GLY	-	expression tag	UNP A0A0S6WSA3
С	-2	ARG	-	expression tag	UNP A0A0S6WSA3
С	-1	GLY	-	expression tag	UNP A0A0S6WSA3
С	0	SER	-	expression tag	UNP A0A0S6WSA3
С	243	LYS	-	expression tag	UNP A0A0S6WSA3
С	244	LEU	-	expression tag	UNP A0A0S6WSA3
С	245	ALA	-	expression tag	UNP A0A0S6WSA3
С	246	ALA	-	expression tag	UNP A0A0S6WSA3
С	247	ALA	-	expression tag	UNP A0A0S6WSA3
С	248	LEU	-	expression tag	UNP A0A0S6WSA3
С	249	GLU	-	expression tag	UNP A0A0S6WSA3
С	250	HIS	-	expression tag	UNP A0A0S6WSA3
С	251	HIS	-	expression tag	UNP A0A0S6WSA3
С	252	HIS	-	expression tag	UNP A0A0S6WSA3
С	253	HIS	-	expression tag	UNP A0A0S6WSA3
С	254	HIS	-	expression tag	UNP A0A0S6WSA3
С	255	HIS	-	expression tag	UNP A0A0S6WSA3

 $\bullet$  Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $\mathrm{C_2H_6O_2}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	В	1	Total C O 4 2 2	0	0
2	С	1	Total C O 4 2 2	0	0

#### • Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	195	Total O 195 195	0	0
3	В	125	Total O 125 125	0	0
3	С	112	Total O 112 112	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: (S)-2-haloacid dehalogenase





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants	151.75Å 151.75Å 151.75Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	42.09 - 2.50	Depositor
Resolution (A)	45.76 - 2.50	EDS
% Data completeness	99.5 (42.09-2.50)	Depositor
(in resolution range)	99.6 (45.76-2.50)	EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.75  (at  2.51Å)	Xtriage
Refinement program	PHENIX 1.19.2-4158	Depositor
D D	0.166 , 0.216	Depositor
$R, R_{free}$	0.166 , $0.214$	DCC
$R_{free}$ test set	2018 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.1	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.37, 48.2	EDS
L-test for twinning <sup>2</sup>	$< L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.032 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6324	wwPDB-VP
Average B, all atoms $(\mathring{A}^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 33.11 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.4286e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

### 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.42	0/2012	0.61	0/2720	
1	В	0.40	0/2003	0.61	0/2708	
1	С	0.43	0/2021	0.62	0/2732	
All	All	0.42	0/6036	0.61	0/8160	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1960	0	1860	7	0
1	В	1951	0	1855	10	0
1	С	1969	0	1867	14	0
2	A	4	0	6	3	0
2	В	4	0	6	1	0
2	С	4	0	6	4	0
3	A	195	0	0	1	0
3	В	125	0	0	2	0
3	С	112	0	0	1	0
All	All	6324	0	5600	31	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:122:ASN:HB2	2:A:301:EDO:H21	1.81	0.62
1:C:14:TYR:HB2	2:C:301:EDO:H22	1.82	0.61
1:C:34:ARG:NH2	1:C:87:GLU:OE2	2.33	0.61
1:C:122:ASN:HB2	2:C:301:EDO:H12	1.85	0.59
1:C:122:ASN:H	2:C:301:EDO:H12	1.70	0.57
1:A:122:ASN:H	2:A:301:EDO:H21	1.71	0.55
1:C:189:ALA:HB1	1:C:194:MET:HB2	1.90	0.53
1:B:93:ARG:O	1:B:96:ARG:HG2	2.12	0.49
1:A:129:GLN:O	1:A:133:GLU:HG3	2.13	0.49
1:B:11:PHE:HE2	1:B:117:LEU:HD22	1.78	0.48
1:A:183:PHE:CD2	1:A:212:MET:HB3	2.48	0.48
1:B:1:MET:N	3:B:405:HOH:O	2.40	0.47
1:B:46:GLN:NE2	1:B:209:GLY:O	2.47	0.47
1:A:122:ASN:H	2:A:301:EDO:C2	2.29	0.46
1:B:14:TYR:HB2	2:B:301:EDO:H11	1.97	0.46
1:B:62:LYS:HE2	1:B:62:LYS:HB2	1.73	0.45
1:C:198:TRP:CE3	1:C:223:ARG:HD2	2.52	0.45
1:C:9:LEU:HD23	1:C:176:LEU:HB3	1.99	0.44
1:C:141:ALA:HB2	1:C:168:LEU:HD11	1.98	0.44
1:B:181:SER:HB2	1:B:211:THR:HG21	2.00	0.44
1:A:9:LEU:HD23	1:A:176:LEU:HB3	2.01	0.43
1:C:186:HIS:ND1	1:C:196:ASN:HB2	2.33	0.43
1:C:235[A]:GLN:NE2	3:C:414:HOH:O	2.50	0.43
1:C:144:SER:O	1:C:148:VAL:HG23	2.19	0.43
1:C:220:TYR:CE1	1:C:223:ARG:HB2	2.55	0.41
1:B:182:LEU:HD23	1:B:186:HIS:HD2	1.85	0.41
1:C:96:ARG:HG2	1:C:130:TYR:CD2	2.55	0.41
1:B:41:ARG:O	1:B:45:LEU:HG	2.20	0.41
1:B:194:MET:HA	3:B:425:HOH:O	2.21	0.41
1:C:122:ASN:H	2:C:301:EDO:C1	2.33	0.41
1:A:46:GLN:NE2	3:A:414:HOH:O	2.53	0.40

There are no symmetry-related clashes.



#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	239/289 (83%)	234 (98%)	5 (2%)	0	100	100
1	В	238/289 (82%)	234 (98%)	4 (2%)	0	100	100
1	С	240/289 (83%)	234 (98%)	6 (2%)	0	100	100
All	All	717/867 (83%)	702 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	205/241~(85%)	203 (99%)	2 (1%)	76 90
1	В	204/241 (85%)	202 (99%)	2 (1%)	76 90
1	С	206/241 (86%)	204 (99%)	2 (1%)	76 90
All	All	615/723~(85%)	609 (99%)	6 (1%)	76 90

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	TRP
1	A	29	ARG
1	В	12	ASP
1	В	20	TRP
1	С	12	ASP



Continued from previous page...

Mol	Chain	Res	Type
1	С	20	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		e Chain Res		es Link	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
2	EDO	В	301	-	3,3,3	0.72	0	2,2,2	0.43	0
2	EDO	A	301	-	3,3,3	0.81	0	2,2,2	0.30	0
2	EDO	С	301	-	3,3,3	0.78	0	2,2,2	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	В	301	-	-	1/1/1/1	-
2	EDO	A	301	-	=	1/1/1/1	-
2	EDO	С	301	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	301	EDO	O1-C1-C2-O2
2	В	301	EDO	O1-C1-C2-O2
2	A	301	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	301	EDO	1	0
2	A	301	EDO	3	0
2	С	301	EDO	4	0

### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	240/289 (83%)	-0.63	1 (0%) 92 93	16, 23, 40, 64	0
1	В	240/289 (83%)	-0.31	9 (3%) 40 43	24, 32, 57, 78	0
1	С	240/289 (83%)	-0.27	13 (5%) 25 27	22, 30, 62, 98	0
All	All	720/867 (83%)	-0.40	23 (3%) 47 51	16, 29, 55, 98	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	36	GLY	4.3
1	С	217	GLU	4.2
1	С	216	HIS	4.0
1	С	206	GLU	3.7
1	С	36	GLY	3.6
1	С	37	THR	3.2
1	С	193	GLY	3.2
1	В	2	ARG	3.1
1	С	38	ASP	3.0
1	В	216	HIS	3.0
1	В	240	ASN	2.8
1	В	1	MET	2.8
1	В	38	ASP	2.8
1	В	37	THR	2.8
1	В	239	ARG	2.7
1	A	36	GLY	2.4
1	В	215	SER	2.4
1	С	213	THR	2.3
1	С	208	PHE	2.2
1	С	209	GLY	2.2
1	С	215	SER	2.1
1	С	1	MET	2.0
1	С	207	GLY	2.0



### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
2	EDO	С	301	4/4	0.90	0.15	26,29,40,41	0
2	EDO	В	301	4/4	0.94	0.20	27,37,42,45	0
2	EDO	A	301	4/4	0.96	0.23	18,23,29,32	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

