

# wwPDB X-ray Structure Validation Summary Report (i)

#### Jun 14, 2023 – 08:09 PM JST

PDB ID	:	8HP6
Title	:	Crystal structure of (S)-2-haloacid dehalogenase D12A mutant
Authors	:	Yang, Q.; Wang, L.; Xu, X.; Xing, X.; Zhou, J.
Deposited on		
Resolution	:	2.20  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.33
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.33

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{l} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	289	<b>78%</b> 6%	17%
1	В	289	% • 79% •	17%
1	С	289	80%	17%



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	240	Total	С	Ν	0	S	0	0	0
	A		1948	1232	341	365	10	0		
1	D	240	Total	С	Ν	0	S	0	1	0
	D	240	1956	1236	343	367	10			
1	C	240	Total	С	Ν	0	S	0	9	0
		240	1965	1241	344	370	10	U		0

• Molecule 1 is a protein called (S)-2-haloacid dehalogenase.

	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	initiating methionine	UNP A0A0S6WSA3
A	-32	GLY	-	expression tag	UNP A0A0S6WSA3
A	-31	SER	-	expression tag	UNP A0A0S6WSA3
A	-30	SER	-	expression tag	UNP A0A0S6WSA3
A	-29	HIS	-	expression tag	UNP A0A0S6WSA3
A	-28	HIS	-	expression tag	UNP A0A0S6WSA3
A	-27	HIS	-	expression tag	UNP A0A0S6WSA3
A	-26	HIS	-	expression tag	UNP A0A0S6WSA3
A	-25	HIS	-	expression tag	UNP A0A0S6WSA3
A	-24	HIS	-	expression tag	UNP A0A0S6WSA3
A	-23	SER	-	expression tag	UNP A0A0S6WSA3
A	-22	SER	-	expression tag	UNP A0A0S6WSA3
A	-21	GLY	-	expression tag	UNP A0A0S6WSA3
A	-20	LEU	-	expression tag	UNP A0A0S6WSA3
A	-19	VAL	-	expression tag	UNP A0A0S6WSA3
A	-18	PRO	-	expression tag	UNP A0A0S6WSA3
A	-17	ARG	-	expression tag	UNP A0A0S6WSA3
A	-16	GLY	-	expression tag	UNP A0A0S6WSA3
A	-15	SER	-	expression tag	UNP A0A0S6WSA3
A	-14	HIS	-	expression tag	UNP A0A0S6WSA3
A	-13	MET	-	expression tag	UNP A0A0S6WSA3
A	-12	ALA	-	expression tag	UNP A0A0S6WSA3
A	-11	SER	-	expression tag	UNP A0A0S6WSA3

There are 144 discrepancies between the modelled and reference sequences:

Continued on next page...



Chain	Residue	vious page Modelled	Actual	Comment	Reference
А	-10	MET	-	expression tag	UNP A0A0S6WSA3
А	-9	THR	-	expression tag	UNP A0A0S6WSA3
А	-8	GLY	-	expression tag	UNP A0A0S6WSA3
А	-7	GLY	-	expression tag	UNP A0A0S6WSA3
А	-6	GLN	-	expression tag	UNP A0A0S6WSA3
А	-5	GLN	-	expression tag	UNP A0A0S6WSA3
А	-4	MET	-	expression tag	UNP A0A0S6WSA3
А	-3	GLY	-	expression tag	UNP A0A0S6WSA3
А	-2	ARG	-	expression tag	UNP A0A0S6WSA3
А	-1	GLY	_	expression tag	UNP A0A0S6WSA3
А	0	SER	-	expression tag	UNP A0A0S6WSA3
А	12	ALA	ASP	engineered mutation	UNP A0A0S6WSA3
А	243	LYS	_	expression tag	UNP A0A0S6WSA3
А	244	LEU	-	expression tag	UNP A0A0S6WSA3
А	245	ALA	-	expression tag	UNP A0A0S6WSA3
А	246	ALA	-	expression tag	UNP A0A0S6WSA3
А	247	ALA	-	expression tag	UNP A0A0S6WSA3
А	248	LEU	-	expression tag	UNP A0A0S6WSA3
А	249	GLU	-	expression tag	UNP A0A0S6WSA3
А	250	HIS	-	expression tag	UNP A0A0S6WSA3
А	251	HIS	-	expression tag	UNP A0A0S6WSA3
А	252	HIS	-	expression tag	UNP A0A0S6WSA3
А	253	HIS	-	expression tag	UNP A0A0S6WSA3
А	254	HIS	_	expression tag	UNP A0A0S6WSA3
А	255	HIS	-	expression tag	UNP A0A0S6WSA3
В	-33	MET	-	initiating methionine	UNP A0A0S6WSA3
В	-32	GLY	-	expression tag	UNP A0A0S6WSA3
В	-31	SER	-	expression tag	UNP A0A0S6WSA3
В	-30	SER	-	expression tag	UNP A0A0S6WSA3
В	-29	HIS	-	expression tag	UNP A0A0S6WSA3
В	-28	HIS	-	expression tag	UNP A0A0S6WSA3
В	-27	HIS	_	expression tag	UNP A0A0S6WSA3
В	-26	HIS	-	expression tag	UNP A0A0S6WSA3
В	-25	HIS	-	expression tag	UNP A0A0S6WSA3
В	-24	HIS	-	expression tag	UNP A0A0S6WSA3
В	-23	SER	-	expression tag	UNP A0A0S6WSA3
В	-22	SER	_	expression tag	UNP A0A0S6WSA3
В	-21	GLY	-	expression tag	UNP A0A0S6WSA3
В	-20	LEU	-	expression tag	UNP A0A0S6WSA3
В	-19	VAL	-	expression tag	UNP A0A0S6WSA3
В	-18	PRO	_	expression tag	UNP A0A0S6WSA3
В	-17	ARG	-	expression tag	UNP A0A0S6WSA3

Continued on next page...



8HP6
------

	Continued from previous page							
Chain	Residue	Modelled	Actual	Comment	Reference			
В	-16	GLY	-	expression tag	UNP A0A0S6WSA3			
В	-15	SER	-	expression tag	UNP A0A0S6WSA3			
В	-14	HIS	-	expression tag	UNP A0A0S6WSA3			
В	-13	MET	-	expression tag	UNP A0A0S6WSA3			
В	-12	ALA	-	expression tag	UNP A0A0S6WSA3			
В	-11	SER	-	expression tag	UNP A0A0S6WSA3			
В	-10	MET	-	expression tag	UNP A0A0S6WSA3			
В	-9	THR	-	expression tag	UNP A0A0S6WSA3			
В	-8	GLY	-	expression tag	UNP A0A0S6WSA3			
В	-7	GLY	-	expression tag	UNP A0A0S6WSA3			
В	-6	GLN	-	expression tag	UNP A0A0S6WSA3			
В	-5	GLN	-	expression tag	UNP A0A0S6WSA3			
В	-4	MET	-	expression tag	UNP A0A0S6WSA3			
В	-3	GLY	-	expression tag	UNP A0A0S6WSA3			
В	-2	ARG	-	expression tag	UNP A0A0S6WSA3			
В	-1	GLY	-	expression tag	UNP A0A0S6WSA3			
В	0	SER	_	expression tag	UNP A0A0S6WSA3			
В	12	ALA	ASP	engineered mutation	UNP A0A0S6WSA3			
В	243	LYS	-	expression tag	UNP A0A0S6WSA3			
В	244	LEU	-	expression tag	UNP A0A0S6WSA3			
В	245	ALA	-	expression tag	UNP A0A0S6WSA3			
В	246	ALA	-	expression tag	UNP A0A0S6WSA3			
В	247	ALA	-	expression tag	UNP A0A0S6WSA3			
В	248	LEU	-	expression tag	UNP A0A0S6WSA3			
В	249	GLU	-	expression tag	UNP A0A0S6WSA3			
В	250	HIS	_	expression tag	UNP A0A0S6WSA3			
В	251	HIS	-	expression tag	UNP A0A0S6WSA3			
В	252	HIS	-	expression tag	UNP A0A0S6WSA3			
В	253	HIS	-	expression tag	UNP A0A0S6WSA3			
В	254	HIS	-	expression tag	UNP A0A0S6WSA3			
В	255	HIS	-	expression tag	UNP A0A0S6WSA3			
С	-33	MET	_	initiating methionine	UNP A0A0S6WSA3			
С	-32	GLY	_	expression tag	UNP A0A0S6WSA3			
С	-31	SER	_	expression tag	UNP A0A0S6WSA3			
С	-30	SER	_	expression tag	UNP A0A0S6WSA3			
С	-29	HIS	-	expression tag	UNP A0A0S6WSA3			
С	-28	HIS	-	expression tag	UNP A0A0S6WSA3			
С	-27	HIS	-	expression tag	UNP A0A0S6WSA3			
С	-26	HIS	_	expression tag	UNP A0A0S6WSA3			
С	-25	HIS	-	expression tag	UNP A0A0S6WSA3			
С	-24	HIS	-	expression tag	UNP A0A0S6WSA3			
С	-23	SER	-	expression tag	UNP A0A0S6WSA3			

Continued on next page...



8HP6
------

Chain	Residue	vious page Modelled	Actual	Comment	Reference
С	-22	SER	-	expression tag	UNP A0A0S6WSA3
С	-21	GLY	-	expression tag	UNP A0A0S6WSA3
С	-20	LEU	_	expression tag	UNP A0A0S6WSA3
С	-19	VAL	-	expression tag	UNP A0A0S6WSA3
С	-18	PRO	-	expression tag	UNP A0A0S6WSA3
С	-17	ARG	-	expression tag	UNP A0A0S6WSA3
С	-16	GLY	-	expression tag	UNP A0A0S6WSA3
С	-15	SER	-	expression tag	UNP A0A0S6WSA3
С	-14	HIS	-	expression tag	UNP A0A0S6WSA3
С	-13	MET	-	expression tag	UNP A0A0S6WSA3
С	-12	ALA	-	expression tag	UNP A0A0S6WSA3
С	-11	SER	-	expression tag	UNP A0A0S6WSA3
С	-10	MET	-	expression tag	UNP A0A0S6WSA3
С	-9	THR	-	expression tag	UNP A0A0S6WSA3
$\mathbf{C}$	-8	GLY	-	expression tag	UNP A0A0S6WSA3
С	-7	GLY	-	expression tag	UNP A0A0S6WSA3
С	-6	GLN	-	expression tag	UNP A0A0S6WSA3
С	-5	GLN	-	expression tag	UNP A0A0S6WSA3
С	-4	MET	-	expression tag	UNP A0A0S6WSA3
С	-3	GLY	-	expression tag	UNP A0A0S6WSA3
С	-2	ARG	-	expression tag	UNP A0A0S6WSA3
С	-1	GLY	-	expression tag	UNP A0A0S6WSA3
С	0	SER	-	expression tag	UNP A0A0S6WSA3
С	12	ALA	ASP	engineered mutation	UNP A0A0S6WSA3
С	243	LYS	-	expression tag	UNP A0A0S6WSA3
С	244	LEU	-	expression tag	UNP A0A0S6WSA3
С	245	ALA	-	expression tag	UNP A0A0S6WSA3
С	246	ALA	-	expression tag	UNP A0A0S6WSA3
С	247	ALA	-	expression tag	UNP A0A0S6WSA3
С	248	LEU	-	expression tag	UNP A0A0S6WSA3
С	249	GLU	-	expression tag	UNP A0A0S6WSA3
С	250	HIS	-	expression tag	UNP A0A0S6WSA3
С	251	HIS	-	expression tag	UNP A0A0S6WSA3
С	252	HIS	-	expression tag	UNP A0A0S6WSA3
С	253	HIS	-	expression tag	UNP A0A0S6WSA3
С	254	HIS	-	expression tag	UNP A0A0S6WSA3
С	255	HIS	-	expression tag	UNP A0A0S6WSA3

• Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na). Continued on next page...



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Na 1 1	0	0
2	В	1	Total Na 1 1	0	0
2	С	1	Total Na 1 1	0	0


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	267	Total O 267 267	0	0
3	В	188	Total O 188 188	0	0
3	С	207	Total         O           207         207	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: (S)-2-haloacid dehalogenase



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants	151.73Å 151.73Å 151.73Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	36.80 - 2.20	Depositor
Resolution (A)	43.80 - 2.20	EDS
% Data completeness	100.0 (36.80-2.20)	Depositor
(in resolution range)	$100.0 \ (43.80-2.20)$	EDS
R <sub>merge</sub>	0.12	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.20 (at 2.20 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
D D.	0.166 , $0.201$	Depositor
$R, R_{free}$	0.167 , $0.201$	DCC
$R_{free}$ test set	2914 reflections $(4.93\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	35.1	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, $35.9$	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.027 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6534	wwPDB-VP
Average B, all atoms $(Å^2)$	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 36.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.3165e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.43	0/2000	0.64	0/2704	
1	В	0.45	0/2008	0.64	0/2715	
1	С	0.42	0/2017	0.60	0/2727	
All	All	0.44	0/6025	0.63	0/8146	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1948	0	1856	9	0
1	В	1956	0	1860	5	0
1	С	1965	0	1866	5	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
3	А	267	0	0	0	0
3	В	188	0	0	0	0
3	С	207	0	0	0	0
All	All	6534	0	5582	17	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:HIS:HB2	1:B:194:MET:HE2	1.93	0.51
1:A:34:ARG:NH2	1:A:87:GLU:OE2	2.40	0.50
1:B:34:ARG:NH2	1:B:87:GLU:OE2	2.41	0.49
1:A:66:ASP:C	1:A:69:PRO:HD2	2.34	0.47
1:B:66:ASP:C	1:B:69:PRO:HD2	2.35	0.46

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	238/289~(82%)	235~(99%)	3~(1%)	0	100	100
1	В	239/289~(83%)	235~(98%)	4 (2%)	0	100	100
1	С	240/289~(83%)	235~(98%)	5 (2%)	0	100	100
All	All	717/867~(83%)	705 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	203/240~(85%)	202 (100%)	1 (0%)	88 94		
1	В	204/240~(85%)	202~(99%)	2(1%)	76 86		
1	С	205/240~(85%)	204 (100%)	1 (0%)	88 94		
All	All	612/720~(85%)	608~(99%)	4 (1%)	84 91		

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	20	TRP
1	В	20	TRP
1	В	38	ASP
1	С	20	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	240/289~(83%)	-0.53	3 (1%) 77 75	26, 33, 50, 74	0
1	В	240/289~(83%)	-0.57	3 (1%) 77 75	28, 37, 53, 83	0
1	С	240/289~(83%)	-0.35	7 (2%) 51 49	28, 38, 60, 93	0
All	All	720/867~(83%)	-0.48	13 (1%) 68 66	26, 35, 55, 93	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	38	ASP	4.6
1	В	38	ASP	4.2
1	С	36	GLY	3.9
1	В	37	THR	3.9
1	В	36	GLY	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
2	NA	А	301	1/1	0.99	0.33	28,28,28,28	0
2	NA	В	301	1/1	0.99	0.21	30,30,30,30	0
2	NA	С	301	1/1	0.99	0.32	30,30,30,30	0

## 6.5 Other polymers (i)

There are no such residues in this entry.

