



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 26, 2023 – 08:10 PM JST

PDB ID : 8HS6  
Title : Brucella melitensis 7-alpha-Hydroxysteroid Dehydrogenase mutant:1-53 truncation  
Authors : Liu, Z.Y.; Zhang, R.Z.  
Deposited on : 2022-12-17  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	 2% 78% 14% 7%
1	B	249	 83% 14% 2%
1	C	249	 80% 12% 8%
1	D	249	 90% 9% 1%
1	E	249	 85% 14% 1% 1%
1	F	249	 81% 10% 9%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	249	 2% 78% 13% 9%
1	H	249	 % 83% 17%
1	I	249	 2% 66% 24% 9%
1	J	249	 % 68% 21% 11%
1	K	249	 % 69% 29% .
1	L	249	 3% 68% 27% . .

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 20848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 7-alpha-hydroxysteroid dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	232	Total 1656	C 1041	N 297	O 310	S 8	0	0	0
1	B	245	Total 1766	C 1112	N 316	O 330	S 8	0	0	0
1	C	229	Total 1633	C 1027	N 292	O 306	S 8	0	0	0
1	D	247	Total 1773	C 1118	N 317	O 330	S 8	0	0	0
1	E	248	Total 1782	C 1123	N 319	O 332	S 8	0	0	0
1	F	227	Total 1629	C 1026	N 293	O 302	S 8	0	0	0
1	G	227	Total 1623	C 1023	N 291	O 301	S 8	0	0	0
1	H	249	Total 1791	C 1128	N 320	O 335	S 8	0	0	0
1	I	226	Total 1612	C 1013	N 290	O 301	S 8	0	0	0
1	J	221	Total 1577	C 993	N 281	O 295	S 8	0	0	0
1	K	246	Total 1766	C 1113	N 316	O 329	S 8	0	0	0
1	L	239	Total 1718	C 1083	N 309	O 318	S 8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	258	ILE	MET	engineered mutation	UNP Q8YIN7
A	262	LYS	THR	engineered mutation	UNP Q8YIN7
B	258	ILE	MET	engineered mutation	UNP Q8YIN7
B	262	LYS	THR	engineered mutation	UNP Q8YIN7
C	258	ILE	MET	engineered mutation	UNP Q8YIN7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	262	LYS	THR	engineered mutation	UNP Q8YIN7
D	258	ILE	MET	engineered mutation	UNP Q8YIN7
D	262	LYS	THR	engineered mutation	UNP Q8YIN7
E	258	ILE	MET	engineered mutation	UNP Q8YIN7
E	262	LYS	THR	engineered mutation	UNP Q8YIN7
F	258	ILE	MET	engineered mutation	UNP Q8YIN7
F	262	LYS	THR	engineered mutation	UNP Q8YIN7
G	258	ILE	MET	engineered mutation	UNP Q8YIN7
G	262	LYS	THR	engineered mutation	UNP Q8YIN7
H	258	ILE	MET	engineered mutation	UNP Q8YIN7
H	262	LYS	THR	engineered mutation	UNP Q8YIN7
I	258	ILE	MET	engineered mutation	UNP Q8YIN7
I	262	LYS	THR	engineered mutation	UNP Q8YIN7
J	258	ILE	MET	engineered mutation	UNP Q8YIN7
J	262	LYS	THR	engineered mutation	UNP Q8YIN7
K	258	ILE	MET	engineered mutation	UNP Q8YIN7
K	262	LYS	THR	engineered mutation	UNP Q8YIN7
L	258	ILE	MET	engineered mutation	UNP Q8YIN7
L	262	LYS	THR	engineered mutation	UNP Q8YIN7

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	76	Total O 76 76	0	0
2	B	59	Total O 59 59	0	0
2	C	65	Total O 65 65	0	0
2	D	53	Total O 53 53	0	0
2	E	48	Total O 48 48	0	0
2	F	49	Total O 49 49	0	0
2	G	52	Total O 52 52	0	0
2	H	63	Total O 63 63	0	0
2	I	17	Total O 17 17	0	0
2	J	16	Total O 16 16	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	K	13	Total	O	0	0
			13	13		
2	L	11	Total	O	0	0
			11	11		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase




- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase









- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase



- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase



- Molecule 1: 7-alpha-hydroxysteroid dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.08Å 99.91Å 103.42Å 87.43° 78.84° 78.37°	Depositor
Resolution (Å)	23.75 – 2.35 23.75 – 2.35	Depositor EDS
% Data completeness (in resolution range)	93.4 (23.75-2.35) 93.4 (23.75-2.35)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.79 (at 2.36Å)	Xtrriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, $R_{free}$	0.217 , 0.266 0.219 , 0.266	Depositor DCC
$R_{free}$ test set	6448 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.8	Xtrriage
Anisotropy	0.216	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 18.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.157 for -h,-l,-k	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	20848	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1682	0.63	0/2276
1	B	0.45	0/1794	0.63	0/2428
1	C	0.48	0/1659	0.64	0/2245
1	D	0.46	0/1802	0.65	0/2442
1	E	0.45	0/1811	0.63	0/2454
1	F	0.45	0/1655	0.63	0/2238
1	G	0.48	0/1648	0.67	0/2228
1	H	0.48	0/1820	0.64	0/2466
1	I	0.37	0/1637	0.57	0/2214
1	J	0.35	0/1602	0.56	0/2168
1	K	0.33	0/1795	0.55	0/2432
1	L	0.37	0/1746	0.55	0/2361
All	All	0.43	0/20651	0.61	0/27952

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1656	0	1671	24	0
1	B	1766	0	1789	23	0
1	C	1633	0	1646	20	0
1	D	1773	0	1804	14	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1782	0	1812	22	0
1	F	1629	0	1645	18	0
1	G	1623	0	1643	27	0
1	H	1791	0	1818	28	0
1	I	1612	0	1624	35	0
1	J	1577	0	1588	27	0
1	K	1766	0	1795	47	0
1	L	1718	0	1745	42	0
2	A	76	0	0	3	0
2	B	59	0	0	0	0
2	C	65	0	0	1	0
2	D	53	0	0	0	0
2	E	48	0	0	1	0
2	F	49	0	0	1	0
2	G	52	0	0	0	0
2	H	63	0	0	0	0
2	I	17	0	0	0	0
2	J	16	0	0	0	0
2	K	13	0	0	1	0
2	L	11	0	0	1	0
All	All	20848	0	20580	301	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 301 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:139:THR:HG22	1:I:140:VAL:HG23	1.60	0.82
1:I:203:VAL:HG13	1:I:204:ARG:HG3	1.64	0.80
1:K:257:ALA:HA	1:K:260:LYS:HE3	1.64	0.80
1:C:264:LEU:HD11	1:C:266:ARG:HE	1.51	0.75
1:K:195:SER:HB3	1:K:212:LYS:HG3	1.68	0.75

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/249 (92%)	219 (96%)	9 (4%)	0	100	100
1	B	241/249 (97%)	235 (98%)	6 (2%)	0	100	100
1	C	225/249 (90%)	219 (97%)	6 (3%)	0	100	100
1	D	245/249 (98%)	239 (98%)	6 (2%)	0	100	100
1	E	246/249 (99%)	239 (97%)	7 (3%)	0	100	100
1	F	221/249 (89%)	212 (96%)	9 (4%)	0	100	100
1	G	221/249 (89%)	214 (97%)	7 (3%)	0	100	100
1	H	247/249 (99%)	239 (97%)	8 (3%)	0	100	100
1	I	222/249 (89%)	216 (97%)	6 (3%)	0	100	100
1	J	217/249 (87%)	209 (96%)	8 (4%)	0	100	100
1	K	244/249 (98%)	239 (98%)	5 (2%)	0	100	100
1	L	235/249 (94%)	227 (97%)	8 (3%)	0	100	100
All	All	2792/2988 (93%)	2707 (97%)	85 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/177 (92%)	158 (98%)	4 (2%)	47	58
1	B	175/177 (99%)	169 (97%)	6 (3%)	37	46

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	160/177 (90%)	156 (98%)	4 (2%)	47	58
1	D	175/177 (99%)	173 (99%)	2 (1%)	73	84
1	E	176/177 (99%)	167 (95%)	9 (5%)	24	27
1	F	160/177 (90%)	153 (96%)	7 (4%)	28	34
1	G	159/177 (90%)	153 (96%)	6 (4%)	33	41
1	H	177/177 (100%)	171 (97%)	6 (3%)	37	46
1	I	157/177 (89%)	151 (96%)	6 (4%)	33	41
1	J	154/177 (87%)	148 (96%)	6 (4%)	32	40
1	K	174/177 (98%)	169 (97%)	5 (3%)	42	52
1	L	169/177 (96%)	162 (96%)	7 (4%)	30	37
All	All	1998/2124 (94%)	1930 (97%)	68 (3%)	37	46

5 of 68 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	203	VAL
1	K	283	PRO
1	L	256	ARG
1	E	287	TRP
1	E	267	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	134	GLN
1	I	183	GLN
1	L	181	HIS
1	J	183	GLN
1	B	59	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/249 (93%)	-0.56	4 (1%) 70 78	9, 18, 45, 81	0
1	B	245/249 (98%)	-0.56	0 100 100	11, 19, 51, 80	0
1	C	229/249 (91%)	-0.60	3 (1%) 77 84	10, 17, 38, 93	0
1	D	247/249 (99%)	-0.55	0 100 100	11, 19, 54, 80	0
1	E	248/249 (99%)	-0.52	2 (0%) 86 91	12, 20, 50, 75	0
1	F	227/249 (91%)	-0.61	3 (1%) 77 84	10, 18, 41, 101	0
1	G	227/249 (91%)	-0.57	4 (1%) 68 77	11, 18, 40, 83	0
1	H	249/249 (100%)	-0.57	2 (0%) 86 91	10, 20, 45, 102	0
1	I	226/249 (90%)	-0.17	5 (2%) 62 72	26, 39, 59, 88	0
1	J	221/249 (88%)	-0.23	3 (1%) 75 83	25, 40, 56, 70	0
1	K	246/249 (98%)	0.17	3 (1%) 79 86	36, 53, 67, 82	0
1	L	239/249 (95%)	0.20	7 (2%) 51 62	36, 53, 71, 86	0
All	All	2836/2988 (94%)	-0.38	36 (1%) 77 84	9, 23, 62, 102	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	249	VAL	4.6
1	C	300	VAL	4.5
1	A	244	ASP	4.4
1	C	265	GLY	4.3
1	G	299	GLY	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.