



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 07:21 pm BST

PDB ID : 1HTR  
Title : CRYSTAL AND MOLECULAR STRUCTURES OF HUMAN PRO-  
GASTRICSIN AT 1.62 ANGSTROMS RESOLUTION  
Authors : Moore, S.A.; Sielecki, A.R.; James, M.N.G.  
Deposited on : 1994-10-21  
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	P	43	
2	B	329	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3129 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROGASTRICSIN (PRO SEGMENT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	P	43	362	240	63	58	1	0	0	0

- Molecule 2 is a protein called GASTRICSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	329	2517	1600	390	515	12	0	6	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	P	12	Total	O	0	0
			12	12		
3	B	238	Total	O	0	0
			238	238		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: PROGASTRICSIN (PRO SEGMENT)

Chain P: 



- Molecule 2: GASTRICSIN

Chain B: 



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.31Å 105.31Å 70.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.62	Depositor
% Data completeness (in resolution range)	96.0 (20.00-1.62)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ, TNT	Depositor
R, $R_{free}$	0.179 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3129	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	P	1.03	0/371	1.67	7/494 (1.4%)
2	B	1.04	1/2615 (0.0%)	1.63	36/3575 (1.0%)
All	All	1.04	1/2986 (0.0%)	1.64	43/4069 (1.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	17	GLU	CD-OE2	5.19	1.31	1.25

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	233	LEU	CB-CG-CD2	-10.76	92.71	111.00
2	B	279	SER	CB-CA-C	-9.46	92.12	110.10
1	P	14	ARG	NE-CZ-NH2	-8.99	115.81	120.30
2	B	38	LEU	CB-CG-CD2	-8.15	97.15	111.00
2	B	189	LEU	CB-CG-CD2	7.65	124.00	111.00
2	B	310	ARG	NE-CZ-NH2	7.17	123.89	120.30
2	B	322	ARG	NE-CZ-NH2	7.00	123.80	120.30
2	B	199	LEU	CB-CG-CD2	-6.83	99.40	111.00
2	B	28	LEU	CB-CG-CD1	-6.79	99.45	111.00
2	B	146	LEU	CB-CG-CD1	-6.29	100.30	111.00
2	B	17	GLU	OE1-CD-OE2	6.27	130.82	123.30
2	B	248	LEU	CB-CG-CD1	-6.22	100.42	111.00
1	P	11	LYS	CB-CA-C	-6.13	98.14	110.40
2	B	234	LEU	CB-CG-CD1	-6.09	100.65	111.00
2	B	148	SER	CB-CA-C	-6.02	98.66	110.10
2	B	73	LEU	CB-CG-CD1	-6.01	100.78	111.00
2	B	222	LEU	CB-CG-CD1	-6.01	100.79	111.00
1	P	19	GLU	CB-CA-C	-5.93	98.54	110.40
2	B	196	GLU	OE1-CD-OE2	-5.92	116.19	123.30
2	B	124	ALA	C-N-CA	-5.92	106.91	121.70
2	B	156	SER	CA-CB-OG	-5.88	95.32	111.20

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	7	MET	CG-SD-CE	-5.84	90.86	100.20
2	B	172	ASP	CB-CG-OD1	5.77	123.49	118.30
2	B	90	THR	CA-CB-CG2	-5.75	104.35	112.40
2	B	249	VAL	CB-CA-C	-5.71	100.55	111.40
1	P	28	ARG	NE-CZ-NH2	-5.64	117.48	120.30
2	B	223	LEU	CB-CA-C	-5.61	99.53	110.20
1	P	43	LEU	CB-CG-CD2	5.56	120.45	111.00
2	B	10	MET	CG-SD-CE	-5.54	91.33	100.20
1	P	38	TYR	N-CA-CB	-5.49	100.72	110.60
1	P	17	MET	CG-SD-CE	-5.47	91.45	100.20
2	B	248	LEU	CB-CA-C	-5.47	99.81	110.20
2	B	162	SER	C-N-CA	-5.38	110.99	122.30
2	B	114	TYR	CD1-CE1-CZ	-5.24	115.09	119.80
2	B	275	SER	CB-CA-C	-5.23	100.17	110.10
2	B	134	THR	N-CA-C	5.17	124.95	111.00
2	B	126	PRO	N-CA-C	5.14	125.47	112.10
2	B	196	GLU	N-CA-CB	-5.14	101.35	110.60
2	B	60	GLU	OE1-CD-OE2	-5.13	117.14	123.30
2	B	95	GLN	CA-CB-CG	-5.12	102.13	113.40
2	B	92	GLN	CG-CD-NE2	5.05	128.83	116.70
2	B	36	SER	CA-CB-OG	-5.04	97.60	111.20
2	B	51	THR	C-N-CA	-5.01	109.19	121.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	362	0	387	19	0
2	B	2517	0	2339	53	0
3	B	238	0	0	3	0
3	P	12	0	0	0	0
All	All	3129	0	2726	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:130:VAL:HG13	2:B:142:GLN:HE21	1.24	1.00
2:B:135:THR:HG22	2:B:137:MET:H	1.28	0.97
1:P:28:ARG:HD3	2:B:244:TYR:CE1	2.03	0.92
2:B:30:LEU:HB3	2:B:120[B]:ILE:HD13	1.54	0.88
1:P:9:LYS:HD3	2:B:159:GLN:HG3	1.55	0.88
2:B:135:THR:HG22	2:B:137:MET:N	1.88	0.87
2:B:130:VAL:HG13	2:B:142:GLN:NE2	1.97	0.78
2:B:92:GLN:O	2:B:93:SER:HB3	1.85	0.76
1:P:30:HIS:HB3	2:B:7:MET:HE2	1.71	0.73
1:P:31:LYS:HD3	1:P:32:TYR:N	2.04	0.73
2:B:207:TRP:CH2	2:B:229:TYR:HB3	2.25	0.72
1:P:9:LYS:CD	2:B:159:GLN:HG3	2.20	0.71
2:B:82:GLY:HA2	2:B:105:GLU:HG3	1.73	0.70
1:P:28:ARG:HD3	2:B:244:TYR:HE1	1.57	0.68
2:B:255:GLN:H	2:B:255:GLN:CD	1.97	0.66
2:B:135:THR:CG2	2:B:137:MET:H	2.08	0.65
1:P:32:TYR:HB2	2:B:7:MET:HE3	1.77	0.64
2:B:73:LEU:HD12	2:B:73:LEU:C	2.23	0.58
1:P:29:THR:HG22	1:P:30:HIS:CE1	2.39	0.57
1:P:28:ARG:HD3	2:B:244:TYR:CD1	2.41	0.56
2:B:92:GLN:O	2:B:93:SER:CB	2.52	0.55
1:P:32:TYR:HA	1:P:36:TRP:CZ3	2.42	0.55
2:B:243:GLU:HG3	2:B:244:TYR:CZ	2.41	0.55
2:B:248:LEU:HD22	2:B:283:TYR:CD2	2.43	0.54
2:B:197:GLU:HB3	2:B:263:ILE:HD12	1.90	0.53
2:B:207:TRP:HB3	2:B:229:TYR:CE2	2.44	0.52
2:B:30:LEU:HB3	2:B:120[B]:ILE:CD1	2.34	0.52
2:B:19:SER:HB2	2:B:90:THR:HB	1.92	0.52
2:B:207:TRP:CZ3	2:B:229:TYR:HB3	2.45	0.52
1:P:32:TYR:HB2	2:B:7:MET:CE	2.40	0.51
1:P:40:PHE:CD1	1:P:40:PHE:N	2.78	0.51
2:B:156:SER:O	2:B:159:GLN:OE1	2.27	0.51
2:B:255:GLN:OE1	2:B:255:GLN:N	2.38	0.51
2:B:49:ALA:O	2:B:53:HIS:HD2	1.95	0.50
2:B:296:GLN:NE2	3:B:602:HOH:O	2.43	0.50
2:B:129:SER:HB2	2:B:131:ASP:OD1	2.12	0.49
2:B:4:TYR:CD2	2:B:39:TRP:HZ2	2.30	0.49
2:B:4:TYR:CE2	2:B:39:TRP:HZ2	2.31	0.49

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:25:GLU:O	1:P:29:THR:HB	2.13	0.48
2:B:91:VAL:O	2:B:92:GLN:HB2	2.13	0.48
2:B:243:GLU:HG3	2:B:244:TYR:CE2	2.49	0.47
1:P:20:LYS:NZ	2:B:114:TYR:HE1	2.13	0.47
2:B:296:GLN:HG2	3:B:480:HOH:O	2.14	0.47
1:P:40:PHE:H	1:P:40:PHE:HD1	1.59	0.47
1:P:16:THR:O	1:P:20:LYS:HG2	2.16	0.46
2:B:47:SER:O	2:B:51:THR:HG23	2.16	0.46
1:P:27:LEU:HD23	1:P:27:LEU:HA	1.41	0.44
1:P:35:ALA:HB3	2:B:293:LEU:HD12	2.00	0.43
2:B:134:THR:O	2:B:134:THR:HG22	2.18	0.43
2:B:248:LEU:HD23	2:B:285:THR:HG22	2.01	0.43
2:B:49:ALA:HB3	2:B:107[B]:GLU:OE1	2.19	0.42
2:B:126:PRO:HD2	2:B:188:GLU:O	2.19	0.42
2:B:280:ASN:O	2:B:281:ASN:HB2	2.19	0.42
2:B:3:THR:HG22	3:B:598:HOH:O	2.20	0.42
1:P:32:TYR:CE1	2:B:5:GLU:HG2	2.56	0.41
2:B:282:GLY:O	2:B:283:TYR:HB3	2.20	0.41
2:B:80:LEU:HD12	2:B:106:ASN:OD1	2.21	0.41
2:B:248:LEU:CD2	2:B:283:TYR:CD2	3.03	0.41
2:B:5:GLU:HA	2:B:6:PRO:HD2	1.91	0.41
2:B:30:LEU:HD23	2:B:120[B]:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	41/43 (95%)	41 (100%)	0	0	100	100
2	B	333/329 (101%)	323 (97%)	9 (3%)	1 (0%)	41	21
All	All	374/372 (100%)	364 (97%)	9 (2%)	1 (0%)	41	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	76	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	P	38/38 (100%)	34 (90%)	4 (10%)	<b>7</b> <b>0</b>
2	B	280/274 (102%)	277 (99%)	3 (1%)	73 56
All	All	318/312 (102%)	311 (98%)	7 (2%)	52 25

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	P	20	LYS
1	P	28	ARG
1	P	31	LYS
1	P	40	PHE
2	B	93	SER
2	B	187	GLN
2	B	240	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	53	HIS
2	B	142	GLN
2	B	240	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.