



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 13, 2024 – 11:50 pm GMT

PDB ID : 6HTV  
Title : Crystal structure of *Leuconostoc citreum* NRRL B-1299 N-terminally truncated dextransucrase DSR-M in complex with isomaltotetraose  
Authors : Claverie, M.; Cioci, G.; Remaud-Simeon, M.; Moulis, C.; Lippens, G.  
Deposited on : 2018-10-04  
Resolution : 3.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1002 (4.14-3.66)
Clashscore	141614	1004 (4.12-3.68)
Ramachandran outliers	138981	1021 (4.14-3.66)
Sidechain outliers	138945	1014 (4.14-3.66)
RSRZ outliers	127900	1275 (4.20-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1290	 76% 6% 18%
2	B	4	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	B	1	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8407 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alternansucrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1064	8361	5229	1414	1701	17	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	4	45	24	21	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	185.31Å 185.31Å 150.77Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.96 – 3.90 47.26 – 3.90	Depositor EDS
% Data completeness (in resolution range)	96.4 (47.96-3.90) 96.5 (47.26-3.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.16 (at 3.88Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.210 , 0.232 0.210 , 0.232	Depositor DCC
$R_{free}$ test set	1319 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	110.1	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 67.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8407	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	117.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.56	0/8542	0.69	0/11613

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8361	0	7915	62	0
2	B	45	0	39	0	0
3	A	1	0	0	0	0
All	All	8407	0	7954	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:853:TYR:CE2	1:A:870:ASP:OD2	2.36	0.78

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:433:SER:HB3	1:A:435:ASP:OD1	1.88	0.72
1:A:725:ASN:OD1	1:A:729:ASN:ND2	2.21	0.72
1:A:809:TYR:CE2	1:A:818:ALA:HB2	2.27	0.69
1:A:819:LYS:HE3	1:A:1120:VAL:HB	1.75	0.67
1:A:853:TYR:HE2	1:A:870:ASP:OD2	1.78	0.66
1:A:798:ALA:CB	1:A:816:LEU:HD12	2.26	0.65
1:A:1120:VAL:CG2	1:A:1123:VAL:HG22	2.27	0.65
1:A:604:PRO:HB3	1:A:724:TYR:CD2	2.31	0.64
1:A:605:SER:OG	1:A:720:ASN:ND2	2.20	0.62
1:A:725:ASN:O	1:A:729:ASN:HB3	2.00	0.61
1:A:388:ASN:OD1	1:A:402:LEU:HD23	2.02	0.60
1:A:819:LYS:HE3	1:A:1120:VAL:CB	2.34	0.57
1:A:853:TYR:CZ	1:A:870:ASP:OD2	2.58	0.56
1:A:433:SER:CB	1:A:435:ASP:OD1	2.54	0.55
1:A:809:TYR:CZ	1:A:818:ALA:HB2	2.40	0.55
1:A:369:VAL:CG1	1:A:376:LYS:HE3	2.36	0.54
1:A:798:ALA:HB1	1:A:816:LEU:HD12	1.89	0.54
1:A:434:TYR:CE1	1:A:1313:MET:HB3	2.43	0.54
1:A:605:SER:HG	1:A:720:ASN:HD22	1.49	0.53
1:A:853:TYR:OH	1:A:870:ASP:OD2	2.23	0.53
1:A:498:LEU:HD23	1:A:523:ALA:HB1	1.91	0.53
1:A:786:ILE:HG22	1:A:787:ARG:HG2	1.90	0.52
1:A:788:ALA:HB3	1:A:791:SER:HB3	1.90	0.52
1:A:809:TYR:HD1	1:A:814:LYS:HG2	1.76	0.51
1:A:502:LYS:HD3	1:A:504:TYR:CE1	2.46	0.50
1:A:870:ASP:O	1:A:883:THR:HG21	2.12	0.50
1:A:504:TYR:CZ	1:A:516:ALA:HB2	2.47	0.49
1:A:808:LEU:HG	1:A:809:TYR:CE2	2.48	0.48
1:A:975:THR:HG22	1:A:980:ILE:HG12	1.95	0.48
1:A:798:ALA:HB3	1:A:816:LEU:HD12	1.94	0.48
1:A:1352:TYR:CD1	1:A:1372:VAL:HG21	2.48	0.47
1:A:502:LYS:HD3	1:A:504:TYR:CZ	2.49	0.47
1:A:808:LEU:CD2	1:A:809:TYR:CE2	2.97	0.47
1:A:939:ILE:HD11	1:A:1024:VAL:CG1	2.44	0.47
1:A:604:PRO:HB3	1:A:724:TYR:CG	2.50	0.47
1:A:819:LYS:NZ	1:A:1120:VAL:HA	2.30	0.47
1:A:787:ARG:NH1	1:A:870:ASP:OD1	2.48	0.46
1:A:1132:ARG:HD3	1:A:1227:GLN:OE1	2.15	0.46
1:A:730:GLY:O	1:A:1049:SER:HA	2.15	0.46
1:A:1414:TYR:HA	1:A:1424:TYR:HA	1.98	0.45
1:A:787:ARG:HD2	1:A:792:GLU:HB2	1.99	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1380:TYR:HB2	1:A:1403:PHE:CZ	2.53	0.44
1:A:786:ILE:HD13	1:A:852:ALA:HB1	2.00	0.44
1:A:434:TYR:CE1	1:A:1313:MET:CB	3.00	0.44
1:A:1120:VAL:HG22	1:A:1123:VAL:HG22	1.99	0.43
1:A:731:ASN:HD22	1:A:1049:SER:HB3	1.83	0.43
1:A:786:ILE:C	1:A:787:ARG:HG2	2.39	0.43
1:A:376:LYS:NZ	1:A:398:ILE:HG13	2.33	0.43
1:A:580:PHE:CE1	1:A:1210:ILE:HD11	2.55	0.42
1:A:819:LYS:NZ	1:A:1114:GLY:CA	2.83	0.42
1:A:377:TYR:CE1	1:A:385:ARG:NH2	2.88	0.42
1:A:1120:VAL:HG22	1:A:1123:VAL:CG2	2.50	0.41
1:A:1117:LEU:O	1:A:1120:VAL:HG13	2.19	0.41
1:A:435:ASP:OD1	1:A:435:ASP:N	2.53	0.41
1:A:809:TYR:CD1	1:A:814:LYS:HE2	2.56	0.41
1:A:867:TYR:CE2	1:A:869:GLY:HA3	2.55	0.41
1:A:786:ILE:CD1	1:A:852:ALA:HB1	2.50	0.41
1:A:868:TYR:CE1	1:A:878:TYR:CE1	3.09	0.41
1:A:369:VAL:HG13	1:A:376:LYS:HE3	2.03	0.41
1:A:603:THR:O	1:A:607:GLN:N	2.54	0.40
1:A:870:ASP:O	1:A:883:THR:CG2	2.70	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1062/1290 (82%)	1030 (97%)	32 (3%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	909/1095 (83%)	890 (98%)	19 (2%)	53 73

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	382	GLN
1	A	397	PHE
1	A	434	TYR
1	A	572	THR
1	A	598	ARG
1	A	675	ARG
1	A	704	ASP
1	A	754	SER
1	A	803	ASP
1	A	874	ASP
1	A	897	THR
1	A	908	VAL
1	A	1053	SER
1	A	1061	LEU
1	A	1109	ARG
1	A	1118	ASP
1	A	1132	ARG
1	A	1178	ASP
1	A	1193	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	742	ASN
1	A	1145	ASN
1	A	1212	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	B	1	2	12,12,12	0.56	0	17,17,17	1.87	3 (17%)
2	GLC	B	2	2	11,11,12	0.76	0	15,15,17	1.73	4 (26%)
2	GLC	B	3	2	11,11,12	0.65	0	15,15,17	0.98	1 (6%)
2	GLC	B	4	2	11,11,12	0.59	0	15,15,17	1.38	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	2/2/22/22	0/1/1/1
2	GLC	B	2	2	-	1/2/19/22	0/1/1/1
2	GLC	B	3	2	-	1/2/19/22	0/1/1/1
2	GLC	B	4	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	O1-C1-O5	5.52	126.96	110.38
2	B	4	GLC	O5-C5-C6	4.02	113.51	107.20
2	B	2	GLC	O5-C1-C2	-3.51	105.35	110.77
2	B	1	GLC	C1-C2-C3	-3.34	103.38	110.31
2	B	2	GLC	O2-C2-C1	3.32	115.95	109.15
2	B	2	GLC	O5-C5-C6	2.72	111.47	107.20
2	B	1	GLC	C3-C4-C5	2.50	114.70	110.24
2	B	2	GLC	C6-C5-C4	-2.31	107.59	113.00
2	B	3	GLC	O5-C5-C6	2.29	110.80	107.20

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	GLC	C4-C5-C6-O6
2	B	1	GLC	O5-C5-C6-O6
2	B	4	GLC	O5-C5-C6-O6
2	B	4	GLC	C4-C5-C6-O6
2	B	3	GLC	O5-C5-C6-O6
2	B	2	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1064/1290 (82%)	-0.32	4 (0%) <b>92</b> <b>87</b>	76, 112, 161, 194	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	376	LYS	2.5
1	A	389	TYR	2.2
1	A	378	TYR	2.2
1	A	1428	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	B	1	12/12	<b>0.69</b>	<b>0.41</b>	136,155,171,172	0
2	GLC	B	3	11/12	0.82	0.32	121,161,174,184	0
2	GLC	B	2	11/12	0.86	0.33	135,149,162,169	0
2	GLC	B	4	11/12	0.87	0.33	124,156,179,182	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	A	1505	1/1	0.99	0.24	66,66,66,66	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.