

Full wwPDB X-ray Structure Validation Report (i)

Nov 7, 2023 – 06:51 PM JST

PDB ID : 8HX9

Title : Crystal structure of 4-amino-4-deoxychorismate synthase from Streptomyces

venezuelae with chorismate

Authors : Nakamichi, Y.; Watanabe, M.

Deposited on : 2023-01-04

Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	702	83%	7%	10%
1	В	702	85%	9%	5%



2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 10154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 4-amino-4-deoxychorismate synthase.

\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	632	Total	С	Ν	О	S	0	1	0
1	Λ	052	4900	3075	880	927	18	0	1	
1	B	665	Total	С	N	О	S	0	0	0
1	D	000	5128	3211	927	973	17	0		U

There are 32 discrepancies between the modelled and reference sequences:

Chain A	Residue	Modelled	A at 1	O 1	
A	4 -		Actual	Comment	Reference
	-15	MET	-	initiating methionine	UNP Q6L8Q5
A	-14	ASN	-	expression tag	UNP Q6L8Q5
A	-13	HIS	-	expression tag	UNP Q6L8Q5
A	-12	LYS	-	expression tag	UNP Q6L8Q5
A	-11	VAL	-	expression tag	UNP Q6L8Q5
A	-10	HIS	-	expression tag	UNP Q6L8Q5
A	-9	HIS	-	expression tag	UNP Q6L8Q5
A	-8	HIS	-	expression tag	UNP Q6L8Q5
A	-7	HIS	-	expression tag	UNP Q6L8Q5
A	-6	HIS	-	expression tag	UNP Q6L8Q5
A	-5	HIS	-	expression tag	UNP Q6L8Q5
A	-4	ILE	-	expression tag	UNP Q6L8Q5
A	-3	GLU	-	expression tag	UNP Q6L8Q5
A	-2	GLY	-	expression tag	UNP Q6L8Q5
A	-1	ARG	-	expression tag	UNP Q6L8Q5
A	0	HIS	-	expression tag	UNP Q6L8Q5
В	-15	MET	-	initiating methionine	UNP Q6L8Q5
В	-14	ASN	-	expression tag	UNP Q6L8Q5
В	-13	HIS	-	expression tag	UNP Q6L8Q5
В	-12	LYS	-	expression tag	UNP Q6L8Q5
В	-11	VAL	_	expression tag	UNP Q6L8Q5
В	-10	HIS	-	expression tag	UNP Q6L8Q5
В	-9	HIS	-	expression tag	UNP Q6L8Q5
В	-8	HIS	-	expression tag	UNP Q6L8Q5
В	-7	HIS	-	expression tag	UNP Q6L8Q5


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	-6	HIS	-	expression tag	UNP Q6L8Q5
В	-5	HIS	-	expression tag	UNP Q6L8Q5
В	-4	ILE	-	expression tag	UNP Q6L8Q5
В	-3	GLU	-	expression tag	UNP Q6L8Q5
В	-2	GLY	-	expression tag	UNP Q6L8Q5
В	-1	ARG	-	expression tag	UNP Q6L8Q5
В	0	HIS	-	expression tag	UNP Q6L8Q5


• Molecule 2 is TRYPTOPHAN (three-letter code: TRP) (formula: $C_{11}H_{12}N_2O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 15	_		_	0	0
2	В	1	Total 15	_	N 2	_	0	0


• Molecule 3 is (3R,4R)-3-[(1-carboxyethenyl)oxy]-4-hydroxycyclohexa-1,5-diene-1-carbo xylic acid (three-letter code: ISJ) (formula: $C_{10}H_{10}O_6$) (labeled as "Ligand of Interest" by depositor).





N	Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
	3	A	1	Total 16	C 10	O 6	0	0

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0

• Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total Mg 2 2	0	0
5	В	1	Total Mg 1 1	0	0

 \bullet Molecule 6 is FORMIC ACID (three-letter code: FMT) (formula: $\mathrm{CH_2O_2}).$



N	Λ ol	Chain	Residues	Atoms		ZeroOcc	AltConf
	6	В	1	Total 3	C O 1 2	0	0

• Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	46	Total O 46 46	0	0
7	В	22	Total O 22 22	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 4-amino-4-deoxychorismate synthase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	99.29Å 170.78Å 162.08Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.47 - 2.03	Depositor
Resolution (A)	47.47 - 2.03	EDS
% Data completeness	99.9 (47.47-2.03)	Depositor
(in resolution range)	99.9 (47.47-2.03)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.57 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
D D.	0.218 , 0.279	Depositor
R, R_{free}	0.232 , 0.292	DCC
R_{free} test set	4446 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	41.9	Xtriage
Anisotropy	0.768	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 46.8	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	0.000 for 1/2 +h-1/2 +k,-3/2 +h-1/2 +k,-l	Xtriage
Estimated twinning fraction	0.017 for 1/2 *h + 1/2 *k, 3/2 *h - 1/2 *k, -1	Amage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10154	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, ISJ, MG, OCS, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.27	0/4997	0.48	0/6781	
1	В	0.27	0/5234	0.48	0/7108	
All	All	0.27	0/10231	0.48	0/13889	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4900	0	4797	27	0
1	В	5128	0	5013	31	0
2	A	15	0	9	0	0
2	В	15	0	9	0	0
3	A	16	0	8	0	0
4	A	6	0	8	0	0
5	A	2	0	0	0	0
5	В	1	0	0	0	0
6	В	3	0	1	0	0
7	A	46	0	0	0	0
7	В	22	0	0	0	0
All	All	10154	0	9845	58	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:309:LEU:HD12	1:A:359:LEU:HD13	1.74	0.70
1:A:398:ILE:HB	1:A:399:PRO:HD3	1.80	0.64
1:B:320:LYS:NZ	1:B:606:ASP:OD1	2.32	0.60
1:A:398:ILE:HD11	1:A:515:LEU:HD11	1.87	0.56
1:A:500:LYS:HD2	1:A:652:ALA:HB2	1.88	0.56
1:A:502:THR:HG23	1:A:563:VAL:HG21	1.87	0.55
1:B:428:GLU:HG2	1:B:604:ILE:HD13	1.90	0.53
1:A:535:LEU:C	1:A:535:LEU:HD23	2.27	0.53
1:B:91:PHE:HB3	1:B:141:LEU:HD21	1.93	0.51
1:A:535:LEU:HD22	1:A:591:GLY:HA3	1.93	0.50
1:B:452:LEU:HB3	1:B:453:PRO:HD3	1.95	0.49
1:A:502:THR:HG23	1:A:563:VAL:CG2	2.43	0.49
1:B:264:ALA:HB2	1:B:330:ARG:HD3	1.94	0.49
1:B:259:LEU:C	1:B:259:LEU:HD23	2.33	0.48
1:A:502:THR:CG2	1:A:563:VAL:HG21	2.44	0.47
1:B:601:THR:HA	1:B:604:ILE:HD12	1.96	0.47
1:A:314:TYR:CE2	1:A:337:ALA:HB3	2.50	0.47
1:B:274:ASP:HB2	1:B:276:THR:HG22	1.97	0.46
1:A:270:VAL:O	1:A:277:THR:HA	2.14	0.46
1:B:207:VAL:HG22	1:B:357:LEU:HD22	1.96	0.46
1:B:636:ARG:NH2	1:B:669:LYS:HD3	2.30	0.46
1:B:530:LEU:C	1:B:530:LEU:HD23	2.36	0.46
1:B:213:LEU:HD22	1:B:352:GLY:HA2	1.98	0.45
1:A:257:GLU:O	1:A:339:LEU:HD12	2.16	0.45
1:A:271:ARG:HA	1:A:277:THR:HG22	1.99	0.45
1:A:259:LEU:HD23	1:A:287:TYR:HE2	1.82	0.45
1:B:259:LEU:HD21	1:B:284:PHE:HE1	1.82	0.45
1:B:356:LEU:C	1:B:357:LEU:HD23	2.36	0.45
1:A:259:LEU:HD12	1:A:259:LEU:N	2.31	0.45
1:A:265:ASP:HB2	1:A:267:VAL:HG22	1.97	0.45
1:B:475:LEU:HD23	1:B:475:LEU:N	2.32	0.44
1:B:511:GLU:O	1:B:515:LEU:HD12	2.18	0.44
1:B:8:ASN:C	1:B:8:ASN:HD22	2.20	0.44
1:A:551:VAL:HG11	1:A:554:LEU:HD13	2.00	0.44
1:B:407:PRO:HB2	1:B:408:LEU:HD22	2.00	0.44
1:A:475:LEU:HD12	1:A:639:VAL:HG13	2.00	0.43
1:A:561:ALA:HB3	1:A:562:PRO:HD3	1.99	0.43

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
		$\operatorname{distance}\ (ext{Å})$	overlap (Å)
1:B:308:ASN:O	1:B:309:LEU:HB2	2.19	0.43
1:B:210:VAL:HG12	1:B:212:VAL:HG22	2.00	0.43
1:B:229:THR:HA	1:B:471:GLU:O	2.19	0.43
1:B:76:PRO:HG2	1:B:188:ALA:HA	2.01	0.43
1:A:405:PHE:HB3	1:A:443:VAL:HG11	2.01	0.43
1:B:224:LEU:HD23	1:B:230:THR:HB	2.01	0.43
1:B:588:PHE:CD1	1:B:589:PRO:HA	2.53	0.43
1:A:309:LEU:CD1	1:A:359:LEU:HD13	2.45	0.42
1:B:69:ALA:O	1:B:73:SER:HB3	2.19	0.42
1:B:561:ALA:HB3	1:B:562:PRO:HD3	2.01	0.42
1:A:499:ILE:HG12	1:A:567:VAL:HG22	2.02	0.41
1:A:565:GLN:HE22	1:A:636[A]:ARG:HH22	1.66	0.41
1:B:259:LEU:HA	1:B:269:SER:O	2.20	0.41
1:A:270:VAL:CG1	1:A:278:THR:HB	2.49	0.41
1:B:320:LYS:HB2	1:B:602:MET:HE3	2.02	0.41
1:A:500:LYS:HA	1:A:500:LYS:HD3	1.96	0.41
1:A:32:PRO:HG2	1:A:35:ALA:HB2	2.02	0.40
1:B:4:LEU:C	1:B:4:LEU:HD13	2.42	0.40
1:B:192:HIS:O	1:B:195:ARG:HB3	2.21	0.40
1:B:76:PRO:HA	1:B:161:PRO:O	2.22	0.40
1:A:270:VAL:HG13	1:A:278:THR:HB	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Favoured Allowed		Perce	ntiles
1	A	620/702 (88%)	595 (96%)	25 (4%)	0	100	100
1	В	660/702 (94%)	638 (97%)	22 (3%)	0	100	100
All	All	1280/1404 (91%)	1233 (96%)	47 (4%)	0	100	100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Rotameric Outliers	
1	A	512/560 (91%)	501 (98%)	11 (2%)	53 55
1	В	534/560 (95%)	518 (97%)	16 (3%)	41 40
All	All	1046/1120 (93%)	1019 (97%)	27 (3%)	46 46

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ASP
1	A	39	ARG
1	A	105	ARG
1	A	118	ARG
1	A	156	ARG
1	A	288	LEU
1	A	359	LEU
1	A	373	ARG
1	A	410	ARG
1	A	421	ARG
1	A	522	ARG
1	В	8	ASN
1	В	9	TYR
1	В	58	ASP
1	В	102	MET
1	В	195	ARG
1	В	196	ARG
1	В	197	HIS
1	В	274	ASP
1	В	410	ARG
1	В	420	LYS
1	В	421	ARG
1	В	430	ARG
1	В	500	LYS
1	В	553	ARG

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	В	640	LEU
1	В	671	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	191	HIS
1	A	286	ASN
1	A	550	HIS
1	В	8	ASN
1	В	84	HIS
1	В	89	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Ch		pe Chain Res Lin		Link	В	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
1	OCS	A	81	1	7,8,9	1.52	1 (14%)	6,11,13	1.75	2 (33%)	
1	OCS	В	81	1	7,8,9	1.42	1 (14%)	6,11,13	1.69	2 (33%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	A	81	1	-	4/4/7/9	-
1	OCS	В	81	1	-	2/4/7/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	A	81	OCS	OD1-SG	3.71	1.56	1.45
1	В	81	OCS	OD3-SG	3.38	1.55	1.45

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	81	OCS	OD2-SG-OD3	2.95	118.47	111.27
1	В	81	OCS	OD3-SG-CB	-2.77	103.64	106.94
1	В	81	OCS	OD2-SG-OD1	2.75	117.99	111.27
1	A	81	OCS	OD1-SG-CB	-2.68	103.75	106.94

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	81	OCS	N-CA-CB-SG
1	A	81	OCS	CA-CB-SG-OD1
1	A	81	OCS	CA-CB-SG-OD2
1	A	81	OCS	CA-CB-SG-OD3
1	В	81	OCS	N-CA-CB-SG
1	В	81	OCS	CA-CB-SG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain	Res	es Link	Bond lengths			Bond angles		
MIOI	Type		nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ISJ	A	702	5	16,16,16	3.67	5 (31%)	14,22,22	1.42	2 (14%)
4	GOL	A	703	-	5,5,5	0.10	0	5,5,5	0.31	0
2	TRP	A	701	-	14,16,16	0.76	0	16,22,22	0.90	1 (6%)
6	FMT	В	702	-	2,2,2	0.95	0	1,1,1	0.25	0
2	TRP	В	701	-	14,16,16	0.75	0	16,22,22	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRP	A	701	-	-	1/7/8/8	0/2/2/2
2	TRP	В	701	-	-	0/7/8/8	0/2/2/2
3	ISJ	A	702	5	-	0/10/25/25	0/1/1/1
4	GOL	A	703	-	-	0/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(A)
3	A	702	ISJ	C12-C13	-13.38	1.36	1.49
3	A	702	ISJ	C2-C3	3.32	1.43	1.35
3	A	702	ISJ	O9-C7	-2.94	1.22	1.30
3	A	702	ISJ	O15-C13	-2.52	1.23	1.30
3	A	702	ISJ	C4-C3	-2.01	1.39	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	702	ISJ	O15-C13-C12	2.74	118.59	113.91
3	A	702	ISJ	C6-C1-C2	2.40	113.26	110.67
2	A	701	TRP	CH2-CZ2-CE2	-2.01	117.19	120.08

There are no chirality outliers.



All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	TRP	O-C-CA-N

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$		$OWAB(Å^2)$	Q < 0.9
1	A	631/702 (89%)	0.47	40 (6%)	20 19	36, 59, 100, 146	0
1	В	664/702 (94%)	0.43	41 (6%)	20 19	39, 66, 96, 121	0
All	All	1295/1404 (92%)	0.45	81 (6%)	20 19	36, 63, 97, 146	0

All (81) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	TRP	10.7
1	A	39	ARG	6.1
1	В	132	LEU	6.1
1	A	40	LEU	5.6
1	A	45	PHE	5.4
1	В	199	ALA	5.4
1	A	226	GLY	5.3
1	A	70	ILE	4.5
1	A	68	ARG	4.3
1	A	398	ILE	4.2
1	A	71	THR	4.1
1	В	362	ARG	4.1
1	A	74	GLY	4.0
1	A	160	LYS	3.8
1	A	43	GLU	3.8
1	A	105	ARG	3.8
1	В	197	HIS	3.7
1	A	267	VAL	3.6
1	В	87	ILE	3.5
1	A	362	ARG	3.4
1	A	141	LEU	3.4
1	A	427	LYS	3.3
1	В	45	PHE	3.3
1	В	198	GLY	3.2

Continued on next page...



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	В	84	HIS	3.2
1	В	77	VAL	3.2
1	В	281	ARG	3.2
1	В	85	GLN	3.1
1	В	301	PRO	3.1
1	В	34	ASP	3.1
1	В	363	GLY	3.0
1	В	65	ILE	2.9
1	A	44	ASP	2.9
1	A	394	MET	2.9
1	В	506	GLY	2.9
1	A	75	LEU	2.9
1	В	68	ARG	2.9
1	В	39	ARG	2.9
1	В	204	GLU	2.8
1	A	69	ALA	2.8
1	В	405	PHE	2.8
1	В	196	ARG	2.7
1	В	134	ALA	2.7
1	A	155	LEU	2.7
1	A	29	VAL	2.7
1	A	162	LEU	2.7
1	В	72	ASP	2.6
1	В	139	ASP	2.6
1	В	88	ALA	2.6
1	В	99	PRO	2.6
1	В	507	GLY	2.6
1	A	195	ARG	2.5
1	В	615	VAL	2.4
1	A	280	THR	2.4
1	В	273	SER	2.4
1	В	430	ARG	2.4
1	В	259	LEU	2.4
1	A	146	TRP	2.4
1	В	43	GLU	2.4
1	В	-1	ARG	2.4
1	A	8	ASN	2.3
1	A	281	ARG	2.3
1	A	259	LEU	2.3
1	A	296	ARG	2.3
1	В	37	TRP	2.3
1	A	391	THR	2.3

| 391 | THR | 2.3 | Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	286	ASN	2.3
1	A	395	VAL	2.2
1	В	67	ARG	2.2
1	В	279	ARG	2.2
1	A	6	ILE	2.2
1	A	268	VAL	2.2
1	A	192	HIS	2.2
1	В	58	ASP	2.2
1	В	141	LEU	2.1
1	A	156	ARG	2.1
1	В	42	VAL	2.1
1	A	103	HIS	2.1
1	В	162	LEU	2.0
1	A	510	GLU	2.0
1	В	364	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	OCS	A	81	9/10	0.96	0.10	73,77,83,87	0
1	OCS	В	81	9/10	0.96	0.14	53,54,59,61	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	GOL	A	703	6/6	0.71	0.20	75,77,79,80	0

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
5	MG	A	705	1/1	0.88	0.27	58,58,58,58	0
5	MG	A	704	1/1	0.89	0.05	49,49,49,49	0
3	ISJ	A	702	16/16	0.92	0.18	51,57,61,63	0
5	MG	В	703	1/1	0.92	0.08	46,46,46,46	0
6	FMT	В	702	3/3	0.92	0.15	50,50,51,51	0
2	TRP	В	701	15/15	0.93	0.12	37,41,47,47	0
2	TRP	A	701	15/15	0.97	0.12	36,38,38,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





Electron density around MG A 705: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around MG A 704:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)















Electron density around MG B 703: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around TRP B 701:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around TRP A 701:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

