



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 01:37 PM EDT

PDB ID : 3HYL
Title : Crystal Structure of Transketolase from Bacillus anthracis
Authors : Maltseva, N.; Kim, Y.; Kwon, K.; Joachimiak, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2009-06-22
Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	690	
1	B	690	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FMT	A	669	-	-	X	-
3	FMT	A	673	-	-	X	-
3	FMT	B	668	-	-	X	-
3	FMT	B	674	-	-	X	-
4	PEG	A	677	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 10868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	663	5087	3198	862	1005	1	21	0	2	0
1	B	663	5096	3205	863	1006	1	21	0	3	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MSE	-	EXPRESSION TAG	UNP C3P4P9
A	-22	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-21	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-20	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-19	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-18	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-17	HIS	-	EXPRESSION TAG	UNP C3P4P9
A	-16	SER	-	EXPRESSION TAG	UNP C3P4P9
A	-15	SER	-	EXPRESSION TAG	UNP C3P4P9
A	-14	GLY	-	EXPRESSION TAG	UNP C3P4P9
A	-13	VAL	-	EXPRESSION TAG	UNP C3P4P9
A	-12	ASP	-	EXPRESSION TAG	UNP C3P4P9
A	-11	LEU	-	EXPRESSION TAG	UNP C3P4P9
A	-10	GLY	-	EXPRESSION TAG	UNP C3P4P9
A	-9	THR	-	EXPRESSION TAG	UNP C3P4P9
A	-8	GLU	-	EXPRESSION TAG	UNP C3P4P9
A	-7	ASN	-	EXPRESSION TAG	UNP C3P4P9
A	-6	LEU	-	EXPRESSION TAG	UNP C3P4P9
A	-5	TYR	-	EXPRESSION TAG	UNP C3P4P9
A	-4	PHE	-	EXPRESSION TAG	UNP C3P4P9
A	-3	GLN	-	EXPRESSION TAG	UNP C3P4P9
A	-2	SER	-	EXPRESSION TAG	UNP C3P4P9
A	-1	ASN	-	EXPRESSION TAG	UNP C3P4P9
A	0	ALA	-	EXPRESSION TAG	UNP C3P4P9
B	-23	MSE	-	EXPRESSION TAG	UNP C3P4P9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-21	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-20	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-19	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-18	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-17	HIS	-	EXPRESSION TAG	UNP C3P4P9
B	-16	SER	-	EXPRESSION TAG	UNP C3P4P9
B	-15	SER	-	EXPRESSION TAG	UNP C3P4P9
B	-14	GLY	-	EXPRESSION TAG	UNP C3P4P9
B	-13	VAL	-	EXPRESSION TAG	UNP C3P4P9
B	-12	ASP	-	EXPRESSION TAG	UNP C3P4P9
B	-11	LEU	-	EXPRESSION TAG	UNP C3P4P9
B	-10	GLY	-	EXPRESSION TAG	UNP C3P4P9
B	-9	THR	-	EXPRESSION TAG	UNP C3P4P9
B	-8	GLU	-	EXPRESSION TAG	UNP C3P4P9
B	-7	ASN	-	EXPRESSION TAG	UNP C3P4P9
B	-6	LEU	-	EXPRESSION TAG	UNP C3P4P9
B	-5	TYR	-	EXPRESSION TAG	UNP C3P4P9
B	-4	PHE	-	EXPRESSION TAG	UNP C3P4P9
B	-3	GLN	-	EXPRESSION TAG	UNP C3P4P9
B	-2	SER	-	EXPRESSION TAG	UNP C3P4P9
B	-1	ASN	-	EXPRESSION TAG	UNP C3P4P9
B	0	ALA	-	EXPRESSION TAG	UNP C3P4P9

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	B	1	3	1	2	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	7	4	3	0	0
4	A	1	7	4	3	0	0
4	A	1	7	4	3	0	0
4	B	1	7	4	3	0	0


- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	1	1	0	0
5	B	1	1	1	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total O S 5 4 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	343	Total O 343 343	0	0
8	B	261	Total O 261 261	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Transketolase



• Molecule 1: Transketolase





4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	138.80Å 70.97Å 145.79Å 90.00° 117.35° 90.00°	Depositor
Resolution (Å)	36.35 – 2.16 36.35 – 2.16	Depositor EDS
% Data completeness (in resolution range)	99.4 (36.35-2.16) 99.4 (36.35-2.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.36 (at 2.16Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.176 , 0.230 0.173 , 0.228	Depositor DCC
R_{free} test set	3423 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtrriage
Anisotropy	0.438	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.011 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10868	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, MG, SO4, FMT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/5174	0.77	4/6988 (0.1%)
1	B	0.64	0/5187	0.76	6/7005 (0.1%)
All	All	0.67	0/10361	0.76	10/13993 (0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	93	ARG	NE-CZ-NH2	-10.79	114.90	120.30
1	A	93	ARG	NE-CZ-NH1	10.20	125.40	120.30
1	B	93	ARG	NE-CZ-NH2	-7.87	116.37	120.30
1	A	665	MSE	CG-SE-CE	-6.16	85.36	98.90
1	B	31	MSE	CG-SE-CE	-5.65	86.46	98.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5087	0	4921	134	0
1	B	5096	0	4930	198	0
2	A	12	0	16	2	0
3	A	15	0	5	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	18	0	6	7	0
4	A	21	0	30	7	0
4	B	7	0	10	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
7	B	5	0	0	0	0
8	A	343	0	0	5	0
8	B	261	0	0	12	0
All	All	10868	0	9918	329	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 329 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:ASP:HB2	8:B:866:HOH:O	1.26	1.30
1:B:616:MSE:SE	8:B:741:HOH:O	2.12	1.16
1:B:465:ILE:HD11	1:B:616:MSE:HE2	1.26	1.12
1:B:616:MSE:HE3	1:B:653:PHE:CD2	1.85	1.12
1:A:130:MSE:SE	8:A:927:HOH:O	2.21	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	663/690 (96%)	637 (96%)	25 (4%)	1 (0%)	47 46
1	B	664/690 (96%)	634 (96%)	30 (4%)	0	100 100
All	All	1327/1380 (96%)	1271 (96%)	55 (4%)	1 (0%)	51 53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	258	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	528/527 (100%)	492 (93%)	36 (7%)	16 10
1	B	529/527 (100%)	483 (91%)	46 (9%)	10 6
All	All	1057/1054 (100%)	975 (92%)	82 (8%)	13 7

5 of 82 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	346	LEU
1	B	513	LYS
1	B	370	GLU
1	B	442	LEU
1	B	551	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	313	ASN
1	B	479	GLN
1	B	512	ASN
1	B	488	ASN
1	B	453	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 3 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	678	-	6,6,6	0.51	0	5,5,5	1.86	2 (40%)
3	FMT	A	671	-	2,2,2	0.62	0	1,1,1	0.04	0
3	FMT	B	672	-	2,2,2	0.57	0	1,1,1	0.09	0
3	FMT	B	675	-	2,2,2	0.64	0	1,1,1	0.13	0
4	PEG	B	669	-	6,6,6	0.59	0	5,5,5	2.30	3 (60%)
3	FMT	B	668	5	2,2,2	0.69	0	1,1,1	0.21	0
3	FMT	A	670	-	2,2,2	0.62	0	1,1,1	0.02	0
4	PEG	A	677	-	6,6,6	0.53	0	5,5,5	1.48	1 (20%)
3	FMT	B	674	-	2,2,2	0.74	0	1,1,1	0.42	0
2	GOL	A	667	-	5,5,5	0.40	0	5,5,5	0.57	0
7	SO4	B	671	-	4,4,4	0.26	0	6,6,6	0.18	0
3	FMT	A	672	-	2,2,2	0.61	0	1,1,1	0.05	0
3	FMT	B	673	-	2,2,2	0.61	0	1,1,1	0.03	0
3	FMT	A	669	-	2,2,2	0.57	0	1,1,1	0.40	0
4	PEG	A	674	-	6,6,6	0.53	0	5,5,5	1.65	1 (20%)
3	FMT	A	673	-	2,2,2	0.67	0	1,1,1	0.56	0
2	GOL	A	668	-	5,5,5	0.39	0	5,5,5	0.56	0
3	FMT	B	667	-	2,2,2	0.66	0	1,1,1	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	678	-	-	1/4/4/4	-
4	PEG	B	669	-	-	2/4/4/4	-
4	PEG	A	677	-	-	2/4/4/4	-
2	GOL	A	668	-	-	3/4/4/4	-
2	GOL	A	667	-	-	0/4/4/4	-
4	PEG	A	674	-	-	3/4/4/4	-

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	669	PEG	O2-C3-C4	3.19	124.18	110.11
4	A	677	PEG	O2-C3-C4	2.43	120.84	110.11
4	B	669	PEG	C3-O2-C2	2.35	123.56	113.26
4	B	669	PEG	O2-C2-C1	2.28	120.14	110.11
4	A	678	PEG	O2-C2-C1	2.26	120.07	110.11

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	669	PEG	C4-C3-O2-C2
2	A	668	GOL	O1-C1-C2-C3
4	A	677	PEG	O1-C1-C2-O2
4	B	669	PEG	O2-C3-C4-O4
2	A	668	GOL	O1-C1-C2-O2

There are no ring outliers.

10 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	678	PEG	3	0
3	B	672	FMT	1	0
3	B	675	FMT	1	0
3	B	668	FMT	3	0
4	A	677	PEG	4	0
3	B	674	FMT	2	0
2	A	667	GOL	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	672	FMT	1	0
3	A	669	FMT	3	0
3	A	673	FMT	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	642/690 (93%)	-0.12	16 (2%) 57 65	17, 42, 77, 120	0
1	B	642/690 (93%)	0.22	31 (4%) 30 39	19, 53, 91, 126	0
All	All	1284/1380 (93%)	0.05	47 (3%) 41 49	17, 47, 87, 126	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	144	TYR	5.1
1	B	142	ASP	5.0
1	A	142	ASP	4.3
1	B	269	VAL	3.9
1	A	144	TYR	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PEG	A	678	7/7	0.36	0.25	76,78,82,83	0
4	PEG	A	674	7/7	0.53	0.35	78,80,80,81	0
3	FMT	B	672	3/3	0.70	0.17	60,60,65,66	0
3	FMT	B	674	3/3	0.77	0.17	46,46,47,50	0
4	PEG	B	669	7/7	0.77	0.26	50,58,66,67	0
3	FMT	A	673	3/3	0.79	0.19	45,45,47,49	0
3	FMT	B	673	3/3	0.84	0.21	65,65,70,71	0
7	SO4	B	671	5/5	0.84	0.37	121,121,122,124	0
2	GOL	A	667	6/6	0.85	0.19	43,53,58,58	0
2	GOL	A	668	6/6	0.86	0.15	64,68,72,74	0
3	FMT	B	675	3/3	0.88	0.12	65,65,66,67	0
3	FMT	A	672	3/3	0.89	0.16	54,54,58,61	0
4	PEG	A	677	7/7	0.90	0.36	68,71,73,74	0
3	FMT	A	669	3/3	0.90	0.16	48,48,51,53	0
3	FMT	B	667	3/3	0.92	0.19	41,41,47,49	0
3	FMT	A	671	3/3	0.92	0.11	53,53,57,59	0
5	MG	B	670	1/1	0.94	0.09	42,42,42,42	0
5	MG	A	675	1/1	0.94	0.10	43,43,43,43	0
3	FMT	A	670	3/3	0.95	0.09	44,44,45,47	0
3	FMT	B	668	3/3	0.96	0.12	43,43,45,47	0
6	CL	A	676	1/1	0.97	0.08	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.