

wwPDB X-ray Structure Validation Summary Report (i)

Nov 15, 2023 – 02:54 AM JST

PDB ID : 8I01

Title : Crystal structure of Escherichia coli glyoxylate carboligase

Authors: Kim, J.H.; Kim, J.S.

Deposited on : 2023-01-10

Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(\mathring{A}))$
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	594	92%	8%
1	В	594	90%	8% •
1	С	594	91%	8% •
1	D	594	92%	8% •
1	Е	594	88%	10% •
1	F	594	92%	8% •



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	UQ0	С	705	-	-	=	X
5	UQ0	С	706	-	-	=	X
5	UQ0	D	707	-	-	=	X



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 30468 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glyoxylate carboligase.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	594	Total	С	N	О	S	0	0	0
1	Λ	094	4542	2878	794	836	34	U	0	
1	В	594	Total	С	N	О	S	0	0	0
1	Ъ	094	4541	2878	793	836	34	U	0	0
1	С	593	Total	С	N	О	S	0	0	0
1		090	4538	2876	793	835	34	0		
1	D	594	Total	С	N	О	S	0	0	0
1	D	094	4542	2878	794	836	34	U	0	
1	Е	593	Total	С	N	О	S	0	0	0
1	l L	090	4538	2876	793	835	34	U	0	
1	F	594	Total	С	N	О	S	0	0	0
	Г	J94	4542	2878	794	836	34	U	U	U

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P0AEP7
A	488	LEU	MET	engineered mutation	UNP P0AEP7
В	0	GLY	-	expression tag	UNP P0AEP7
В	488	LEU	MET	engineered mutation	UNP P0AEP7
С	0	GLY	-	expression tag	UNP P0AEP7
С	488	LEU	MET	engineered mutation	UNP P0AEP7
D	0	GLY	-	expression tag	UNP P0AEP7
D	488	LEU	MET	engineered mutation	UNP P0AEP7
Е	0	GLY	-	expression tag	UNP P0AEP7
Е	488	LEU	MET	engineered mutation	UNP P0AEP7
F	0	GLY	-	expression tag	UNP P0AEP7
F	488	LEU	MET	engineered mutation	UNP P0AEP7

• Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
	A	1	53	27	9	15	2	0	U
2	В	1	Total	С	N	О	Р	0	0
2	Б	1	53	27	9	15	2	U	0
2	C	1	Total	С	N	О	Р	0	0
2		1	53	27	9	15	2		
2	D	1	Total	С	N	О	Р	0	0
	D	1	53	27	9	15	2	U	U
2	Е	1	Total	С	N	О	Р	0	0
	Ľ	1	53	27	9	15	2	U	U
2	F	1	Total	С	N	О	Р	0	0
	1'	1	53	27	9	15	2	0	

• Molecule 3 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: $C_{12}H_{19}N_4O_7P_2S$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues		A	ton	ıs			ZeroOcc	AltConf
3	A	1	Total	С	N	О	Р	S	0	0
3	Λ	1	26	12	4	7	2	1	0	0
3	В	1	Total	С	N	О	Р	S	0	0
	D	1	26	12	4	7	2	1		U
3	С	1	Total	С	N	О	Р	S	0	0
0	C	1	26	12	4	7	2	1		0
3	D	1	Total	С	N	О	Р	S	0	0
	D	1	26	12	4	7	2	1	U	U
3	E	1	Total	С	N	Ο	Р	S	0	0
	נו	1	26	12	4	7	2	1	U	U
3	F	1	Total	С	N	О	Р	S	0	0
)	1'	1	26	12	4	7	2	1		

 \bullet Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Mg 2 2	0	0
4	В	2	Total Mg 2 2	0	0
4	С	1	Total Mg 1 1	0	0
4	D	2	Total Mg 2 2	0	0
4	E	2	Total Mg 2 2	0	0
4	F	1	Total Mg 1 1	0	0



• Molecule 5 is 2,3-DIMETHOXY-5-METHYL-1,4-BENZOQUINONE (three-letter code: UQ0) (formula: $C_9H_{10}O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 13 9 4	0	0
5	A	1	Total C O 13 9 4	0	0
5	В	1	Total C O 13 9 4	0	0
5	В	1	Total C O 13 9 4	0	0
5	С	1	Total C O 13 9 4	0	0
5	С	1	Total C O 13 9 4	0	0
5	С	1	Total C O 13 9 4	0	0
5	D	1	Total C O 13 9 4	0	0
5	D	1	Total C O 13 9 4	0	0
5	D	1	Total C O 13 9 4	0	0
5	Е	1	Total C O 13 9 4	0	0
5	F	1	Total C O 13 9 4	0	0

• Molecule 6 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	531	Total O 531 531	0	0
6	В	486	Total O 486 486	0	0
6	С	402	Total O 402 402	0	0
6	D	400	Total O 400 400	0	0
6	E	396	Total O 396 396	0	0
6	F	370	Total O 370 370	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glyoxylate carboligase








• Molecule 1: Glyoxylate carboligase



• Molecule 1: Glyoxylate carboligase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	189.62Å 189.62Å 248.63Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.58 - 2.15	Depositor
Resolution (A)	48.58 - 2.15	EDS
% Data completeness	99.7 (48.58-2.15)	Depositor
(in resolution range)	99.7 (48.58-2.15)	EDS
R_{merge}	0.99	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.91 (at 2.16Å)	Xtriage
Refinement program	PHENIX v2.0	Depositor
D D.	0.191 , 0.231	Depositor
R, R_{free}	0.192 , 0.231	DCC
R_{free} test set	9711 reflections (4.00%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 33.5	EDS
L-test for twinning ²	$ < L >=0.53, < L^2>=0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	30468	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 66.55 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.9354e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FAD, UQ0, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.47	0/4634	0.69	$2/6290 \ (0.0\%)$	
1	В	0.47	$2/4633 \ (0.0\%)$	0.70	5/6289 (0.1%)	
1	С	0.43	0/4630	0.64	0/6285	
1	D	0.42	0/4634	0.63	$3/6290 \ (0.0\%)$	
1	Е	0.42	0/4630	0.64	$1/6285 \ (0.0\%)$	
1	F	0.41	0/4634	0.62	0/6290	
All	All	0.44	$2/27795 \ (0.0\%)$	0.65	11/37729 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	В	0	2
1	С	0	1
All	All	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
1	В	82	ASP	CG-OD2	6.91	1.41	1.25
1	В	82	ASP	CB-CG	6.03	1.64	1.51

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	488	LEU	CA-CB-CG	13.75	146.93	115.30
1	В	82	ASP	CB-CG-OD1	-9.57	109.68	118.30



Continued from previous page...

Mol	Chain	Res	Type	Atoms	${f Z}$	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	Ε	488	LEU	CA-CB-CG	7.92	133.51	115.30
1	В	82	ASP	CB-CG-OD2	6.69	124.32	118.30
1	D	488	LEU	CA-CB-CG	6.23	129.64	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	296	ARG	Sidechain
1	В	296	ARG	Sidechain
1	В	592	TYR	Peptide
1	С	296	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4542	0	4550	26	0
1	В	4541	0	4545	38	1
1	С	4538	0	4547	34	0
1	D	4542	0	4550	34	0
1	Е	4538	0	4547	48	0
1	F	4542	0	4550	38	0
2	A	53	0	31	0	0
2	В	53	0	31	1	0
2	С	53	0	31	1	0
2	D	53	0	31	3	0
2	Ε	53	0	31	3	0
2	F	53	0	31	3	0
3	A	26	0	16	1	0
3	В	26	0	16	1	1
3	С	26	0	16	0	0
3	D	26	0	16	0	0
3	E	26	0	16	0	0
3	F	26	0	16	0	0
4	A	2	0	0	0	0
4	В	2	0	0	0	0



Continued from previous page						
	\mathcal{C}	lontinued	l fr	rom	previous	page

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	С	1	0	0	0	0
4	D	2	0	0	0	0
4	${ m E}$	2	0	0	0	0
4	F	1	0	0	0	0
5	A	26	0	0	1	2
5	В	26	0	0	2	2
5	С	39	0	0	3	0
5	D	39	0	0	2	0
5	Ε	13	0	0	0	0
5	F	13	0	0	0	0
6	A	531	0	0	6	1
6	В	486	0	0	9	1
6	С	402	0	0	5	0
6	D	400	0	0	2	0
6	Ε	396	0	0	6	0
6	F	370	0	0	8	0
All	All	30468	0	27571	207	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 207 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:D:292:TYR:O	1:D:296:ARG:NH1	2.03	0.91
1:B:384:GLY:HA2	1:B:593:GLU:HG2	1.59	0.85
1:E:361:ARG:HD3	1:E:572:GLU:OE2	1.77	0.85
1:F:393:ILE:HA	1:F:397:GLN:HG2	1.61	0.83
1:E:237:THR:HG22	1:E:240:GLY:H	1.46	0.80

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:82:ASP:OD2	3:B:702:TPP:CM2[8_554]	1.43	0.77
5:A:705:UQ0:CM2	5:B:705:UQ0:CM3[3_555]	1.49	0.71
5:A:706:UQ0:O1	6:A:820:HOH:O[8_554]	2.11	0.09
5:B:706:UQ0:O1	6:B:886:HOH:O[8_554]	2.12	0.08



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	tiles
1	A	592/594~(100%)	583 (98%)	9 (2%)	0	100	100
1	В	592/594 (100%)	581 (98%)	11 (2%)	0	100	100
1	С	591/594 (100%)	583 (99%)	8 (1%)	0	100	100
1	D	592/594 (100%)	582 (98%)	10 (2%)	0	100	100
1	E	591/594 (100%)	580 (98%)	11 (2%)	0	100	100
1	F	592/594 (100%)	584 (99%)	8 (1%)	0	100	100
All	All	3550/3564 (100%)	3493 (98%)	57 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	$475/475 \; (100\%)$	462 (97%)	13 (3%)	44 46
1	В	474/475 (100%)	461 (97%)	13 (3%)	44 46
1	С	475/475 (100%)	457 (96%)	18 (4%)	33 31
1	D	475/475 (100%)	463 (98%)	12 (2%)	47 49
1	E	475/475 (100%)	458 (96%)	17 (4%)	35 33
1	F	475/475 (100%)	464 (98%)	11 (2%)	50 53
All	All	2849/2850 (100%)	2765 (97%)	84 (3%)	42 42

5 of 84 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	${ m E}$	108	ARG
1	Ε	561	ILE
1	Ε	124	LYS
1	Ε	299	VAL
1	F	108	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	569	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 34 ligands modelled in this entry, 10 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain Res	Res 1	Link	Bond lengths			В	Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2														
2	FAD	A	701	-	53,58,58	0.63	1 (1%)	68,89,89	0.67	1 (1%)														
3	TPP	В	702	4	22,27,27	1.19	2 (9%)	29,40,40	1.23	3 (10%)														
2	FAD	В	701	-	53,58,58	0.59	1 (1%)	68,89,89	0.70	2 (2%)														



Mol	Type	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	cles
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TPP	D	702	4	22,27,27	0.65	0	29,40,40	0.86	1 (3%)
5	UQ0	В	705	-	13,13,13	3.37	5 (38%)	16,18,18	1.51	3 (18%)
2	FAD	С	701	-	53,58,58	0.47	0	68,89,89	0.66	1 (1%)
3	TPP	С	702	4	22,27,27	0.57	0	29,40,40	0.73	1 (3%)
3	TPP	F	702	4	22,27,27	0.81	0	29,40,40	0.90	1 (3%)
2	FAD	D	701	-	53,58,58	0.53	0	68,89,89	0.57	1 (1%)
5	UQ0	D	705	-	13,13,13	3.50	5 (38%)	16,18,18	1.32	1 (6%)
3	TPP	Е	702	4	22,27,27	0.67	0	29,40,40	0.87	0
2	FAD	F	701	-	53,58,58	0.51	0	68,89,89	0.63	1 (1%)
5	UQ0	A	706	-	13,13,13	3.45	7 (53%)	16,18,18	2.54	2 (12%)
5	UQ0	С	705	-	13,13,13	3.42	8 (61%)	16,18,18	1.13	1 (6%)
5	UQ0	D	706	-	13,13,13	3.53	5 (38%)	16,18,18	1.30	2 (12%)
5	UQ0	D	707	-	13,13,13	3.51	6 (46%)	16,18,18	1.37	1 (6%)
5	UQ0	Е	705	-	13,13,13	3.36	6 (46%)	16,18,18	1.31	2 (12%)
3	TPP	A	702	4	22,27,27	0.54	0	29,40,40	0.83	1 (3%)
5	UQ0	С	706	-	13,13,13	3.58	7 (53%)	16,18,18	1.69	1 (6%)
5	UQ0	В	706	-	13,13,13	3.49	8 (61%)	16,18,18	1.97	2 (12%)
2	FAD	Е	701	-	53,58,58	0.49	0	68,89,89	0.61	1 (1%)
5	UQ0	A	705	-	13,13,13	3.51	5 (38%)	16,18,18	1.40	1 (6%)
5	UQ0	С	704	-	13,13,13	3.44	6 (46%)	16,18,18	1.26	1 (6%)
5	UQ0	F	704	-	13,13,13	3.47	8 (61%)	16,18,18	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	701	-	-	7/30/50/50	0/6/6/6
3	TPP	В	702	4	-	2/16/17/17	0/2/2/2
2	FAD	В	701	-	-	4/30/50/50	0/6/6/6
3	TPP	D	702	4	-	2/16/17/17	0/2/2/2
5	UQ0	В	705	-	-	0/4/24/24	0/1/1/1
2	FAD	С	701	-	-	7/30/50/50	0/6/6/6
3	TPP	С	702	4	-	2/16/17/17	0/2/2/2
3	TPP	F	702	4	-	1/16/17/17	0/2/2/2



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	D	701	-	-	6/30/50/50	0/6/6/6
5	UQ0	D	705	-	-	0/4/24/24	0/1/1/1
3	TPP	Е	702	4	-	1/16/17/17	0/2/2/2
2	FAD	F	701	-	-	6/30/50/50	0/6/6/6
5	UQ0	A	706	-	-	2/4/24/24	0/1/1/1
5	UQ0	С	705	-	-	2/4/24/24	0/1/1/1
5	UQ0	D	706	-	-	0/4/24/24	0/1/1/1
5	UQ0	D	707	-	-	1/4/24/24	0/1/1/1
5	UQ0	Е	705	-	-	0/4/24/24	0/1/1/1
3	TPP	A	702	4	-	2/16/17/17	0/2/2/2
5	UQ0	С	706	-	-	2/4/24/24	0/1/1/1
5	UQ0	В	706	-	-	1/4/24/24	0/1/1/1
2	FAD	Е	701	-	-	6/30/50/50	0/6/6/6
5	UQ0	A	705	-	-	0/4/24/24	0/1/1/1
5	UQ0	С	704	-	-	0/4/24/24	0/1/1/1
5	UQ0	F	704	-	-	0/4/24/24	0/1/1/1

The worst 5 of 80 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
5	D	705	UQ0	C6-C5	9.57	1.54	1.35
5	F	704	UQ0	C6-C5	9.32	1.53	1.35
5	В	705	UQ0	C6-C5	9.25	1.53	1.35
5	D	707	UQ0	C6-C5	9.08	1.53	1.35
5	С	704	UQ0	C6-C5	9.07	1.53	1.35

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	A	706	UQ0	CM5-C5-C4	8.96	123.56	117.45
5	В	706	UQ0	CM5-C5-C4	6.17	121.66	117.45
5	С	706	UQ0	CM5-C5-C4	5.72	121.35	117.45
5	D	707	UQ0	CM5-C5-C4	3.88	120.09	117.45
5	D	706	UQ0	CM5-C5-C4	3.53	119.86	117.45

There are no chirality outliers.

5 of 54 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	FAD	C5B-O5B-PA-O3P



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	В	701	FAD	C5B-O5B-PA-O1A
2	С	701	FAD	C5B-O5B-PA-O1A
2	D	701	FAD	C5B-O5B-PA-O1A
2	D	701	FAD	C5B-O5B-PA-O3P


There are no ring outliers.

14 monomers are involved in 25 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	702	TPP	1	1
2	В	701	FAD	1	0
5	В	705	UQ0	0	1
2	С	701	FAD	1	0
2	D	701	FAD	3	0
2	F	701	FAD	3	0
5	A	706	UQ0	1	1
5	С	705	UQ0	3	0
5	D	706	UQ0	1	0
5	D	707	UQ0	1	0
3	A	702	TPP	1	0
5	В	706	UQ0	2	1
2	Е	701	FAD	3	0
5	A	705	UQ0	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	594/594 (100%)	-0.59	0 100 100	17, 21, 34, 54	0
1	В	594/594 (100%)	-0.37	1 (0%) 95 96	17, 23, 38, 62	0
1	С	593/594 (99%)	-0.60	2 (0%) 94 95	20, 28, 42, 66	0
1	D	594/594 (100%)	-0.49	6 (1%) 82 86	20, 29, 45, 72	0
1	E	593/594 (99%)	-0.56	2 (0%) 94 95	21, 30, 46, 66	0
1	F	594/594 (100%)	-0.29	10 (1%) 70 76	21, 31, 50, 74	0
All	All	3562/3564~(99%)	-0.48	21 (0%) 89 91	17, 27, 44, 74	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	485	ALA	4.3
1	F	504	VAL	3.3
1	F	593	GLU	3.1
1	F	3	LYS	2.9
1	С	3	LYS	2.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
5	UQ0	D	706	13/13	0.30	0.40	78,86,103,106	0
5	UQ0	С	705	13/13	0.42	0.63	93,104,121,123	0
5	UQ0	С	706	13/13	0.57	0.48	73,78,92,94	0
5	UQ0	D	707	13/13	0.57	0.44	91,97,106,112	0
5	UQ0	В	706	13/13	0.62	0.31	77,82,96,97	0
5	UQ0	A	706	13/13	0.64	0.35	64,69,77,84	0
5	UQ0	Е	705	13/13	0.70	0.25	72,79,82,84	0
5	UQ0	С	704	13/13	0.73	0.25	58,62,65,65	0
5	UQ0	A	705	13/13	0.82	0.21	46,54,58,60	0
5	UQ0	D	705	13/13	0.89	0.19	43,47,55,57	0
5	UQ0	F	704	13/13	0.89	0.15	46,55,65,67	0
5	UQ0	В	705	13/13	0.91	0.12	36,41,47,51	0
3	TPP	F	702	26/26	0.93	0.15	26,36,41,43	0
3	TPP	В	702	26/26	0.95	0.10	21,26,30,33	0
4	MG	D	703	1/1	0.97	0.06	28,28,28,28	0
4	MG	F	703	1/1	0.97	0.04	33,33,33,33	0
2	FAD	С	701	53/53	0.97	0.08	21,24,26,27	0
3	TPP	D	702	26/26	0.97	0.09	24,30,33,35	0
3	TPP	Е	702	26/26	0.97	0.10	27,31,36,38	0
2	FAD	E	701	53/53	0.97	0.08	23,26,31,32	0
4	MG	В	703	1/1	0.97	0.05	23,23,23,23	0
4	MG	A	704	1/1	0.98	0.06	17,17,17,17	1
3	TPP	A	702	26/26	0.98	0.07	18,20,23,25	0
2	FAD	A	701	53/53	0.98	0.11	19,19,21,24	0
3	TPP	С	702	26/26	0.98	0.07	21,26,29,31	0
2	FAD	D	701	53/53	0.98	0.09	22,24,27,28	0
2	FAD	В	701	53/53	0.98	0.11	16,18,22,23	0
2	FAD	F	701	53/53	0.98	0.10	19,25,30,32	0
4	MG	A	703	1/1	0.98	0.12	19,19,19,19	0
4	MG	Е	703	1/1	0.99	0.05	26,26,26,26	0
4	MG	Е	704	1/1	0.99	0.05	29,29,29,29	0
4	MG	С	703	1/1	0.99	0.11	21,21,21,21	0
4	MG	В	704	1/1	0.99	0.07	18,18,18,18	1
4	MG	D	704	1/1	0.99	0.05	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers




as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around UQ0 C 705: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)























Electron density around UQ0 E 705: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)













Electron density around UQ0 D 705: $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around TPP B 702:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around FAD C 701:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around TPP D 702:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around TPP E 702: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around FAD E 701: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)









Electron density around TPP A 702: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around FAD A 701: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around TPP C 702:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FAD D 701:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around FAD B 701:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FAD F 701:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

