

wwPDB X-ray Structure Validation Summary Report (i)

Nov 13, 2023 – 05:35 PM JST

PDB ID : 8I09

Title : Crystal structure of serine acetyltransferase from Salmonella typhimurium

complexed with butyl gallate

Authors: Toyomoto, T.; Ono, K.; Shiba, T.; Momitani, K.; Zhang, T.; Tsutsuki, H.;

Ishikawa, T.; Hoso, K.; Hamada, K.; Rahman, A.; Zhong, H.; Akaike, T.;

Yamamoto, K.; Matsuoka, M.; Hanaoka, K.; Niidome, T.; Sawa, T.

Deposited on : 2023-01-10

Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36




1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.80 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# ext{Entries}, ext{ resolution range}(\mathring{A}))$
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	280	.% 82%	5% • 12%
1	В	280	.% 82%	5% 12%
1	С	280	79%	9% 12%
1	D	280	81%	6% • 12%
1	Е	280	79%	8% • 12%
1	F	280	79%	7% • 12%





Continued from previous page...

Mol	Chain	Length	Quality of chain		
	~	222	<u>%</u>		
1	G	280	80%	7% •	12%
			<u>%</u>		
1	Н	280	81%	6% •	12%
			2%		
1	I	280	81%	6% •	12%
			. <mark>%</mark>		
1	J	280	83%		12%
			2%		
1	K	280	81%	6% •	12%
			2%		
1	L	280	80%	8%	12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CYS	J	401	-	X	-	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 23191 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Serine acetyltransferase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	246	Total	С	N	О	S	0	1	0
1	A	240	1852	1178	328	336	10	U	1	
1	В	246	Total	С	N	О	S	0	0	0
1	Ъ	240	1844	1173	327	335	9	U	U	
1	С	246	Total	\mathbf{C}	N	O	S	0	1	0
1		240	1852	1178	328	336	10	O	1	
1	D	246	Total	\mathbf{C}	N	Ο	S	0	0	0
1	D	240	1842	1171	327	335	9	O	U	
1	E	245	Total	\mathbf{C}	N	Ο	S	0	0	0
1	L	240	1837	1168	326	334	9	O	U	
1	F	245	Total	\mathbf{C}	N	O	S	0	0	0
1	I.	240	1837	1168	326	334	9			
1	G	246	Total	$^{\mathrm{C}}$	Ν	O	S	0	1	0
1	d	240	1850	1176	328	336	10	O	1	
1	Н	247	Total	\mathbf{C}	N	O	S	0	1	0
	11	211	1857	1181	329	337	10	0	1	0
1	I	246	Total	С	Ν	O	S	0	1	0
1	1	240	1850	1176	328	336	10	O	1	0
1	J	246	Total	С	N	О	S	0	0	0
1	3	240	1844	1173	327	335	9	O	U	
1	K	246	Total	С	N	О	S	0	0	0
1	17	240	1844	1173	327	335	9	U	U	
1	L	246	Total	\mathbf{C}	N	О	S	0	1	0
	П	240	1852	1178	328	336	10		1	

There are 108 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
A	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
A	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
A	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
A	-3	HIS	-	expression tag	UNP A0A0D6I3Y9



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
A	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
A	0	ASP	-	expression tag	UNP A0A0D6I3Y9
A	1	PRO	-	expression tag	UNP A0A0D6I3Y9
В	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
В	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
В	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
В	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
В	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
В	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
В	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
В	0	ASP	-	expression tag	UNP A0A0D6I3Y9
В	1	PRO	-	expression tag	UNP A0A0D6I3Y9
С	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
С	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
С	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
С	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
С	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
С	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
С	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
С	0	ASP	-	expression tag	UNP A0A0D6I3Y9
С	1	PRO	-	expression tag	UNP A0A0D6I3Y9
D	-7	MET	ı	initiating methionine	UNP A0A0D6I3Y9
D	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
D	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
D	-4	HIS	ı	expression tag	UNP A0A0D6I3Y9
D	-3	HIS	ı	expression tag	UNP A0A0D6I3Y9
D	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
D	-1	HIS	ı	expression tag	UNP A0A0D6I3Y9
D	0	ASP	-	expression tag	UNP A0A0D6I3Y9
D	1	PRO	-	expression tag	UNP A0A0D6I3Y9
E	-7	MET	ı	initiating methionine	UNP A0A0D6I3Y9
Е	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
Е	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
Е	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
Е	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
Е	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
Е	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
Е	0	ASP	_	expression tag	UNP A0A0D6I3Y9
Е	1	PRO	-	expression tag	UNP A0A0D6I3Y9
F	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
F	-6	HIS	-	expression tag	UNP A0A0D6I3Y9



Continued from previous page...


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
F	-5	HIS	_	expression tag	UNP A0A0D6I3Y9
F	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
F	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
F	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
F	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
F	0	ASP	_	expression tag	UNP A0A0D6I3Y9
F	1	PRO	-	expression tag	UNP A0A0D6I3Y9
G	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
G	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
G	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
G	-4	HIS	_	expression tag	UNP A0A0D6I3Y9
G	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
G	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
G	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
G	0	ASP	-	expression tag	UNP A0A0D6I3Y9
G	1	PRO	-	expression tag	UNP A0A0D6I3Y9
Н	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
Н	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
Н	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
Н	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
Н	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
Н	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
Н	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
Н	0	ASP	-	expression tag	UNP A0A0D6I3Y9
Н	1	PRO	-	expression tag	UNP A0A0D6I3Y9
I	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
I	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
I	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
I	-4	HIS	_	expression tag	UNP A0A0D6I3Y9
I	-3	HIS	_	expression tag	UNP A0A0D6I3Y9
I	-2	HIS	_	expression tag	UNP A0A0D6I3Y9
I	-1	HIS	_	expression tag	UNP A0A0D6I3Y9
I	0	ASP	_	expression tag	UNP A0A0D6I3Y9
I	1	PRO	-	expression tag	UNP A0A0D6I3Y9
J	-7	MET	_	initiating methionine	UNP A0A0D6I3Y9
J	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
J	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
J	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
J	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
J	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
J	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
J	0	ASP	-	expression tag	UNP A0A0D6I3Y9



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
J	1	PRO	-	expression tag	UNP A0A0D6I3Y9
K	-7	MET	- initiating methionine		UNP A0A0D6I3Y9
K	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
K	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
K	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
K	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
K	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
K	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
K	0	ASP	-	expression tag	UNP A0A0D6I3Y9
K	1	PRO	-	expression tag	UNP A0A0D6I3Y9
L	-7	MET	-	initiating methionine	UNP A0A0D6I3Y9
L	-6	HIS	-	expression tag	UNP A0A0D6I3Y9
L	-5	HIS	-	expression tag	UNP A0A0D6I3Y9
L	-4	HIS	-	expression tag	UNP A0A0D6I3Y9
L	-3	HIS	-	expression tag	UNP A0A0D6I3Y9
L	-2	HIS	-	expression tag	UNP A0A0D6I3Y9
L	-1	HIS	-	expression tag	UNP A0A0D6I3Y9
L	0	ASP	-	expression tag	UNP A0A0D6I3Y9
L	1	PRO	-	expression tag	UNP A0A0D6I3Y9

• Molecule 2 is CYSTEINE (three-letter code: CYS) (formula: $C_3H_7NO_2S$).



\mathbf{Mol}	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total 7					0	0
2	A	1	Total 7	C 3	N 1	O 2	S 1	0	0



Continued from previous page...

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
2	В	1	Total	С	N	O	S	0	0	
		_	7	3	1	2	1	,	Ů,	
2	D	1	Total	С	N	О	S	0	0	
	D	1	7	3	1	2	1	Ŭ	Ŭ	
2	E	1	Total	С	N	Ο	S	0	0	
	L	1	7	3	1	2	1	U	U	
2	F	1	Total	С	Ν	Ο	\mathbf{S}	0	0	
2	Г	1	7	3	1	2	1		0	
2	G	1	Total	С	N	О	S	0	0	
	G	G	1	7	3	1	2	1	0	0
2	TT	1	Total	С	N	О	S	0	0	
2	Н	1	7	3	1	2	1	0		
0	т	1	Total	С	N	О	S	0	0	
2	I	1	7	3	1	2	1	0	0	
2	J	1	Total	С	N	О	S	0	0	
2	J	1	7	3	1	2	1	0	0	
2	J	1	Total	С	N	О	S	0	0	
	J	1	7	3	1	2	1	0	U	
0	K	1	Total	С	N	О	S	0	0	
2	N.	1	7	3	1	2	1	0	0	

• Molecule 3 is butyl 3,4,5-tris(oxidanyl)benzoate (three-letter code: NF0) (formula: $C_{11}H_{14}O_5$) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 16	C 11	O 5	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total C O 16 11 5	0	0
3	С	1	Total C O 16 11 5	0	0
3	Е	1	Total C O 16 11 5	0	0
3	Е	1	Total C O 16 11 5	0	0
3	F	1	Total C O 16 11 5	0	0
3	G	1	Total C O 16 11 5	0	0
3	Н	1	Total C O 16 11 5	0	0
3	I	1	Total C O 16 11 5	0	0
3	J	1	Total C O 16 11 5	0	0
3	L	1	Total C O 16 11 5	0	0
3	L	1	Total C O 16 11 5	0	0

 \bullet Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: $\mathrm{O_4P}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
1	٨	1	Total O P	0	0
4	Λ	1	5 4 1	U	
1	F	1	Total O P	0	0
4	I'	1	5 4 1	0	0
1	G	1	Total O P	0	0
4	G	1	5 4 1		0
1	I/	1	Total O P	0	0
4	K	1	5 4 1		U

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
5	A	71	Total O	0	0	
9	Λ	7.1	71 71	U	U	
5	В	67	Total O	0	0	
	D	01	67 67	U	U	
5	C	67	Total O	0	0	
		01	67 67	Ů	Ü	
5	D	52	Total O	0	0	
		92	52 52		Ü	
5	E	54	Total O	0	0	
	_	0 -	54 54	,	-	
5	F	51	Total O	0	0	
			51 51			
5	G	62	Total O	0	0	
			62 62			
5	Н	62	Total O 62 62	0	0	
			Total O			
5	I	60	60 60	0	0	
			Total O			
5	J	63	63 63	0	0	
			Total O			
5	K	61	61 61	0	0	
			Total O			
5	L	64	64 64	0	0	



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Serine acetyltransferase







HIS HIS HIS PHE GLU TYR GLY ASP GLY

• Molecule 1: Serine acetyltransferase







• Molecule 1: Serine acetyltransferase



• Molecule 1: Serine acetyltransferase




• Molecule 1: Serine acetyltransferase






GLN HIS PHE ASN GLY ILE HIS HIS PHE GLU THR PHE GLU TYR GLV


• Molecule 1: Serine acetyltransferase



• Molecule 1: Serine acetyltransferase



• Molecule 1: Serine acetyltransferase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	177.36Å 102.65Å 158.72Å	Donositon
a, b, c, α , β , γ	90.00° 111.85° 90.00°	Depositor
Resolution (Å)	19.96 - 1.80	Depositor
Resolution (A)	19.96 - 1.80	EDS
% Data completeness	99.3 (19.96-1.80)	Depositor
(in resolution range)	98.8 (19.96-1.80)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.42 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
D D	0.204 , 0.241	Depositor
R, R_{free}	0.213 , 0.247	DCC
R_{free} test set	12032 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 28.1	EDS
L-test for twinning ²	$< L > = 0.52, < L^2> = 0.35$	Xtriage
	0.438 for 1/2 *h + 3/2 *k, 1/2 *h - 1/2 *k, -1/2 *h	
Estimated twinning fraction	1/2*k-l	Xtriage
	0.438 for 1/2 +h- 3/2 +k, -1/2 +h- 1/2 +k, -1/2 +h	
D.D. L.:	+1/2*k-l	EDG
F_o, F_c correlation	0.97	EDS
Total number of atoms	23191	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 47.80 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.4291e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, NF0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond lengths		Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.53	0/1889	0.70	1/2564~(0.0%)	
1	В	0.55	0/1881	0.73	1/2554~(0.0%)	
1	С	0.51	0/1889	0.73	1/2564~(0.0%)	
1	D	0.50	0/1878	0.71	1/2550~(0.0%)	
1	Е	0.53	0/1873	0.71	1/2542 (0.0%)	
1	F	0.52	0/1873	0.70	1/2543~(0.0%)	
1	G	0.51	0/1886	0.74	2/2560 (0.1%)	
1	Н	0.52	0/1894	0.73	1/2572~(0.0%)	
1	I	0.49	0/1886	0.71	2/2560 (0.1%)	
1	J	0.50	0/1881	0.72	0/2554	
1	K	0.53	0/1881	0.72	0/2554	
1	L	0.51	0/1889	0.75	0/2564	
All	All	0.52	0/22600	0.72	11/30681 (0.0%)	

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	Н	1	PRO	N-CA-CB	6.49	111.09	103.30
1	G	2	PRO	N-CA-CB	5.97	110.46	103.30
1	D	2	PRO	N-CA-CB	5.89	110.37	103.30
1	С	91	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	I	91	ARG	NE-CZ-NH2	-5.59	117.50	120.30

There are no chirality outliers.

There are no planarity outliers.



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1852	0	1898	8	0
1	В	1844	0	1890	6	0
1	С	1852	0	1898	12	0
1	D	1842	0	1884	15	0
1	Ε	1837	0	1883	8	0
1	F	1837	0	1882	12	0
1	G	1850	0	1892	10	0
1	Н	1857	0	1899	7	0
1	I	1850	0	1892	10	0
1	J	1844	0	1890	8	0
1	K	1844	0	1890	10	0
1	L	1852	0	1898	11	0
2	A	14	0	8	0	0
2	В	7	0	4	0	0
2	D	7	0	4	0	0
2	Ε	7	0	4	0	0
2	F	7	0	4	0	0
2	G	7	0	4	0	0
2	Н	7	0	4	0	0
2	I	7	0	4	0	0
2	J	14	0	8	1	0
2	K	7	0	4	0	0
3	A	16	0	0	0	0
3	В	16	0	0	0	0
3	С	16	0	0	0	0
3	Ε	32	0	0	0	0
3	F	16	0	0	1	0
3	G	16	0	0	0	0
3	Н	16	0	0	0	0
3	I	16	0	0	0	0
3	J	16	0	0	0	0
3	L	32	0	0	0	0
4	A	5	0	0	0	0
4	F	5	0	0	0	0
4	G	5	0	0	0	0
4	K	5	0	0	0	0
5	A	71	0	0	1	0



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	В	67	0	0	2	0
5	С	67	0	0	2	0
5	D	52	0	0	0	0
5	Е	54	0	0	2	0
5	F	51	0	0	2	0
5	G	62	0	0	2	0
5	Н	62	0	0	2	0
5	I	60	0	0	2	0
5	J	63	0	0	2	0
5	K	61	0	0	2	0
5	L	64	0	0	1	0
All	All	23191	0	22744	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:D:4:GLU:OE2	1:D:8:ILE:HG12	1.80	0.81
1:J:111:LEU:HD21	1:J:154:ILE:HB	1.70	0.71
1:K:111:LEU:HD21	1:K:154:ILE:HB	1.76	0.67
1:F:3:CYS:SG	1:F:4:GLU:N	2.69	0.66
1:E:92:ASP:O	5:E:601:HOH:O	2.14	0.64

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/280 (88%)	239 (98%)	6 (2%)	0	100 100	



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	В	244/280 (87%)	240 (98%)	4 (2%)	0	100	100
1	С	245/280 (88%)	240 (98%)	5 (2%)	0	100	100
1	D	244/280 (87%)	240 (98%)	4 (2%)	0	100	100
1	E	243/280 (87%)	237 (98%)	3 (1%)	3 (1%)	13	3
1	F	243/280 (87%)	237 (98%)	6 (2%)	0	100	100
1	G	245/280 (88%)	238 (97%)	5 (2%)	2 (1%)	19	7
1	Н	246/280 (88%)	238 (97%)	8 (3%)	0	100	100
1	I	245/280 (88%)	237 (97%)	7 (3%)	1 (0%)	34	21
1	J	244/280 (87%)	238 (98%)	6 (2%)	0	100	100
1	K	244/280 (87%)	235 (96%)	6 (2%)	3 (1%)	13	3
1	L	245/280 (88%)	237 (97%)	8 (3%)	0	100	100
All	All	2933/3360 (87%)	2856 (97%)	68 (2%)	9 (0%)	41	27

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Е	6	LEU
1	G	188	THR
1	I	188	THR
1	K	5	GLU
1	K	6	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	$191/220\ (87\%)$	184 (96%)	7 (4%)	34	19	
1	В	190/220~(86%)	185 (97%)	5 (3%)	46	32	
1	С	191/220 (87%)	182 (95%)	9 (5%)	26	12	
1	D	189/220~(86%)	181 (96%)	8 (4%)	30	15	
1	Е	189/220 (86%)	180 (95%)	9 (5%)	25	11	



Continued from previous page...

Page 20

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	F	189/220 (86%)	180 (95%)	9 (5%)	25 11
1	G	190/220 (86%)	184 (97%)	6 (3%)	39 25
1	Н	191/220 (87%)	182 (95%)	9 (5%)	26 12
1	I	190/220 (86%)	183 (96%)	7 (4%)	34 19
1	J	190/220~(86%)	185 (97%)	5 (3%)	46 32
1	K	190/220 (86%)	183 (96%)	7 (4%)	34 19
1	L	191/220 (87%)	184 (96%)	7 (4%)	34 19
All	All	2281/2640 (86%)	2193 (96%)	88 (4%)	32 17

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	114	TYR
1	J	160	THR
1	Н	187	LYS
1	I	114	TYR
1	K	23	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	135	GLN
1	L	38	HIS
1	F	133	GLN
1	F	38	HIS
1	L	133	GLN

RNA (i) 5.3.3

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

28 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	T	Clasia.	Das	T :1-	Вс	ond leng	ths	Bond angles		
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CYS	A	401	-	5,6,6	1.23	0	5,7,7	1.81	2 (40%)
3	NF0	С	501	-	16,16,16	1.38	1 (6%)	21,21,21	1.33	2 (9%)
2	CYS	J	402	-	5,6,6	1.17	0	5,7,7	1.72	2 (40%)
3	NF0	L	502	-	16,16,16	1.37	1 (6%)	21,21,21	1.74	3 (14%)
2	CYS	I	502	-	5,6,6	1.11	1 (20%)	5,7,7	1.59	2 (40%)
2	CYS	В	401	-	5,6,6	1.28	0	5,7,7	1.57	2 (40%)
2	CYS	G	401	_	5,6,6	1.01	1 (20%)	5,7,7	1.72	2 (40%)
2	CYS	F	502	_	5,6,6	1.05	0	5,7,7	1.65	2 (40%)
2	CYS	A	402	-	5,6,6	1.10	0	5,7,7	1.48	1 (20%)
3	NF0	I	501	-	16,16,16	1.47	2 (12%)	21,21,21	1.51	3 (14%)
3	NF0	Н	501	-	16,16,16	1.46	2 (12%)	21,21,21	2.17	5 (23%)
2	CYS	Е	502	-	5,6,6	1.05	0	5,7,7	1.57	2 (40%)
4	PO4	F	503	-	4,4,4	0.79	0	6,6,6	0.59	0
3	NF0	J	403	-	16,16,16	1.35	2 (12%)	21,21,21	1.49	2 (9%)
3	NF0	F	501	-	16,16,16	1.32	1 (6%)	21,21,21	1.27	2 (9%)
4	PO4	A	404	-	4,4,4	0.85	0	6,6,6	0.55	0
3	NF0	E	503	_	16,16,16	1.29	1 (6%)	21,21,21	1.08	2 (9%)
3	NF0	A	403	-	16,16,16	1.41	2 (12%)	21,21,21	1.74	3 (14%)
4	PO4	K	402	-	4,4,4	0.83	0	6,6,6	0.46	0
2	CYS	Н	502	-	5,6,6	1.00	0	5,7,7	1.48	1 (20%)
4	PO4	G	403	-	4,4,4	0.90	0	6,6,6	0.45	0
3	NF0	Е	501	-	16,16,16	1.36	2 (12%)	21,21,21	1.95	4 (19%)
3	NF0	В	402	-	16,16,16	1.33	1 (6%)	21,21,21	2.08	4 (19%)



Mol	Tuno	Chain	Res	Res Link Bond lengths			Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NF0	L	501	-	16,16,16	1.47	3 (18%)	21,21,21	2.19	6 (28%)
2	CYS	J	401	-	5,6,6	1.28	0	5,7,7	2.14	2 (40%)
3	NF0	G	402	-	16,16,16	1.42	1 (6%)	21,21,21	1.41	3 (14%)
2	CYS	D	401	-	5,6,6	1.01	0	5,7,7	1.68	2 (40%)
2	CYS	K	401	-	5,6,6	1.23	0	5,7,7	1.76	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CYS	A	401	-	-	5/6/6/6	-
3	NF0	С	501	-	-	5/9/9/9	0/1/1/1
2	CYS	J	402	-	-	1/6/6/6	-
3	NF0	L	502	-	-	5/9/9/9	0/1/1/1
2	CYS	I	502	-	-	0/6/6/6	-
2	CYS	В	401	-	-	0/6/6/6	-
2	CYS	G	401	_	-	4/6/6/6	-
2	CYS	F	502	-	-	1/6/6/6	-
2	CYS	A	402	-	-	0/6/6/6	-
3	NF0	I	501	_	-	5/9/9/9	0/1/1/1
3	NF0	Н	501	-	-	4/9/9/9	0/1/1/1
2	CYS	E	502	-	-	1/6/6/6	-
3	NF0	J	403	-	-	5/9/9/9	0/1/1/1
3	NF0	F	501	-	-	5/9/9/9	0/1/1/1
3	NF0	E	503	-	-	5/9/9/9	0/1/1/1
3	NF0	A	403	-	-	5/9/9/9	0/1/1/1
2	CYS	Н	502	-	-	0/6/6/6	-
3	NF0	Е	501	-	-	4/9/9/9	0/1/1/1
3	NF0	В	402	-	-	4/9/9/9	0/1/1/1
3	NF0	L	501	-	-	4/9/9/9	0/1/1/1
2	CYS	J	401	-	-	6/6/6/6	-
3	NF0	G	402	-	-	6/9/9/9	0/1/1/1
2	CYS	D	401	-		0/6/6/6	
2	CYS	K	401	_	-	2/6/6/6	-

The worst 5 of 21 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	С	501	NF0	CAF-CAG	-4.69	1.39	1.50
3	F	501	NF0	CAF-CAG	-4.56	1.39	1.50
3	I	501	NF0	CAF-CAG	-4.38	1.39	1.50
3	G	402	NF0	CAF-CAG	-4.36	1.39	1.50
3	Н	501	NF0	CAF-CAG	-4.28	1.39	1.50

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	В	402	NF0	OAH-CAG-CAF	6.74	123.88	112.14
3	A	403	NF0	OAH-CAG-CAF	6.23	122.98	112.14
3	Е	501	NF0	OAH-CAG-CAF	6.20	122.94	112.14
3	L	501	NF0	OAH-CAG-CAF	5.86	122.35	112.14
3	Н	501	NF0	OAH-CAG-CAF	5.84	122.32	112.14

There are no chirality outliers.

5 of 77 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	CYS	O-C-CA-N
2	A	401	CYS	C-CA-CB-SG
2	G	401	CYS	O-C-CA-N
2	G	401	CYS	N-CA-CB-SG
2	G	401	CYS	C-CA-CB-SG

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	501	NF0	1	0
2	J	401	CYS	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient




equivalents in the CSD to analyse the geometry.
























































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(m \AA^2)$	Q<0.9
1	A	246/280 (87%)	-0.28	3 (1%) 79 76	23, 33, 59, 83	0
1	В	246/280 (87%)	-0.29	2 (0%) 86 84	22, 32, 58, 79	0
1	С	246/280 (87%)	-0.26	3 (1%) 79 76	22, 33, 60, 85	0
1	D	246/280 (87%)	-0.18	4 (1%) 72 68	26, 39, 68, 100	0
1	Е	245/280 (87%)	-0.24	1 (0%) 92 90	25, 39, 65, 84	0
1	F	245/280 (87%)	-0.23	6 (2%) 59 54	25, 39, 65, 92	0
1	G	246/280 (87%)	-0.13	4 (1%) 72 68	25, 35, 62, 86	0
1	Н	247/280 (88%)	-0.20	3 (1%) 79 76	24, 34, 64, 81	0
1	I	246/280 (87%)	-0.15	6 (2%) 59 54	25, 35, 64, 81	0
1	J	246/280 (87%)	-0.18	2 (0%) 86 84	24, 35, 61, 82	0
1	K	246/280 (87%)	-0.18	6 (2%) 59 54	24, 35, 62, 82	0
1	L	246/280 (87%)	-0.16	5 (2%) 65 61	24, 35, 61, 86	0
All	All	2951/3360 (87%)	-0.21	45 (1%) 73 70	22, 36, 63, 100	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	247	PRO	5.3
1	L	57	ILE	5.0
1	G	187	LYS	4.9
1	J	57	ILE	4.4
1	K	246	LYS	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	CYS	J	402	7/7	0.41	0.33	91,95,97,97	0
2	CYS	A	402	7/7	0.45	0.22	70,78,78,80	0
2	CYS	K	401	7/7	0.45	0.31	69,78,82,86	0
2	CYS	A	401	7/7	0.52	0.27	65,71,74,77	0
2	CYS	D	401	7/7	0.52	0.36	87,92,96,97	0
2	CYS	F	502	7/7	0.56	0.33	94,95,96,96	0
3	NF0	Ε	501	16/16	0.60	0.16	52,55,57,57	0
3	NF0	Н	501	16/16	0.60	0.18	41,48,51,51	0
2	CYS	I	502	7/7	0.61	0.34	72,80,82,82	0
2	CYS	В	401	7/7	0.63	0.20	56,64,65,68	0
2	CYS	J	401	7/7	0.63	0.34	77,84,85,85	0
3	NF0	В	402	16/16	0.64	0.14	40,48,53,53	0
3	NF0	L	501	16/16	0.65	0.14	41,47,53,53	0
2	CYS	Н	502	7/7	0.66	0.21	69,74,77,77	0
2	CYS	G	401	7/7	0.67	0.30	82,90,95,99	0
4	PO4	F	503	5/5	0.71	0.24	107,107,109,110	0
3	NF0	Ι	501	16/16	0.72	0.17	40,46,49,50	0
3	NF0	С	501	16/16	0.73	0.14	42,48,54,55	0
3	NF0	L	502	16/16	0.74	0.14	39,46,51,51	0
3	NF0	J	403	16/16	0.74	0.13	41,48,50,53	0
3	NF0	A	403	16/16	0.75	0.14	42,51,55,56	0
2	CYS	Ε	502	7/7	0.76	0.24	77,83,84,85	0
3	NF0	Ε	503	16/16	0.78	0.14	50,55,58,58	0
3	NF0	G	402	16/16	0.79	0.15	45,51,54,54	0
4	PO4	A	404	5/5	0.82	0.26	114,116,117,118	0
3	NF0	F	501	16/16	0.83	0.14	52,56,58,59	0
4	PO4	K	402	5/5	0.85	0.26	86,86,86,92	0
4	PO4	G	403	5/5	0.90	0.38	111,113,115,115	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different



orientation to approximate a three-dimensional view.


Electron density around NF0 E 501: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)


Electron density around NF0 H 501: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around NF0 B 402:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NF0 L 501:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NF0 I 501:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NF0 C 501:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NF0 J 403: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NF0 G 402:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NF0 F 501:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

