

wwPDB X-ray Structure Validation Summary Report (i)

Feb 10, 2024 – 06:07 PM EST

PDB ID 2I39

> Title Crystal structure of Vaccinia virus N1L protein

Authors Aoyagi, M.; Aleshin, A.E.; Stec, B.; Liddington, R.C.

2006-08-17 Deposited on

2.20 Å(reported) Resolution

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

> The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467

> 1.8.5 (274361), CSD as541be (2020) Mogul

Xtriage (Phenix) 1.13

EDS 2.36

20191225.v01 (using entries in the PDB archive December 25th 2019) Percentile statistics

> Refmac 5.8.0158

CCP4 7.0.044 (Gargrove)

Ideal geometry (proteins) Engh & Huber (2001) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \ resolution} \\ (\#{\rm Entries, \ resolution \ range(\AA)}) \end{array}$
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	137	64%	20%		15%
1	В	137	61%	23%	•	13%
1	С	137	69%	18%		13%
1	D	137	66%	21%		13%
1	Е	137	63%	22%	•	14%



Mol	Chain	Length	Quality of chain		
			.% -		
1	F	137	72%	15%	14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MPD	A	501	X	-	-	-
2	MPD	Е	502	X	-	-	-



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein N1.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	117	Total	С	N	О	S	0	0	0
1	A	111	974	615	165	186	8	0	0	U
1	В	119	Total	С	N	О	S	0	0	0
1	Б	119	983	620	167	188	8	0	0	
1	С	119	Total	С	N	О	S	0	0	0
1		119	983	620	167	188	8			
1	D	119	Total	С	N	О	S	0	0	0
1	ע	119	983	620	167	188	8	0		
1	Е	118	Total	С	N	О	S	0	0	0
1	12	110	978	617	166	187	8	0	0	U
1	F	110	Total	С	N	О	S	0	0	0
	Г	118	978	617	166	187	8		U	U

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	cloning artifact	UNP P21054
A	-18	GLY	-	cloning artifact	UNP P21054
A	-17	SER	-	cloning artifact	UNP P21054
A	-16	SER	-	cloning artifact	UNP P21054
A	-15	HIS	-	expression tag	UNP P21054
A	-14	HIS	-	expression tag	UNP P21054
A	-13	HIS	-	expression tag	UNP P21054
A	-12	HIS	-	expression tag	UNP P21054
A	-11	HIS	-	expression tag	UNP P21054
A	-10	HIS	-	expression tag	UNP P21054
A	-9	SER	-	cloning artifact	UNP P21054
A	-8	SER	-	cloning artifact	UNP P21054
A	-7	GLY	-	cloning artifact	UNP P21054
A	-6	LEU	=	cloning artifact	UNP P21054
A	-5	VAL	=	cloning artifact	UNP P21054
A	-4	PRO	=	cloning artifact	UNP P21054
A	-3	ARG	=	cloning artifact	UNP P21054



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	-2	GLY	-	cloning artifact	UNP P21054
A	-1	SER	-	cloning artifact	UNP P21054
A	0	HIS	-	cloning artifact	UNP P21054
В	-19	MET	-	cloning artifact	UNP P21054
В	-18	GLY	-	cloning artifact	UNP P21054
В	-17	SER	-	cloning artifact	UNP P21054
В	-16	SER	-	cloning artifact	UNP P21054
В	-15	HIS	-	expression tag	UNP P21054
В	-14	HIS	-	expression tag	UNP P21054
В	-13	HIS	-	expression tag	UNP P21054
В	-12	HIS	-	expression tag	UNP P21054
В	-11	HIS	-	expression tag	UNP P21054
В	-10	HIS	-	expression tag	UNP P21054
В	-9	SER	_	cloning artifact	UNP P21054
В	-8	SER	-	cloning artifact	UNP P21054
В	-7	GLY	_	cloning artifact	UNP P21054
В	-6	LEU	-	cloning artifact	UNP P21054
В	-5	VAL	-	cloning artifact	UNP P21054
В	-4	PRO	-	cloning artifact	UNP P21054
В	-3	ARG	-	cloning artifact	UNP P21054
В	-2	GLY	-	cloning artifact	UNP P21054
В	-1	SER	-	cloning artifact	UNP P21054
В	0	HIS	-	cloning artifact	UNP P21054
С	-19	MET	-	cloning artifact	UNP P21054
С	-18	GLY	-	cloning artifact	UNP P21054
С	-17	SER	-	cloning artifact	UNP P21054
С	-16	SER	-	cloning artifact	UNP P21054
С	-15	HIS	-	expression tag	UNP P21054
С	-14	HIS	-	expression tag	UNP P21054
С	-13	HIS	-	expression tag	UNP P21054
С	-12	HIS	-	expression tag	UNP P21054
С	-11	HIS	-	expression tag	UNP P21054
С	-10	HIS	-	expression tag	UNP P21054
С	-9	SER	-	cloning artifact	UNP P21054
С	-8	SER	-	cloning artifact	UNP P21054
С	-7	GLY	-	cloning artifact	UNP P21054
С	-6	LEU	-	cloning artifact	UNP P21054
С	-5	VAL		cloning artifact	UNP P21054
С	-4	PRO		cloning artifact	UNP P21054
С	-3	ARG		cloning artifact	UNP P21054
С	-2	GLY	-	cloning artifact	UNP P21054
С	-1	SER	-	cloning artifact	UNP P21054



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
С	0	HIS	-	cloning artifact	UNP P21054
D	-19	MET	-	cloning artifact	UNP P21054
D	-18	GLY	-	cloning artifact	UNP P21054
D	-17	SER	-	cloning artifact	UNP P21054
D	-16	SER	-	cloning artifact	UNP P21054
D	-15	HIS	-	expression tag	UNP P21054
D	-14	HIS	-	expression tag	UNP P21054
D	-13	HIS	-	expression tag	UNP P21054
D	-12	HIS	-	expression tag	UNP P21054
D	-11	HIS	-	expression tag	UNP P21054
D	-10	HIS	-	expression tag	UNP P21054
D	-9	SER	-	cloning artifact	UNP P21054
D	-8	SER	-	cloning artifact	UNP P21054
D	-7	GLY	-	cloning artifact	UNP P21054
D	-6	LEU	-	cloning artifact	UNP P21054
D	-5	VAL	-	cloning artifact	UNP P21054
D	-4	PRO	-	cloning artifact	UNP P21054
D	-3	ARG	-	cloning artifact	UNP P21054
D	-2	GLY	-	cloning artifact	UNP P21054
D	-1	SER	-	cloning artifact	UNP P21054
D	0	HIS	-	cloning artifact	UNP P21054
Е	-19	MET	_	cloning artifact	UNP P21054
Е	-18	GLY	-	cloning artifact	UNP P21054
Е	-17	SER	-	cloning artifact	UNP P21054
Е	-16	SER	_	cloning artifact	UNP P21054
Е	-15	HIS	-	expression tag	UNP P21054
Е	-14	HIS	-	expression tag	UNP P21054
Е	-13	HIS	-	expression tag	UNP P21054
Е	-12	HIS	-	expression tag	UNP P21054
Е	-11	HIS	-	expression tag	UNP P21054
Е	-10	HIS	-	expression tag	UNP P21054
Е	-9	SER	-	cloning artifact	UNP P21054
Е	-8	SER	-	cloning artifact	UNP P21054
Е	-7	GLY	-	cloning artifact	UNP P21054
Е	-6	LEU	-	cloning artifact	UNP P21054
Е	-5	VAL	-	cloning artifact	UNP P21054
Е	-4	PRO	-	cloning artifact	UNP P21054
Е	-3	ARG	-	cloning artifact	UNP P21054
Е	-2	GLY	-	cloning artifact	UNP P21054
Е	-1	SER	-	cloning artifact	UNP P21054
Е	0	HIS	-	cloning artifact	UNP P21054
F	-19	MET	-	cloning artifact	UNP P21054



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-18	GLY	-	cloning artifact	UNP P21054
F	-17	SER	-	cloning artifact	UNP P21054
F	-16	SER	-	cloning artifact	UNP P21054
F	-15	HIS	-	expression tag	UNP P21054
F	-14	HIS	-	expression tag	UNP P21054
F	-13	HIS	-	expression tag	UNP P21054
F	-12	HIS	-	expression tag	UNP P21054
F	-11	HIS	-	expression tag	UNP P21054
F	-10	HIS	-	expression tag	UNP P21054
F	-9	SER	-	cloning artifact	UNP P21054
F	-8	SER	-	cloning artifact	UNP P21054
F	-7	GLY	-	cloning artifact	UNP P21054
F	-6	LEU	-	cloning artifact	UNP P21054
F	-5	VAL	-	cloning artifact	UNP P21054
F	-4	PRO	-	cloning artifact	UNP P21054
F	-3	ARG	-	cloning artifact	UNP P21054
F	-2	GLY	-	cloning artifact	UNP P21054
F	-1	SER	-	cloning artifact	UNP P21054
F	0	HIS	-	cloning artifact	UNP P21054

 \bullet Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 8 6 2	0	0



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	Е	1	Total 8	C 6	O 2	0	0


• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	57	Total O 57 57	0	0
3	В	43	Total O 43 43	0	0
3	С	53	Total O 53 53	0	0
3	D	56	Total O 56 56	0	0
3	Е	41	Total O 41 41	0	0
3	F	32	Total O 32 32	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	68.64Å 109.99Å 69.58Å	Depositor
a, b, c, α , β , γ	90.00° 110.92° 90.00°	Depositor
Resolution (Å)	30.00 - 2.20	Depositor
resolution (A)	29.13 - 2.19	EDS
% Data completeness	96.1 (30.00-2.20)	Depositor
(in resolution range)	95.0 (29.13-2.19)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$< I/\sigma(I) > 1$	1.96 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.202 , 0.247	Depositor
it, it _{free}	0.195 , 0.239	DCC
R_{free} test set	2397 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.32 \; , 36.8$	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6177	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	Bond lengths		angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.34	0/989	0.57	0/1333
1	В	0.32	0/998	0.54	0/1345
1	С	0.36	0/998	0.55	0/1345
1	D	0.33	0/998	0.54	0/1345
1	Е	0.36	0/993	0.52	0/1338
1	F	0.34	0/993	0.55	0/1338
All	All	0.34	0/5969	0.54	0/8044

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	974	0	968	18	0
1	В	983	0	973	24	0
1	С	983	0	973	14	0
1	D	983	0	973	18	0
1	Е	978	0	971	30	0
1	F	978	0	971	14	0
2	A	8	0	14	0	0



I 'ontanued	trom	mmonia	maaa
Continued	110111	mea_{mons}	DU_0U_0
00,000,000	.,	p. 0000 ao	pagain

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
2	Ε	8	0	13	0	0
3	A	57	0	0	1	0
3	В	43	0	0	1	0
3	С	53	0	0	2	0
3	D	56	0	0	2	0
3	Ε	41	0	0	2	0
3	F	32	0	0	0	0
All	All	6177	0	5856	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 108 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:72:MET:HE2	1:A:98:LEU:HD12	1.34	1.08
1:A:66:ILE:HD12	1:A:108:ASP:HB3	1.59	0.83
1:B:90:ARG:HD2	3:B:148:HOH:O	1.87	0.73
1:E:92:GLU:OE2	1:F:2:ARG:NE	2.23	0.71
1:E:17:GLN:HE22	1:F:22:ASP:H	1.39	0.69

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	115/137 (84%)	114 (99%)	1 (1%)	0	100	100
1	В	117/137 (85%)	116 (99%)	1 (1%)	0	100	100
1	С	117/137 (85%)	116 (99%)	1 (1%)	0	100	100
1	D	117/137 (85%)	117 (100%)	0	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	E	116/137 (85%)	114 (98%)	2 (2%)	0	100	100
1	F	116/137 (85%)	115 (99%)	1 (1%)	0	100	100
All	All	698/822 (85%)	692 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percenti	les
1	A	109/126~(86%)	103 (94%)	6 (6%)	21 26	3
1	В	109/126 (86%)	101 (93%)	8 (7%)	14 15	5
1	С	109/126 (86%)	101 (93%)	8 (7%)	14 15	5
1	D	109/126 (86%)	105 (96%)	4 (4%)	34 43	}
1	E	109/126 (86%)	105 (96%)	4 (4%)	34 43	}
1	F	109/126 (86%)	108 (99%)	1 (1%)	78 88	}
All	All	654/756~(86%)	623 (95%)	31 (5%)	26 33	}

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	4	LEU
1	Ε	48	MET
1	С	38	ASP
1	Ε	65	ASN
1	D	55	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

\mathbf{Mol}	Chain	Res	Type
1	F	13	ASN
1	F	21	ASN



Mol	Chain	Res	Type
1	F	17	GLN
1	D	13	ASN
1	Е	65	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain Res	Link	B	Bond lengths			Bond angles		
	Type		nes	Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2									
2	MPD	Е	502	-	7,7,7	1.45	2 (28%)	9,10,10	1.48	2 (22%)									
2	MPD	A	501	-	7,7,7	1.38	2 (28%)	9,10,10	1.43	2 (22%)									

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	E	502	-	1/1/2/2	0/5/5/5	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	A	501	-	1/1/2/2	0/5/5/5	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	Ε	502	MPD	C1-C2	2.53	1.60	1.52
2	A	501	MPD	C1-C2	2.52	1.60	1.52
2	Е	502	MPD	O2-C2	-2.25	1.39	1.44
2	A	501	MPD	O2-C2	-2.11	1.39	1.44

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	Е	502	MPD	O2-C2-C1	-2.84	98.98	108.08
2	A	501	MPD	O2-C2-C1	-2.58	99.79	108.08
2	A	501	MPD	O2-C2-C3	2.21	118.09	109.80
2	Е	502	MPD	O2-C2-C3	2.16	117.89	109.80

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	501	MPD	C4
2	Е	502	MPD	C4

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	117/137 (85%)	0.01	3 (2%) 56 53	23, 38, 60, 96	0
1	В	119/137 (86%)	0.09	3 (2%) 57 55	26, 43, 64, 77	0
1	С	119/137 (86%)	-0.14	4 (3%) 45 43	25, 37, 54, 74	0
1	D	119/137 (86%)	0.08	5 (4%) 36 34	26, 40, 60, 70	0
1	E	118/137 (86%)	-0.15	2 (1%) 70 68	23, 38, 60, 89	0
1	F	118/137 (86%)	-0.08	1 (0%) 86 85	23, 41, 62, 75	0
All	All	710/822 (86%)	-0.03	18 (2%) 57 55	23, 40, 63, 96	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	115	GLY	5.7
1	F	115	GLY	4.1
1	С	115	GLY	3.9
1	Е	115	GLY	3.9
1	D	115	GLY	3.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	MPD	A	501	8/8	0.58	0.40	82,83,84,84	0
2	MPD	Е	502	8/8	0.88	0.20	48,50,53,54	0

6.5 Other polymers (i)

There are no such residues in this entry.

