



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 30, 2021 – 05:36 PM EST

PDB ID : 3I3V  
Title : Crystal Structure of probable secreted solute-binding lipoprotein from *Streptomyces coelicolor*  
Authors : Damodharan, L.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2009-07-01  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	
1	B	405	
1	C	405	
1	D	405	

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 12244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable secreted solute-binding lipoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	391	2980	1883	528	564	5	0	0	0
1	B	390	2964	1875	527	557	5	0	0	0
1	C	394	2993	1890	528	570	5	0	0	0
1	D	391	2978	1882	528	563	5	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	MSE	-	insertion	UNP Q9RL34
A	34	SER	-	insertion	UNP Q9RL34
A	35	LEU	-	insertion	UNP Q9RL34
A	430	GLU	-	insertion	UNP Q9RL34
A	431	GLY	-	insertion	UNP Q9RL34
A	432	HIS	-	insertion	UNP Q9RL34
A	433	HIS	-	insertion	UNP Q9RL34
A	434	HIS	-	insertion	UNP Q9RL34
A	435	HIS	-	insertion	UNP Q9RL34
A	436	HIS	-	insertion	UNP Q9RL34
A	437	HIS	-	insertion	UNP Q9RL34
B	33	MSE	-	insertion	UNP Q9RL34
B	34	SER	-	insertion	UNP Q9RL34
B	35	LEU	-	insertion	UNP Q9RL34
B	430	GLU	-	insertion	UNP Q9RL34
B	431	GLY	-	insertion	UNP Q9RL34
B	432	HIS	-	insertion	UNP Q9RL34
B	433	HIS	-	insertion	UNP Q9RL34
B	434	HIS	-	insertion	UNP Q9RL34
B	435	HIS	-	insertion	UNP Q9RL34
B	436	HIS	-	insertion	UNP Q9RL34

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	437	HIS	-	insertion	UNP Q9RL34
C	33	MSE	-	insertion	UNP Q9RL34
C	34	SER	-	insertion	UNP Q9RL34
C	35	LEU	-	insertion	UNP Q9RL34
C	430	GLU	-	insertion	UNP Q9RL34
C	431	GLY	-	insertion	UNP Q9RL34
C	432	HIS	-	insertion	UNP Q9RL34
C	433	HIS	-	insertion	UNP Q9RL34
C	434	HIS	-	insertion	UNP Q9RL34
C	435	HIS	-	insertion	UNP Q9RL34
C	436	HIS	-	insertion	UNP Q9RL34
C	437	HIS	-	insertion	UNP Q9RL34
D	33	MSE	-	insertion	UNP Q9RL34
D	34	SER	-	insertion	UNP Q9RL34
D	35	LEU	-	insertion	UNP Q9RL34
D	430	GLU	-	insertion	UNP Q9RL34
D	431	GLY	-	insertion	UNP Q9RL34
D	432	HIS	-	insertion	UNP Q9RL34
D	433	HIS	-	insertion	UNP Q9RL34
D	434	HIS	-	insertion	UNP Q9RL34
D	435	HIS	-	insertion	UNP Q9RL34
D	436	HIS	-	insertion	UNP Q9RL34
D	437	HIS	-	insertion	UNP Q9RL34


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	99	Total O 99 99	0	0
2	B	76	Total O 76 76	0	0
2	C	85	Total O 85 85	0	0
2	D	69	Total O 69 69	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable secreted solute-binding lipoprotein




- Molecule 1: Probable secreted solute-binding lipoprotein




- Molecule 1: Probable secreted solute-binding lipoprotein





● Molecule 1: Probable secreted solute-binding lipoprotein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.31Å 91.31Å 370.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.70 – 2.30 45.31 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.0 (36.70-2.30) 98.1 (45.31-2.30)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.71 (at 2.29Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.220 , 0.259 0.226 , 0.267	Depositor DCC
$R_{free}$ test set	3997 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.2	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 36.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12244	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5812e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3040	0.67	0/4138
1	B	0.34	0/3023	0.67	1/4116 (0.0%)
1	C	0.34	0/3053	0.65	1/4158 (0.0%)
1	D	0.36	0/3037	0.67	0/4134
All	All	0.35	0/12153	0.66	2/16546 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	388	LEU	N-CA-C	-5.69	95.64	111.00
1	C	37	SER	N-CA-C	5.54	125.95	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	330	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2980	0	2949	82	0
1	B	2964	0	2938	118	0
1	C	2993	0	2948	121	0
1	D	2978	0	2951	119	0
2	A	99	0	0	5	0
2	B	76	0	0	3	0
2	C	85	0	0	7	0
2	D	69	0	0	4	0
All	All	12244	0	11786	436	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 436 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:357:ARG:HB3	1:D:359:MSE:HE3	1.20	1.14
1:D:125:ASP:O	1:D:126:ASP:HB3	1.51	1.09
1:D:399:GLN:HG2	1:D:400:PRO:HD3	1.34	1.02
1:B:357:ARG:HB3	1:B:359:MSE:HE3	1.42	1.01
1:A:357:ARG:HB2	1:A:359:MSE:CE	1.96	0.96

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	389/405 (96%)	371 (95%)	15 (4%)	3 (1%)	19 23
1	B	388/405 (96%)	361 (93%)	20 (5%)	7 (2%)	8 7

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	392/405 (97%)	369 (94%)	18 (5%)	5 (1%)	12	12
1	D	389/405 (96%)	366 (94%)	17 (4%)	6 (2%)	10	10
All	All	1558/1620 (96%)	1467 (94%)	70 (4%)	21 (1%)	12	12

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	ASP
1	A	196	PRO
1	B	180	ASP
1	C	127	TRP
1	C	310	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/314 (97%)	278 (91%)	27 (9%)	9	11
1	B	302/314 (96%)	268 (89%)	34 (11%)	6	6
1	C	306/314 (98%)	263 (86%)	43 (14%)	3	3
1	D	305/314 (97%)	276 (90%)	29 (10%)	8	10
All	All	1218/1256 (97%)	1085 (89%)	133 (11%)	6	7

5 of 133 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	422	LEU
1	C	129	LEU
1	D	336	GLU
1	C	36	ASP
1	C	102	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	85	GLN
1	C	207	GLN
1	D	308	HIS
1	C	163	ASN
1	A	399	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	386/405 (95%)	-0.23	0 <a href="#">100</a> <a href="#">100</a>	21, 34, 48, 65	0
1	B	385/405 (95%)	-0.02	6 (1%) <a href="#">72</a> <a href="#">77</a>	24, 40, 58, 66	0
1	C	389/405 (96%)	-0.06	5 (1%) <a href="#">77</a> <a href="#">81</a>	25, 37, 55, 66	0
1	D	386/405 (95%)	-0.07	3 (0%) <a href="#">86</a> <a href="#">89</a>	23, 37, 54, 67	0
All	All	1546/1620 (95%)	-0.09	14 (0%) <a href="#">84</a> <a href="#">88</a>	21, 37, 55, 67	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	180	ASP	7.0
1	C	310	GLY	4.3
1	D	310	GLY	4.0
1	D	169	ARG	2.6
1	B	307	ALA	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.