



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 12:58 am BST

PDB ID : 2I9D  
Title : chloramphenicol acetyltransferase  
Authors : Duke, N.E.C.; Li, H.; Abdullah, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2006-09-05  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 5133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chloramphenicol acetyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	214	Total 1749	C 1123	N 297	O 324	S 1	Se 4	0	0	0
1	B	193	Total 1584	C 1021	N 272	O 286	S 1	Se 4	0	0	0
1	C	193	Total 1584	C 1021	N 272	O 286	S 1	Se 4	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	CLONING ARTIFACT	UNP Q8A336
A	-1	ASN	-	CLONING ARTIFACT	UNP Q8A336
A	0	ALA	-	CLONING ARTIFACT	UNP Q8A336
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
A	85	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
A	186	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
A	190	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
B	-2	SER	-	CLONING ARTIFACT	UNP Q8A336
B	-1	ASN	-	CLONING ARTIFACT	UNP Q8A336
B	0	ALA	-	CLONING ARTIFACT	UNP Q8A336
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
B	85	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
B	186	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
B	190	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
C	-2	SER	-	CLONING ARTIFACT	UNP Q8A336
C	-1	ASN	-	CLONING ARTIFACT	UNP Q8A336
C	0	ALA	-	CLONING ARTIFACT	UNP Q8A336
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
C	85	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
C	186	MSE	MET	MODIFIED RESIDUE	UNP Q8A336
C	190	MSE	MET	MODIFIED RESIDUE	UNP Q8A336

- Molecule 2 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	A	77	Total 77	O 77	0	0
2	B	71	Total 71	O 71	0	0
2	C	68	Total 68	O 68	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.14Å 116.14Å 130.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.44 – 2.30 43.43 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.44-2.30) 99.8 (43.43-2.30)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.17 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.188 , 0.230 0.189 , 0.230	Depositor DCC
$R_{free}$ test set	2264 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 40.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.036 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5133	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.71	0/1791	0.72	2/2415 (0.1%)
1	B	0.74	0/1620	0.71	1/2179 (0.0%)
1	C	0.78	0/1620	0.76	2/2179 (0.1%)
All	All	0.74	0/5031	0.73	5/6773 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	27	LEU	CA-CB-CG	9.15	136.34	115.30
1	A	27	LEU	CB-CG-CD2	-6.23	100.41	111.00
1	B	27	LEU	CA-CB-CG	6.17	129.48	115.30
1	C	2	LYS	N-CA-C	5.62	126.17	111.00
1	C	27	LEU	CA-CB-CG	5.30	127.48	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	213	LEU	Peptide
1	B	213	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1749	0	1696	24	0
1	B	1584	0	1542	26	1
1	C	1584	0	1542	33	1
2	A	77	0	0	1	0
2	B	71	0	0	0	0
2	C	68	0	0	4	0
All	All	5133	0	4780	72	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:ARG:HH11	1:B:39:ARG:HG2	1.23	1.00
1:B:41:ARG:HH12	1:B:214:LYS:HG2	1.31	0.95
1:B:180:ARG:HH22	1:C:180:ARG:NH1	1.67	0.91
1:B:180:ARG:HH22	1:C:180:ARG:HH11	1.26	0.84
1:B:39:ARG:HG2	1:B:39:ARG:NH1	1.89	0.84
1:C:172:LEU:HG	1:C:190:MSE:HB3	1.61	0.81
1:B:39:ARG:HH11	1:B:39:ARG:CG	1.99	0.75
1:A:128:PRO:HD2	1:B:14:GLU:HG3	1.70	0.73
1:C:176:LYS:HG2	1:C:178:ILE:HD12	1.75	0.67
1:C:26:GLN:NE2	1:C:167:ASN:H	1.94	0.66
1:B:180:ARG:NH2	1:C:180:ARG:HH11	1.95	0.64
1:C:26:GLN:HE22	1:C:167:ASN:H	1.46	0.63
1:C:89:ILE:HG12	1:C:100:THR:HG23	1.81	0.62
1:B:41:ARG:HH22	1:B:214:LYS:HD3	1.69	0.58
1:C:32:GLU:HG3	2:C:270:HOH:O	2.02	0.58
1:C:158:THR:HG22	2:C:217:HOH:O	2.03	0.58
1:A:158:THR:HG22	1:C:158:THR:HG21	1.86	0.57
1:B:158:THR:HG21	1:C:158:THR:HG21	1.87	0.56
1:B:83:ILE:CD1	1:B:142:GLY:HA2	2.35	0.56
1:A:5:ILE:HD11	1:A:79:LEU:HB2	1.89	0.55
1:C:112:TYR:CE2	1:C:116:ARG:HG3	2.43	0.53

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:112:TYR:CE2	1:B:116:ARG:HG3	2.45	0.52
1:C:20:ARG:HG3	1:C:21:HIS:CD2	2.45	0.52
1:B:83:ILE:HD12	1:B:142:GLY:C	2.30	0.51
1:A:158:THR:CG2	1:C:158:THR:HG21	2.41	0.51
1:B:98:PHE:O	1:B:100:THR:HG22	2.10	0.51
1:A:134:GLU:HG3	2:A:248:HOH:O	2.11	0.51
1:C:68:ARG:HG2	1:C:83:ILE:HG13	1.93	0.50
1:C:68:ARG:NH2	2:C:275:HOH:O	2.43	0.50
1:A:68:ARG:HG2	1:A:83:ILE:CG1	2.42	0.49
1:A:178:ILE:HG22	1:A:180:ARG:HD2	1.93	0.49
1:A:128:PRO:HD2	1:B:14:GLU:CG	2.40	0.49
1:C:4:ILE:HD13	1:C:78:VAL:HG12	1.94	0.49
1:B:56:ALA:HB2	1:B:212:PHE:CD2	2.48	0.49
1:C:15:ASN:ND2	1:C:15:ASN:H	2.11	0.49
1:B:53:TYR:HA	1:B:213:LEU:HD21	1.95	0.48
1:A:4:ILE:N	1:A:4:ILE:HD12	2.29	0.48
1:B:4:ILE:HD13	1:B:78:VAL:HG12	1.96	0.48
1:C:-2:SER:OG	1:C:-1:ASN:N	2.47	0.48
1:C:1:MSE:C	1:C:1:MSE:HE3	2.33	0.48
1:B:50:PHE:HA	1:B:53:TYR:CE2	2.50	0.47
1:C:68:ARG:HG2	1:C:83:ILE:CG1	2.45	0.47
1:C:172:LEU:HD11	1:C:190:MSE:HE3	1.97	0.46
1:A:5:ILE:HD13	1:A:69:TYR:CE1	2.51	0.45
1:C:176:LYS:HG2	1:C:178:ILE:CD1	2.42	0.45
1:A:68:ARG:HD2	1:A:80:TYR:O	2.16	0.45
1:C:153:TYR:HB2	1:C:178:ILE:HD13	1.98	0.45
1:C:173:ASN:ND2	2:C:263:HOH:O	2.47	0.45
1:B:207:LYS:HD3	1:B:207:LYS:HA	1.59	0.45
1:A:88:PRO:HD2	1:A:146:LEU:O	2.17	0.45
1:B:213:LEU:O	1:B:214:LYS:HB2	2.18	0.44
1:A:15:ASN:H	1:A:15:ASN:HD22	1.66	0.44
1:B:12:ARG:HA	1:B:15:ASN:HD22	1.82	0.44
1:C:39:ARG:HD2	1:C:184:LEU:HD22	1.99	0.44
1:A:127:ASP:HB3	1:A:130:ALA:HB2	2.00	0.44
1:A:68:ARG:HG2	1:A:83:ILE:HG12	1.98	0.44
1:C:15:ASN:HD22	1:C:15:ASN:H	1.65	0.43
1:A:153:TYR:O	1:B:31:SER:OG	2.29	0.43
1:B:27:LEU:HG	1:B:192:ILE:HD11	2.01	0.43
1:A:6:ASP:OD2	1:A:9:ASN:ND2	2.52	0.43
1:B:-2:SER:HA	1:B:-1:ASN:HA	1.74	0.43
1:A:131:ALA:O	1:A:135:GLU:HG2	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:SER:HB2	1:C:158:THR:HG23	2.02	0.42
1:A:15:ASN:ND2	1:A:15:ASN:H	2.17	0.42
1:A:22:PHE:CE1	1:C:162:GLU:HG3	2.54	0.41
1:A:69:TYR:CE2	1:A:79:LEU:HD13	2.55	0.41
1:C:68:ARG:HD2	1:C:80:TYR:O	2.21	0.41
1:A:211:ASP:O	1:A:214:LYS:HG2	2.20	0.41
1:A:36:GLY:O	1:A:40:GLN:NE2	2.54	0.41
1:C:83:ILE:HA	1:C:83:ILE:HD13	1.85	0.40
1:C:190:MSE:HE2	1:C:192:ILE:CG2	2.52	0.40
1:B:150:PRO:O	1:B:176:LYS:HG3	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2:LYS:O	1:C:1:MSE:O[5_554]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/217 (98%)	207 (98%)	3 (1%)	2 (1%)	17	20
1	B	187/217 (86%)	184 (98%)	3 (2%)	0	100	100
1	C	187/217 (86%)	185 (99%)	0	2 (1%)	14	15
All	All	586/651 (90%)	576 (98%)	6 (1%)	4 (1%)	22	26

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	2	LYS
1	C	1	MSE

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	9	ASN
1	A	10	TRP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/182 (101%)	174 (95%)	10 (5%)	22	30
1	B	167/182 (92%)	155 (93%)	12 (7%)	14	18
1	C	167/182 (92%)	157 (94%)	10 (6%)	19	26
All	All	518/546 (95%)	486 (94%)	32 (6%)	18	25

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	27	LEU
1	A	39	ARG
1	A	40	GLN
1	A	114	GLU
1	A	116	ARG
1	A	138	ASP
1	A	178	ILE
1	A	180	ARG
1	A	203	SER
1	B	20	ARG
1	B	27	LEU
1	B	39	ARG
1	B	40	GLN
1	B	82	THR
1	B	83	ILE
1	B	100	THR
1	B	116	ARG
1	B	178	ILE
1	B	204	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	207	LYS
1	B	214	LYS
1	C	2	LYS
1	C	11	GLU
1	C	27	LEU
1	C	82	THR
1	C	100	THR
1	C	116	ARG
1	C	158	THR
1	C	178	ILE
1	C	190	MSE
1	C	194	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	40	GLN
1	A	47	GLN
1	A	105	HIS
1	B	15	ASN
1	B	47	GLN
1	B	106	ASN
1	C	15	ASN
1	C	21	HIS
1	C	26	GLN
1	C	47	GLN
1	C	173	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	210/217 (96%)	0.04	6 (2%) 51 58	20, 33, 49, 57	0
1	B	189/217 (87%)	0.04	8 (4%) 36 43	17, 29, 51, 60	0
1	C	189/217 (87%)	-0.13	7 (3%) 41 48	20, 28, 44, 51	0
All	All	588/651 (90%)	-0.02	21 (3%) 42 49	17, 30, 49, 60	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	98	PHE	4.1
1	C	20	ARG	3.6
1	B	117	LEU	3.5
1	C	21	HIS	3.3
1	A	106	ASN	3.0
1	A	9	ASN	3.0
1	A	157	ILE	2.9
1	C	98	PHE	2.8
1	C	9	ASN	2.6
1	B	214	LYS	2.5
1	C	113	GLN	2.5
1	A	43	LYS	2.4
1	B	40	GLN	2.4
1	C	158	THR	2.2
1	B	120	ASP	2.1
1	B	158	THR	2.1
1	C	74	ASP	2.0
1	B	192	ILE	2.0
1	A	113	GLN	2.0
1	A	158	THR	2.0
1	B	113	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.