



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 3, 2021 – 12:05 AM EDT

PDB ID : 3IBB
Title : Propionyl-CoA Carboxylase Beta Subunit, D422A
Authors : Diacovich, L.; Arabolaza, A.; Shillito, E.M.; Lin, T.-W.; Mitchell, D.L.; Pham, H.; Melgar, M.M.
Deposited on : 2009-07-15
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	530	22% (green) 66% (yellow) 10% (orange) . (grey)
1	B	530	25% (green) 61% (yellow) 12% (orange) . (grey)
1	C	530	27% (green) 62% (yellow) 9% (orange) . (grey)
1	D	530	23% (green) 63% (yellow) 12% (orange) . (grey)
1	E	530	24% (green) 64% (yellow) 11% (orange) . (grey)
1	F	530	25% (green) 59% (yellow) 13% (orange) . (grey)

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 23700 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Propionyl-CoA carboxylase complex B subunit.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	521	3950	2480	698	759	13	0	0	0
1	B	521	3950	2480	698	759	13	0	0	0
1	C	521	3950	2480	698	759	13	0	0	0
1	D	521	3950	2480	698	759	13	0	0	0
1	E	521	3950	2480	698	759	13	0	0	0
1	F	521	3950	2480	698	759	13	0	0	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	422	ALA	ASP	engineered mutation	UNP Q9X4K7
B	422	ALA	ASP	engineered mutation	UNP Q9X4K7
C	422	ALA	ASP	engineered mutation	UNP Q9X4K7
D	422	ALA	ASP	engineered mutation	UNP Q9X4K7
E	422	ALA	ASP	engineered mutation	UNP Q9X4K7
F	422	ALA	ASP	engineered mutation	UNP Q9X4K7




● Molecule 1: Propionyl-CoA carboxylase complex B subunit





• Molecule 1: Propionyl-CoA carboxylase complex B subunit

Chain F: 25% 59% 13%



L513	R514	R515	R516	R517	L520	P521	P522	K523	K524	H525	G526	M527	L528	P529	L530
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.06Å 183.33Å 228.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.50 45.83 – 3.46	Depositor EDS
% Data completeness (in resolution range)	94.4 (50.00-3.50) 94.5 (45.83-3.46)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.04 (at 3.48Å)	Xtrriage
Refinement program	REFMAC 5.2.0019, CNS	Depositor
R, R_{free}	0.190 , 0.269 0.298 , 0.309	Depositor DCC
R_{free} test set	2294 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	65.3	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 23.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	23700	wwPDB-VP
Average B, all atoms (Å ²)	2.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.45	1/4030 (0.0%)	0.82	5/5474 (0.1%)
1	B	0.39	1/4030 (0.0%)	0.77	2/5474 (0.0%)
1	C	0.37	1/4030 (0.0%)	0.78	2/5474 (0.0%)
1	D	0.41	1/4030 (0.0%)	0.79	5/5474 (0.1%)
1	E	0.39	1/4030 (0.0%)	0.78	5/5474 (0.1%)
1	F	0.37	0/4030	0.74	3/5474 (0.1%)
All	All	0.40	5/24180 (0.0%)	0.78	22/32844 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	530	LEU	C-OXT	5.86	1.34	1.23
1	A	238	HIS	C-N	-5.53	1.21	1.34
1	E	414	ARG	C-N	5.39	1.46	1.34
1	D	363	CYS	C-N	5.35	1.46	1.34
1	B	476	GLU	C-N	-5.09	1.22	1.34

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	E	108	VAL	CB-CA-C	-5.97	100.06	111.40
1	B	150	VAL	CB-CA-C	-5.94	100.12	111.40
1	D	277	VAL	N-CA-C	-5.83	95.27	111.00
1	D	513	LEU	CB-CA-C	-5.78	99.22	110.20
1	A	278	THR	CB-CA-C	5.72	127.04	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3950	0	3880	836	0
1	B	3950	0	3880	780	0
1	C	3950	0	3880	784	0
1	D	3950	0	3881	807	0
1	E	3950	0	3881	772	0
1	F	3950	0	3881	786	0
All	All	23700	0	23283	4332	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 92.

The worst 5 of 4332 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:497:ILE:HD13	1:F:505:HIS:CD2	1.40	1.54
1:C:379:PHE:CE2	1:D:205:ILE:HD11	1.38	1.54
1:A:318:PHE:CZ	1:A:351:THR:HB	1.42	1.53
1:F:311:PHE:CE1	1:F:325:GLY:HA3	1.45	1.49
1:F:215:THR:CG2	1:F:217:GLU:HG2	1.43	1.47

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	519/530 (98%)	503 (97%)	14 (3%)	2 (0%)	34 72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	519/530 (98%)	501 (96%)	18 (4%)	0	100	100
1	C	519/530 (98%)	501 (96%)	17 (3%)	1 (0%)	47	81
1	D	519/530 (98%)	501 (96%)	17 (3%)	1 (0%)	47	81
1	E	519/530 (98%)	505 (97%)	14 (3%)	0	100	100
1	F	519/530 (98%)	503 (97%)	13 (2%)	3 (1%)	25	64
All	All	3114/3180 (98%)	3014 (97%)	93 (3%)	7 (0%)	47	81

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	416	ALA
1	F	145	ARG
1	F	516	LYS
1	A	389	GLY
1	F	381	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/420 (98%)	349 (85%)	62 (15%)	3	17
1	B	411/420 (98%)	339 (82%)	72 (18%)	2	10
1	C	411/420 (98%)	353 (86%)	58 (14%)	3	19
1	D	411/420 (98%)	337 (82%)	74 (18%)	1	9
1	E	411/420 (98%)	342 (83%)	69 (17%)	2	12
1	F	411/420 (98%)	325 (79%)	86 (21%)	1	6
All	All	2466/2520 (98%)	2045 (83%)	421 (17%)	2	12

5 of 421 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	317	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	196	MET
1	F	421	TYR
1	D	370	VAL
1	D	508	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 92 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	199	GLN
1	E	483	ASN
1	E	231	ASN
1	E	315	GLN
1	F	70	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.