



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 03:34 am BST

PDB ID : 1ICV  
Title : THE STRUCTURE OF ESCHERICHIA COLI NITROREDUCTASE COM-  
PLEXED WITH NICOTINIC ACID  
Authors : Lovering, A.L.; Hyde, E.I.; Searle, P.F.; White, S.A.  
Deposited on : 2001-04-02  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	217	75% (green), 19% (yellow), 5% (orange)
1	B	217	71% (green), 24% (yellow), 5% (orange)
1	C	217	77% (green), 20% (yellow), 5% (orange)
1	D	217	72% (green), 23% (yellow), 5% (orange)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FMN	B	703	X	X	-	-

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
2	FMN	C	705	X	X	-	-
2	FMN	D	707	X	X	-	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7511 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called OXYGEN-INSENSITIVE NAD(P)H NITROREDUCTASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	216	1677	1065	287	320	1	4	0	0	0
1	B	216	1677	1065	287	320	1	4	0	0	0
1	C	216	1677	1065	287	320	1	4	0	0	0
1	D	216	1677	1065	287	320	1	4	0	0	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P38489
A	75	MSE	MET	MODIFIED RESIDUE	UNP P38489
A	90	MSE	MET	MODIFIED RESIDUE	UNP P38489
A	127	MSE	MET	MODIFIED RESIDUE	UNP P38489
A	139	MSE	MET	MODIFIED RESIDUE	UNP P38489
B	1	MSE	MET	MODIFIED RESIDUE	UNP P38489
B	75	MSE	MET	MODIFIED RESIDUE	UNP P38489
B	90	MSE	MET	MODIFIED RESIDUE	UNP P38489
B	127	MSE	MET	MODIFIED RESIDUE	UNP P38489
B	139	MSE	MET	MODIFIED RESIDUE	UNP P38489
C	1	MSE	MET	MODIFIED RESIDUE	UNP P38489
C	75	MSE	MET	MODIFIED RESIDUE	UNP P38489
C	90	MSE	MET	MODIFIED RESIDUE	UNP P38489
C	127	MSE	MET	MODIFIED RESIDUE	UNP P38489
C	139	MSE	MET	MODIFIED RESIDUE	UNP P38489
D	1	MSE	MET	MODIFIED RESIDUE	UNP P38489
D	75	MSE	MET	MODIFIED RESIDUE	UNP P38489
D	90	MSE	MET	MODIFIED RESIDUE	UNP P38489
D	127	MSE	MET	MODIFIED RESIDUE	UNP P38489
D	139	MSE	MET	MODIFIED RESIDUE	UNP P38489

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula:  $C_{17}H_{21}N_4O_9P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	C	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	D	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is NICOTINIC ACID (three-letter code: NIO) (formula:  $C_6H_5NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	6	1	2		
3	B	1	Total	C	N	O	0	0
			9	6	1	2		
3	C	1	Total	C	N	O	0	0
			9	6	1	2		
3	D	1	Total	C	N	O	0	0
			9	6	1	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	164	Total	O	0	0
			164	164		
4	B	179	Total	O	0	0
			179	179		
4	C	157	Total	O	0	0
			157	157		
4	D	143	Total	O	0	0
			143	143		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: OXYGEN-INSENSITIVE NAD(P)H NITROREDUCTASE

Chain A: 





- Molecule 1: OXYGEN-INSENSITIVE NAD(P)H NITROREDUCTASE

Chain B: 



- Molecule 1: OXYGEN-INSENSITIVE NAD(P)H NITROREDUCTASE

Chain C: 



- Molecule 1: OXYGEN-INSENSITIVE NAD(P)H NITROREDUCTASE

Chain D: 



K141	Q142	V143	Y144	F150	G153	V154	L157	P163	I164	E165	D173	L178	K179	E180	K181	G182	Y183	V189	P190	V191	S206	R207	L208	P209	Q210	N211	I212	T213	L214	T215	E216	V217
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.74Å 119.57Å 143.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 2.40	Depositor
% Data completeness (in resolution range)	(Not available) (100.00-2.40)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	4.90	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.198 , 0.242	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7511	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, NIO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/1708	0.56	0/2306
1	B	0.38	0/1708	0.57	0/2306
1	C	0.36	0/1708	0.58	0/2306
1	D	0.36	0/1708	0.57	0/2306
All	All	0.36	0/6832	0.57	0/9224

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1677	0	1661	53	0
1	B	1677	0	1661	48	0
1	C	1677	0	1661	42	0
1	D	1677	0	1661	55	0
2	A	31	0	18	1	0
2	B	31	0	19	2	0
2	C	31	0	16	3	0
2	D	31	0	16	4	0
3	A	9	0	5	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	9	0	5	0	0
3	C	9	0	5	0	0
3	D	9	0	5	0	0
4	A	164	0	0	5	0
4	B	179	0	0	2	0
4	C	157	0	0	3	0
4	D	143	0	0	4	0
All	All	7511	0	6733	177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:49:ILE:HG23	1:C:83:VAL:HB	1.23	1.15
1:C:75:MSE:HE3	1:C:188:VAL:HG11	1.46	0.97
1:D:49:ILE:CG2	1:D:83:VAL:HB	1.95	0.94
1:C:60:VAL:HG12	1:C:75:MSE:HE1	1.51	0.92
1:C:60:VAL:HG12	1:C:75:MSE:CE	2.05	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/217 (99%)	212 (99%)	2 (1%)	0	100	100
1	B	214/217 (99%)	209 (98%)	5 (2%)	0	100	100
1	C	214/217 (99%)	210 (98%)	3 (1%)	1 (0%)	29	41
1	D	214/217 (99%)	208 (97%)	5 (2%)	1 (0%)	29	41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	856/868 (99%)	839 (98%)	15 (2%)	2 (0%)	47	62

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	66	GLY
1	D	66	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/174 (102%)	164 (92%)	14 (8%)	12	19
1	B	178/174 (102%)	164 (92%)	14 (8%)	12	19
1	C	178/174 (102%)	165 (93%)	13 (7%)	14	22
1	D	178/174 (102%)	160 (90%)	18 (10%)	7	11
All	All	712/696 (102%)	653 (92%)	59 (8%)	11	17

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	157	LEU
1	C	76	LEU
1	D	157	LEU
1	B	178	LEU
1	C	33	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	149	ASN
1	C	29	GLN
1	D	29	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	146	ASN
1	C	149	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMN	C	705	-	31,33,33	3.45	15 (48%)	40,50,50	8.04	29 (72%)
2	FMN	B	703	-	31,33,33	3.44	16 (51%)	40,50,50	8.27	27 (67%)
2	FMN	A	701	-	31,33,33	3.40	15 (48%)	40,50,50	6.70	21 (52%)
2	FMN	D	707	-	31,33,33	3.40	14 (45%)	40,50,50	10.37	29 (72%)
3	NIO	B	702	-	7,9,9	3.07	3 (42%)	8,11,11	1.34	1 (12%)
3	NIO	A	704	-	7,9,9	2.99	3 (42%)	8,11,11	1.36	1 (12%)
3	NIO	C	708	-	7,9,9	3.00	3 (42%)	8,11,11	1.39	1 (12%)
3	NIO	D	706	-	7,9,9	3.04	3 (42%)	8,11,11	1.38	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	C	705	-	3/3/4/4	6/18/18/18	0/3/3/3
2	FMN	B	703	-	3/3/4/4	7/18/18/18	0/3/3/3
2	FMN	A	701	-	-	5/18/18/18	0/3/3/3
2	FMN	D	707	-	2/2/4/4	8/18/18/18	0/3/3/3
3	NIO	B	702	-	-	0/0/4/4	0/1/1/1
3	NIO	A	704	-	-	0/0/4/4	0/1/1/1
3	NIO	C	708	-	-	0/0/4/4	0/1/1/1
3	NIO	D	706	-	-	0/0/4/4	0/1/1/1

The worst 5 of 72 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	705	FMN	C4A-C10	11.99	1.50	1.38
2	A	701	FMN	C4A-C10	11.79	1.50	1.38
2	B	703	FMN	C4A-C10	11.56	1.50	1.38
2	D	707	FMN	C4A-C10	11.27	1.50	1.38
2	D	707	FMN	C1'-N10	7.37	1.55	1.48

The worst 5 of 110 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	707	FMN	C4A-N5-C5A	30.44	147.19	116.77
2	B	703	FMN	C4A-N5-C5A	28.47	145.22	116.77
2	D	707	FMN	C4'-C3'-C2'	26.33	168.13	113.36
2	C	705	FMN	C4A-N5-C5A	25.39	142.14	116.77
2	A	701	FMN	C4A-N5-C5A	22.13	138.89	116.77

5 of 8 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	705	FMN	C4'
2	C	705	FMN	C2'
2	C	705	FMN	C3'
2	B	703	FMN	C4'
2	B	703	FMN	C2'

5 of 26 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	705	FMN	C2'-C1'-N10-C9A
2	C	705	FMN	C2'-C1'-N10-C10
2	C	705	FMN	C1'-C2'-C3'-O3'
2	C	705	FMN	C3'-C4'-C5'-O5'
2	B	703	FMN	C2'-C1'-N10-C10

There are no ring outliers.


4 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	705	FMN	3	0
2	B	703	FMN	2	0
2	A	701	FMN	1	0
2	D	707	FMN	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.