



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 11:39 PM BST

PDB ID : 6ICY
Title : Crystal structure of H7 hemagglutinin mutant H7-AGTL (V186G, P221T)
from the influenza virus A/Anhui/1/2013 (H7N9)
Authors : Gao, G.F.; Xu, Y.; Qi, J.X.
Deposited on : 2018-09-07
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2392	1484	433	460	15	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	176	GLY	VAL	engineered mutation	UNP R4NN21
A	211	THR	PRO	engineered mutation	UNP R4NN21

- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	168	1364	843	236	278	7	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	80	Total	O	0	0
			80	80		
4	B	28	Total	O	0	0
			28	28		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemagglutinin HA1 chain



- Molecule 2: Hemagglutinin HA2 chain



4 Data and refinement statistics i

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	116.54Å 116.54Å 295.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.35 – 2.40 38.35 – 2.40	Depositor EDS
% Data completeness (in resolution range)	90.0 (38.35-2.40) 90.1 (38.35-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 2.39Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.244 , 0.267 0.244 , 0.266	Depositor DCC
R_{free} test set	1327 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	36.8	Xtrriage
Anisotropy	0.040	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.007 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.019 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.010 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3906	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.25	0/2437	0.48	1/3293 (0.0%)
2	B	0.24	0/1388	0.51	3/1871 (0.2%)
All	All	0.25	0/3825	0.49	4/5164 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	328	ALA	CB-CA-C	-10.16	94.86	110.10
2	B	448	ARG	CB-CA-C	-6.99	96.41	110.40
2	B	448	ARG	N-CA-C	5.97	127.12	111.00
1	A	203	PHE	CB-CA-C	5.38	121.15	110.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	201	GLN	Peptide
2	B	328	ALA	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2392	0	2346	39	0
2	B	1364	0	1261	24	1
3	A	28	0	26	1	0
3	B	14	0	13	0	0
4	A	80	0	0	20	0
4	B	28	0	0	9	0
All	All	3906	0	3646	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:377:ILE:HG22	4:B:602:HOH:O	1.60	1.00
2:B:411:GLU:OE2	4:B:601:HOH:O	1.83	0.96
1:A:164:SER:OG	4:A:702:HOH:O	1.87	0.93
1:A:179:ALA:N	4:A:707:HOH:O	2.02	0.92
2:B:411:GLU:HG2	4:B:601:HOH:O	1.73	0.88
2:B:377:ILE:C	4:B:602:HOH:O	2.12	0.87
1:A:118:GLY:O	4:A:705:HOH:O	1.95	0.84
1:A:177:SER:C	4:A:707:HOH:O	2.17	0.83
2:B:377:ILE:O	4:B:602:HOH:O	1.96	0.83
1:A:186:GLY:N	4:A:709:HOH:O	2.19	0.75
1:A:95:GLU:OE2	4:A:708:HOH:O	2.05	0.74
2:B:328:ALA:N	2:B:329:GLY:HA3	2.05	0.71
1:A:201:GLN:HA	1:A:202:SER:HB3	1.74	0.69
1:A:186:GLY:N	4:A:715:HOH:O	2.26	0.69
1:A:185:TYR:C	4:A:709:HOH:O	2.32	0.66
2:B:388:ASP:OD2	2:B:406:ARG:NH2	2.27	0.65
2:B:411:GLU:CG	4:B:601:HOH:O	2.40	0.65
1:A:122:ASN:HB2	4:A:716:HOH:O	1.96	0.64
1:A:178:THR:N	4:A:707:HOH:O	2.31	0.61
1:A:168:ILE:O	1:A:224:TRP:HA	2.01	0.59
2:B:461:ILE:HD11	4:B:628:HOH:O	2.04	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:LEU:HD21	2:B:443:VAL:HG21	1.86	0.56
1:A:141:TRP:N	4:A:704:HOH:O	1.93	0.55
1:A:313:VAL:N	4:A:703:HOH:O	1.87	0.54
2:B:375:ARG:NH2	4:B:603:HOH:O	2.20	0.53
1:A:201:GLN:CA	1:A:202:SER:HB3	2.39	0.52
2:B:327:ILE:N	2:B:433:ASP:OD1	2.41	0.52
1:A:105:GLY:HA2	1:A:254:SER:HB3	1.91	0.52
1:A:152:PRO:CD	4:A:705:HOH:O	2.57	0.51
1:A:178:THR:C	4:A:707:HOH:O	2.43	0.51
1:A:194:VAL:HA	1:A:234:THR:O	2.11	0.51
1:A:264:ASP:OD2	4:A:710:HOH:O	2.19	0.50
2:B:441:GLU:O	2:B:445:ARG:HG2	2.11	0.49
1:A:169:VAL:HG22	1:A:224:TRP:HB3	1.94	0.49
1:A:48:VAL:HG23	1:A:73:ALA:HB2	1.95	0.48
1:A:297:ARG:NH1	2:B:380:THR:OG1	2.45	0.48
1:A:2:ILE:HD11	2:B:345:PHE:HB3	1.95	0.48
2:B:450:ASN:OD1	2:B:450:ASN:N	2.49	0.46
1:A:163:LYS:NZ	3:A:602:NAG:H3	2.31	0.45
1:A:171:GLY:HA2	1:A:221:ASP:O	2.17	0.44
1:A:268:GLU:OE2	4:A:712:HOH:O	2.21	0.44
1:A:3:CYS:HA	2:B:458:CYS:HA	1.98	0.44
1:A:167:LEU:HB3	1:A:248:SER:HB2	1.99	0.44
1:A:191:LEU:HB2	4:A:740:HOH:O	2.18	0.44
1:A:146:THR:OG1	1:A:147:ASP:N	2.49	0.43
2:B:351:GLN:NE2	4:B:606:HOH:O	2.51	0.43
2:B:462:PHE:HB2	2:B:487:ALA:HB1	1.98	0.43
2:B:327:ILE:C	2:B:329:GLY:HA3	2.39	0.43
2:B:324:PHE:CE2	2:B:434:SER:HB2	2.53	0.43
1:A:105:GLY:O	4:A:711:HOH:O	2.21	0.42
1:A:141:TRP:CB	4:A:704:HOH:O	2.67	0.42
1:A:169:VAL:O	1:A:244:PRO:HB3	2.18	0.42
1:A:290:ARG:HE	2:B:388:ASP:HB2	1.85	0.42
2:B:324:PHE:HB2	2:B:433:ASP:OD2	2.20	0.41
1:A:180:GLU:N	4:A:707:HOH:O	2.21	0.41
2:B:340:ASP:HB2	2:B:357:ALA:HB2	2.02	0.41
1:A:170:TRP:CE2	1:A:223:HIS:HB2	2.56	0.40
1:A:297:ARG:H	1:A:297:ARG:HG2	1.61	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:323:LEU:O	2:B:434:SER:OG[3_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/321 (97%)	293 (94%)	18 (6%)	1 (0%)	41	55
2	B	166/177 (94%)	158 (95%)	8 (5%)	0	100	100
All	All	478/498 (96%)	451 (94%)	26 (5%)	1 (0%)	47	62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/268 (98%)	259 (99%)	3 (1%)	73	87
2	B	144/152 (95%)	142 (99%)	2 (1%)	67	82
All	All	406/420 (97%)	401 (99%)	5 (1%)	71	85

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	HIS
1	A	147	ASP
1	A	297	ARG
2	B	323	LEU
2	B	450	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	501	2	14,14,15	0.31	0	17,19,21	0.48	0
3	NAG	A	601	1	14,14,15	0.26	0	17,19,21	0.41	0
3	NAG	A	602	1	14,14,15	0.26	0	17,19,21	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	501	2	-	1/6/23/26	0/1/1/1
3	NAG	A	601	1	-	2/6/23/26	0/1/1/1
3	NAG	A	602	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	NAG	O5-C5-C6-O6
3	A	601	NAG	O5-C5-C6-O6
3	A	602	NAG	C4-C5-C6-O6
3	A	601	NAG	C4-C5-C6-O6
3	B	501	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/321 (97%)	0.16	23 (7%) 15 13	18, 51, 111, 133	0
2	B	168/177 (94%)	0.79	33 (19%) 1 0	19, 79, 134, 145	0
All	All	482/498 (96%)	0.38	56 (11%) 4 4	18, 59, 125, 145	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	379	LYS	6.9
1	A	202	SER	5.8
2	B	460	GLU	5.3
1	A	211	THR	5.2
2	B	464	LYS	5.1
1	A	118	GLY	5.0
1	A	11	ASN	5.0
2	B	489	GLN	4.8
2	B	353	GLU	4.4
1	A	146	THR	4.0
2	B	356	ALA	3.9
1	A	203	PHE	3.8
2	B	487	ALA	3.8
1	A	119	ILE	3.7
1	A	198	ASN	3.6
2	B	480	HIS	3.6
1	A	155	THR	3.4
2	B	482	LYS	3.4
2	B	348	GLN	3.3
1	A	189	ASN	3.3
1	A	116	TYR	3.1
2	B	462	PHE	3.1
2	B	352	GLY	3.1
1	A	117	SER	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	461	ILE	3.0
2	B	483	TYR	3.0
1	A	241	PHE	2.9
2	B	380	THR	2.8
2	B	452	GLU	2.8
2	B	445	ARG	2.7
1	A	152	PRO	2.7
1	A	145	ASN	2.7
1	A	214	ASN	2.7
2	B	465	CYS	2.6
1	A	209	ALA	2.5
1	A	12	GLY	2.5
1	A	216	LEU	2.5
2	B	488	MET	2.4
2	B	323	LEU	2.4
1	A	2	ILE	2.4
2	B	469	CYS	2.4
1	A	193	THR	2.4
2	B	340	ASP	2.3
2	B	481	SER	2.3
2	B	468	ASP	2.3
1	A	122	ASN	2.2
2	B	451	ALA	2.2
2	B	355	THR	2.2
2	B	463	HIS	2.2
2	B	479	ASP	2.2
2	B	478	TYR	2.1
2	B	344	GLY	2.1
2	B	347	HIS	2.1
1	A	10	SER	2.1
2	B	381	ASN	2.0
2	B	351	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	602	14/15	0.71	0.36	78,92,107,113	0
3	NAG	A	601	14/15	0.72	0.38	74,91,119,125	0
3	NAG	B	501	14/15	0.92	0.15	48,60,66,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.