



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 2, 2023 – 10:00 PM EDT

PDB ID : 1IDW  
Title : STRUCTURE OF THE HYBRID RNA/DNA R-GCUUCGGC-D[CL]U IN PRESENCE OF RH(NH<sub>3</sub>)<sub>6</sub>+++  
Authors : Cruse, W.; Saludjian, P.; Neuman, A.; Prange, T.  
Deposited on : 2001-04-05  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.34


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6793 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	9	
1	B	9	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	UCL	B	9[A]	-	-	X	-
1	UCL	B	9[B]	X	-	X	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 455 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is DNA/RNA hybrid called 5'-R(\*GP\*CP\*UP\*UP\*CP\*GP\*GP\*C)-D(P\*(UCL))-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	Cl	N	O	P			
1	A	9	Total 186	C 84	Cl 1	N 30	O 63	P 8	0	0	0
1	B	9	Total 206	C 93	Cl 2	N 32	O 70	P 9	0	1	0

- Molecule 2 is RHODIUM HEXAMINE ION (three-letter code: RHD) (formula: H<sub>18</sub>N<sub>6</sub>Rh).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	N	Rh		
2	A	1	Total 7	N 6	Rh 1	0	0
2	A	1	Total 7	N 6	Rh 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	B	1	Total Cl 2 2	0	1

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	28	Total O 28 28	0	0
4	B	18	Total O 18 18	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: 5'-R(\*GP\*CP\*UP\*UP\*CP\*GP\*GP\*C)-D(P\*(UCL))-3'

Chain A: 



- Molecule 1: 5'-R(\*GP\*CP\*UP\*UP\*CP\*GP\*GP\*C)-D(P\*(UCL))-3'

Chain B: 



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.82Å 19.38Å 50.30Å 90.00° 109.90° 90.00°	Depositor
Resolution (Å)	6.00 – 1.80	Depositor
% Data completeness (in resolution range)	95.0 (6.00-1.80)	Depositor
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.04	Depositor
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.172 , 0.202	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	455	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: UCL, RHD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.33	0/184	2.28	13/285 (4.6%)
1	B	1.49	3/184 (1.6%)	2.44	20/285 (7.0%)
All	All	1.42	3/368 (0.8%)	2.36	33/570 (5.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	8	C	C4'-C3'	-5.83	1.46	1.52
1	B	8	C	C3'-C2'	-5.63	1.46	1.52
1	B	8	C	C2'-O2'	-5.03	1.35	1.41

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	U	C5-C4-O4	-8.22	120.97	125.90
1	B	8	C	O4'-C1'-N1	-8.02	101.78	108.20
1	B	1	G	C6-N1-C2	-8.02	120.29	125.10
1	A	3	U	C2-N3-C4	-7.87	122.28	127.00
1	B	3	U	C5-C4-O4	-7.23	121.56	125.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	9[B]	UCL	C4'

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	186	0	97	2	0
1	B	206	0	103	40	0
2	A	14	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	28	0	0	0	0
4	B	18	0	0	0	0
All	All	455	0	200	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 69.

The worst 5 of 42 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:9[B]:UCL:C3'	1:B:9[B]:UCL:C4'	1.80	1.55
1:B:8:C:H3'	1:B:9[B]:UCL:C5	1.82	1.07
1:B:8:C:C4'	1:B:9[B]:UCL:O1P	2.21	0.84
1:B:8:C:C3'	1:B:9[B]:UCL:C6	2.59	0.81
1:B:8:C:C4'	1:B:9[A]:UCL:O1P	2.25	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.



### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	UCL	A	9	1	19,21,22	1.94	5 (26%)	24,30,33	4.44	13 (54%)
1	UCL	B	9[B]	1	19,21,22	3.69	6 (31%)	24,30,33	4.13	14 (58%)
1	UCL	B	9[A]	1	19,21,22	3.05	5 (26%)	24,30,33	4.70	17 (70%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	UCL	A	9	1	-	3/7/21/22	0/2/2/2
1	UCL	B	9[B]	1	1/1/4/4	6/7/21/22	0/2/2/2
1	UCL	B	9[A]	1	-	5/7/21/22	0/2/2/2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	9[B]	UCL	C3'-C4'	9.90	1.80	1.53
1	B	9[A]	UCL	P-O5'	-8.24	1.39	1.62
1	B	9[B]	UCL	P-O5'	-8.06	1.39	1.62

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	9[B]	UCL	C2'-C3'	-7.79	1.32	1.52
1	B	9[A]	UCL	O3'-C3'	-6.48	1.29	1.43

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	UCL	C6-N1-C2	-11.29	109.87	121.30
1	B	9[B]	UCL	C6-C5-CL	9.77	130.44	120.86
1	B	9[B]	UCL	C4-C5-CL	-9.46	111.15	117.93
1	B	9[A]	UCL	C4-C5-CL	-9.42	111.18	117.93
1	A	9	UCL	C6-C5-C4	-9.39	115.42	120.65

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	9[B]	UCL	C4'

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	9[A]	UCL	C4'-C5'-O5'-P
1	B	9[B]	UCL	C4'-C5'-O5'-P
1	B	9[B]	UCL	C3'-C4'-C5'-O5'
1	B	9[B]	UCL	C2'-C1'-N1-C6
1	B	9[A]	UCL	C2'-C1'-N1-C2

There are no ring outliers.

3 monomers are involved in 42 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	9	UCL	2	0
1	B	9[B]	UCL	21	0
1	B	9[A]	UCL	19	0

## 5.5 Carbohydrates

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	RHD	A	12	-	0,6,6	-	-	-		
2	RHD	A	11	-	0,6,6	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.