



# wwPDB NMR Structure Validation Summary Report ⓘ

Mar 3, 2022 – 01:03 PM EST

PDB ID : 2IFE  
Title : TRANSLATION INITIATION FACTOR IF3 FROM ESCHERICHIA COLI  
RIBOSOME BINDING DOMAIN (RESIDUES 84-180)  
Authors : De Cock, E.; Garcia, C.; Dardel, F.  
Deposited on : 1998-12-16

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.27  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	100	 43% 15% . . 27% 9%

## 2 Ensemble composition and analysis

This entry contains 24 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 2 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:92-A:99, A:106-A:130, A:140-A:161, A:169-A:177 (64)	0.23	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 5, 8, 10, 12, 13, 14, 15, 16, 17, 20, 21, 22, 23
2	1, 3, 4, 6, 7, 9, 18, 19, 24
Single-model clusters	11

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1527 atoms, of which 784 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PROTEIN (TRANSLATION INITIATION FACTOR IF3).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	91	1527	470	784	135	134	4	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	81	MET	SER	SEE REMARK 999	UNP P0A707
A	82	GLU	LYS	SEE REMARK 999	UNP P0A707
A	83	PHE	GLU	SEE REMARK 999	UNP P0A707

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: PROTEIN (TRANSLATION INITIATION FACTOR IF3)



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 19. Colouring as in section 4.1 above.

- Molecule 1: PROTEIN (TRANSLATION INITIATION FACTOR IF3)



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *DISTANCE GEOMETRY AND RESTRAINED SIMULATED ANNEALING*.

Of the 200 calculated structures, 24 were deposited, based on the following criterion: *LOWEST TARGET FUNCTION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.1
DIANA	structure solution	
X-PLOR	structure solution	3.1

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.19±0.01	0±0/536 ( 0.0± 0.0%)	1.77±0.09	13±3/718 ( 1.8± 0.4%)
All	All	1.19	0/12864 ( 0.0%)	1.77	310/17232 ( 1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.7±0.9
All	All	0	88

There are no bond-length outliers.

5 of 39 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	107	TYR	CB-CG-CD2	-16.36	111.19	121.00	21	17
1	A	98	PHE	CB-CG-CD2	-14.22	110.84	120.80	13	4
1	A	98	PHE	CB-CG-CD1	14.16	130.71	120.80	13	4
1	A	117	PHE	CB-CG-CD2	-12.79	111.84	120.80	21	13
1	A	117	PHE	CB-CG-CD1	11.42	128.79	120.80	10	13

There are no chirality outliers.

5 of 6 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	107	TYR	Sidechain	24
1	A	156	ALA	Peptide	21
1	A	161	PHE	Sidechain,Peptide	18

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	142	MET	Peptide	17
1	A	160	SER	Peptide	2

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	530	569	569	6±2
All	All	12720	13656	13656	138

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

5 of 46 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:156:ALA:HB1	1:A:174:LEU:HD11	0.93	1.38	3	6
1:A:128:LEU:HD21	1:A:141:GLY:HA2	0.72	1.62	14	5
1:A:128:LEU:HD13	1:A:130:PHE:CZ	0.66	2.26	11	2
1:A:128:LEU:HD11	1:A:141:GLY:HA2	0.64	1.69	6	8
1:A:156:ALA:CB	1:A:174:LEU:HD11	0.63	2.24	8	7

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	64/100 (64%)	57±1 (89±2%)	6±1 (9±2%)	1±0 (2±1%)	12	54
All	All	1536/2400 (64%)	1370 (89%)	139 (9%)	27 (2%)	12	54

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occur-



rence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	153	GLN	24
1	A	144	VAL	1
1	A	169	GLN	1
1	A	106	ASP	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	59/90 (66%)	50±2 (84±4%)	9±2 (16±4%)	5	43
All	All	1416/2160 (66%)	1196 (84%)	220 (16%)	5	43

5 of 23 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	108	GLN	24
1	A	152	LEU	24
1	A	107	TYR	22
1	A	118	LEU	21
1	A	128	LEU	19

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided