



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 19, 2023 – 09:32 PM EDT

PDB ID : 5IFM  
Title : Human NONO (p54nrb) Homodimer  
Authors : Knott, G.J.; Bond, C.S.  
Deposited on : 2016-02-26  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	86% 11% .
1	B	261	84% 15% .
1	C	261	82% 15% ..
1	D	261	82% 14% ..
1	E	261	85% 13% .

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	261	
1	G	261	
1	H	261	
1	I	261	
1	J	261	
1	K	261	
1	L	261	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	D	401	-	-	X	-
5	GOL	B	402	-	-	X	-
5	GOL	C	405	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 25317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-POU domain-containing octamer-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	254	2065	1301	369	386	9	0	0	0
1	B	259	2114	1333	380	390	11	0	1	0
1	C	256	2090	1316	374	391	9	0	1	0
1	D	257	2099	1320	379	391	9	0	1	0
1	E	256	2084	1312	374	389	9	0	0	0
1	F	255	2075	1307	373	386	9	0	0	0
1	G	255	2084	1312	373	390	9	0	1	0
1	H	257	2090	1317	375	388	10	0	0	0
1	I	255	2075	1307	372	387	9	0	0	0
1	J	259	2106	1328	377	390	11	0	0	0
1	K	259	2106	1328	377	390	11	0	0	0
1	L	255	2075	1307	372	387	9	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	GLY	-	expression tag	UNP Q15233
B	52	GLY	-	expression tag	UNP Q15233
C	52	GLY	-	expression tag	UNP Q15233
D	52	GLY	-	expression tag	UNP Q15233
E	52	GLY	-	expression tag	UNP Q15233

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	52	GLY	-	expression tag	UNP Q15233
G	52	GLY	-	expression tag	UNP Q15233
H	52	GLY	-	expression tag	UNP Q15233
I	52	GLY	-	expression tag	UNP Q15233
J	52	GLY	-	expression tag	UNP Q15233
K	52	GLY	-	expression tag	UNP Q15233
L	52	GLY	-	expression tag	UNP Q15233

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	C	2	Total Cl 2 2	0	0
2	D	1	Total Cl 1 1	0	0
2	E	1	Total Cl 1 1	0	0
2	F	2	Total Cl 2 2	0	0
2	G	1	Total Cl 1 1	0	0
2	H	1	Total Cl 1 1	0	0
2	L	1	Total Cl 1 1	0	0

- Molecule 3 is PROLINE (three-letter code: PRO) (formula: C<sub>5</sub>H<sub>9</sub>NO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	5	1	2		
3	C	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	K	0	0
			1	1		
4	B	1	Total	K	0	0
			1	1		
4	C	1	Total	K	0	0
			1	1		
4	D	1	Total	K	0	0
			1	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	33	Total O 33 33	0	0
6	B	26	Total O 26 26	0	0
6	C	24	Total O 24 24	0	0
6	D	37	Total O 37 37	0	0
6	E	15	Total O 15 15	0	0
6	F	7	Total O 7 7	0	0
6	G	19	Total O 19 19	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
6	H	10	Total O 10 10	0	0
6	I	5	Total O 5 5	0	0
6	J	4	Total O 4 4	0	0
6	K	7	Total O 7 7	0	0
6	L	7	Total O 7 7	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Non-POU domain-containing octamer-binding protein




- Molecule 1: Non-POU domain-containing octamer-binding protein



- Molecule 1: Non-POU domain-containing octamer-binding protein





- Molecule 1: Non-POU domain-containing octamer-binding protein










- Molecule 1: Non-POU domain-containing octamer-binding protein





- Molecule 1: Non-POU domain-containing octamer-binding protein




- Molecule 1: Non-POU domain-containing octamer-binding protein




- Molecule 1: Non-POU domain-containing octamer-binding protein





- Molecule 1: Non-POU domain-containing octamer-binding protein








- Molecule 1: Non-POU domain-containing octamer-binding protein



- Molecule 1: Non-POU domain-containing octamer-binding protein



- Molecule 1: Non-POU domain-containing octamer-binding protein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.15Å 407.18Å 68.96Å 90.00° 97.75° 90.00°	Depositor
Resolution (Å)	48.15 – 2.60 48.15 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.7 (48.15-2.60) 98.8 (48.15-2.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.85 (at 2.61Å)	Xtrriage
Refinement program	BUSTER-TNT 2.10.0	Depositor
R, $R_{free}$	0.197 , 0.234 0.213 , 0.254	Depositor DCC
$R_{free}$ test set	5520 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.7	Xtrriage
Anisotropy	0.220	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 56.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.038 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	25317	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, K, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/2104	0.73	0/2823
1	B	0.55	0/2157	0.74	0/2893
1	C	0.56	0/2133	0.76	0/2862
1	D	0.60	0/2139	0.78	0/2869
1	E	0.50	0/2124	0.70	0/2850
1	F	0.48	0/2115	0.69	0/2838
1	G	0.51	0/2124	0.70	0/2850
1	H	0.50	0/2130	0.73	0/2858
1	I	0.47	0/2115	0.69	0/2838
1	J	0.46	0/2146	0.68	1/2879 (0.0%)
1	K	0.47	0/2146	0.71	0/2879
1	L	0.49	0/2115	0.71	0/2838
All	All	0.51	0/25548	0.72	1/34277 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	119	ARG	CG-CD-NE	5.30	122.93	111.80

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	0	2064	23	0
1	B	2114	0	2124	35	0
1	C	2090	0	2085	32	0
1	D	2099	0	2094	32	0
1	E	2084	0	2079	20	0
1	F	2075	0	2073	15	0
1	G	2084	0	2076	21	0
1	H	2090	0	2091	23	0
1	I	2075	0	2071	18	0
1	J	2106	0	2111	13	0
1	K	2106	0	2111	17	0
1	L	2075	0	2071	17	0
2	A	1	0	0	1	0
2	C	2	0	0	1	0
2	D	1	0	0	2	0
2	E	1	0	0	1	0
2	F	2	0	0	1	0
2	G	1	0	0	1	0
2	H	1	0	0	1	0
2	L	1	0	0	0	0
3	A	8	0	7	2	0
3	C	8	0	7	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	B	12	0	16	8	0
5	C	12	0	16	7	0
5	D	6	0	8	2	0
6	A	33	0	0	1	0
6	B	26	0	0	0	0
6	C	24	0	0	1	0
6	D	37	0	0	0	0
6	E	15	0	0	1	0
6	F	7	0	0	0	0
6	G	19	0	0	0	0
6	H	10	0	0	0	0
6	I	5	0	0	0	0
6	J	4	0	0	0	0
6	K	7	0	0	0	0
6	L	7	0	0	0	0
All	All	25317	0	25104	191	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 191 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:285:VAL:HG21	1:L:285:VAL:HG21	1.43	0.98
1:B:84:ASP:HB2	1:B:135:ARG:HH22	1.31	0.95
1:C:61:ASN:HA	5:C:405:GOL:H32	1.55	0.88
1:I:285:VAL:HG21	1:J:285:VAL:HG21	1.57	0.84
1:L:156:PRO:HD2	1:L:159:VAL:HG21	1.58	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/261 (97%)	242 (96%)	10 (4%)	0	100	100
1	B	258/261 (99%)	249 (96%)	9 (4%)	0	100	100
1	C	255/261 (98%)	248 (97%)	5 (2%)	2 (1%)	19	39
1	D	256/261 (98%)	248 (97%)	6 (2%)	2 (1%)	19	39
1	E	254/261 (97%)	249 (98%)	5 (2%)	0	100	100
1	F	253/261 (97%)	246 (97%)	7 (3%)	0	100	100
1	G	254/261 (97%)	246 (97%)	8 (3%)	0	100	100
1	H	255/261 (98%)	246 (96%)	9 (4%)	0	100	100
1	I	253/261 (97%)	245 (97%)	8 (3%)	0	100	100
1	J	257/261 (98%)	250 (97%)	6 (2%)	1 (0%)	34	57
1	K	257/261 (98%)	249 (97%)	7 (3%)	1 (0%)	34	57
1	L	253/261 (97%)	244 (96%)	9 (4%)	0	100	100
All	All	3057/3132 (98%)	2962 (97%)	89 (3%)	6 (0%)	47	71

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	85	ILE
1	D	53	GLU
1	J	107	LYS
1	C	107	LYS
1	D	107	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	222/228 (97%)	216 (97%)	6 (3%)	44 71
1	B	228/228 (100%)	223 (98%)	5 (2%)	52 76
1	C	225/228 (99%)	214 (95%)	11 (5%)	25 48
1	D	225/228 (99%)	209 (93%)	16 (7%)	14 29
1	E	224/228 (98%)	215 (96%)	9 (4%)	31 57
1	F	223/228 (98%)	211 (95%)	12 (5%)	22 44
1	G	224/228 (98%)	206 (92%)	18 (8%)	12 24
1	H	225/228 (99%)	211 (94%)	14 (6%)	18 37
1	I	223/228 (98%)	208 (93%)	15 (7%)	16 33
1	J	227/228 (100%)	212 (93%)	15 (7%)	16 33
1	K	227/228 (100%)	211 (93%)	16 (7%)	15 30
1	L	223/228 (98%)	208 (93%)	15 (7%)	16 33
All	All	2696/2736 (98%)	2544 (94%)	152 (6%)	21 42

5 of 152 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	259	GLN
1	L	132	MET
1	J	301	GLU
1	K	252	GLU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	L	263	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	308	GLN
1	H	307	HIS
1	L	154	ASN
1	D	280	GLN
1	B	308	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 14 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PRO	A	402	-	8,8,8	0.93	0	10,10,10	1.16	1 (10%)
5	GOL	B	401	-	5,5,5	0.38	0	5,5,5	0.89	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PRO	C	403	-	8,8,8	0.86	0	10,10,10	1.02	0
5	GOL	B	402	-	5,5,5	0.30	0	5,5,5	0.49	0
5	GOL	C	405	-	5,5,5	0.25	0	5,5,5	1.01	0
5	GOL	C	404	-	5,5,5	0.31	0	5,5,5	0.68	0
5	GOL	D	402	-	5,5,5	0.26	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PRO	A	402	-	-	2/4/11/11	0/1/1/1
5	GOL	B	401	-	-	2/4/4/4	-
3	PRO	C	403	-	-	2/4/11/11	0/1/1/1
5	GOL	B	402	-	-	2/4/4/4	-
5	GOL	C	405	-	-	2/4/4/4	-
5	GOL	C	404	-	-	3/4/4/4	-
5	GOL	D	402	-	-	4/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	PRO	C-CA-N	2.03	114.73	106.73

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	402	GOL	O1-C1-C2-C3
5	C	405	GOL	C1-C2-C3-O3
5	D	402	GOL	O1-C1-C2-C3
5	D	402	GOL	C1-C2-C3-O3
5	C	405	GOL	O2-C2-C3-O3

There are no ring outliers.

7 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	PRO	2	0
5	B	401	GOL	1	0
3	C	403	PRO	1	0
5	B	402	GOL	7	0
5	C	405	GOL	4	0
5	C	404	GOL	3	0
5	D	402	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	254/261 (97%)	-0.13	1 (0%) 92 91	22, 50, 81, 113	0
1	B	259/261 (99%)	-0.08	8 (3%) 49 42	23, 49, 96, 148	0
1	C	256/261 (98%)	-0.09	6 (2%) 60 54	22, 46, 86, 132	0
1	D	257/261 (98%)	-0.22	4 (1%) 72 68	25, 46, 100, 148	0
1	E	256/261 (98%)	-0.03	10 (3%) 39 32	28, 63, 116, 138	0
1	F	255/261 (97%)	-0.04	5 (1%) 65 60	35, 64, 119, 140	0
1	G	255/261 (97%)	-0.17	1 (0%) 92 91	25, 55, 103, 129	0
1	H	257/261 (98%)	-0.07	5 (1%) 66 62	35, 63, 103, 126	0
1	I	255/261 (97%)	0.24	16 (6%) 20 15	41, 71, 123, 136	0
1	J	259/261 (99%)	0.44	24 (9%) 8 6	45, 80, 143, 165	0
1	K	259/261 (99%)	0.17	12 (4%) 32 26	37, 66, 108, 124	0
1	L	255/261 (97%)	0.02	8 (3%) 49 42	38, 72, 109, 141	0
All	All	3077/3132 (98%)	0.00	100 (3%) 47 40	22, 60, 115, 165	0

The worst 5 of 100 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	81	LEU	6.8
1	J	309	VAL	6.3
1	L	305	HIS	5.4
1	B	307	HIS	5.1
1	L	303	ALA	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	K	C	406	1/1	0.76	0.20	76,76,76,76	0
2	CL	F	402	1/1	0.80	0.15	79,79,79,79	0
5	GOL	B	401	6/6	0.80	0.30	38,41,46,50	0
5	GOL	B	402	6/6	0.81	0.36	48,55,58,59	0
5	GOL	C	404	6/6	0.81	0.26	49,51,57,61	0
5	GOL	D	402	6/6	0.81	0.34	56,69,71,72	0
2	CL	L	401	1/1	0.82	0.18	73,73,73,73	0
4	K	A	403	1/1	0.86	0.15	73,73,73,73	0
5	GOL	C	405	6/6	0.89	0.22	30,42,46,47	0
3	PRO	C	403	8/8	0.89	0.22	64,67,107,128	0
4	K	D	403	1/1	0.90	0.16	71,71,71,71	0
3	PRO	A	402	8/8	0.91	0.29	45,47,62,82	0
4	K	B	403	1/1	0.92	0.17	71,71,71,71	0
2	CL	D	401	1/1	0.92	0.22	90,90,90,90	0
2	CL	C	401	1/1	0.93	0.13	56,56,56,56	0
2	CL	F	401	1/1	0.94	0.26	64,64,64,64	0
2	CL	C	402	1/1	0.94	0.20	59,59,59,59	0
2	CL	A	401	1/1	0.96	0.25	48,48,48,48	0
2	CL	G	401	1/1	0.96	0.13	70,70,70,70	0
2	CL	H	401	1/1	0.96	0.09	63,63,63,63	0
2	CL	E	401	1/1	0.99	0.14	52,52,52,52	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.