wwPDB X-ray Structure Validation Summary Report (i)

Jun 19, 2024 – 08:38 AM EDT

PDB ID	:	4IIG
Title	:	Crystal structure of beta-glucosidase 1 from Aspergillus aculeatus in complex with D-glucose
Authors	:	Suzuki, K.; Sumitani, J.; Kawaguchi, T.; Fushinobu, S.
Deposited on	:	2012-12-20
Resolution	:	2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	841	% 91% 8% •
1	B	841	89% 9% •
2	C	5	100%
3	D	3	67% 33%
3	G	3	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	L	3	100%
3	N	3	100%
4	E	2	50% 50%
5	F	10	10% 90%
5	M	10	20% 80%
6	H	7	100%
6	O	7	86% 14%
7	I	7	86% 14%
8	J	7	14% 86%
9	K	6	17% 67% 17%
10	P	8	75% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
12	MRD	A	941	-	-	X	-

2 Entry composition (i)

There are 14 unique types of molecules in this entry. The entry contains 15202 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	834	6387	4031	1097	1241	18	0	0	0
1	B	832	6375	4023	1095	1239	18	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	5	61	34	2	25	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	3	39	22	2	15	0	0	0
3	G	3	39	22	2	15	0	0	0
3	L	3	39	22	2	15	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	N	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	E	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	F	10	116	64	2	50	0	0	0
5	M	10	116	64	2	50	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	H	7	83	46	2	35	0	0	0
6	O	7	83	46	2	35	0	0	0

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
7	I	7	83	46	2	35	0	0	0

- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
8	J	7	83	46	2	35	0	0	0

- Molecule 9 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
9	K	6	72	40	2	30	0	0	0

- Molecule 10 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
10	P	8	94	52	2	40	0	0	0

- Molecule 11 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
11	A	1	14	8	1	5	0	0
			Total	C	N	O		
11	A	1	14	8	1	5	0	0
			Total	C	N	O		
11	B	1	14	8	1	5	0	0
			Total	C	N	O		
11	B	1	14	8	1	5	0	0
			Total	C	N	O		
11	B	1	14	8	1	5	0	0

- Molecule 12 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula:

$C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
12	A	1	8	6	2		0	0
			Total	C	O			
12	A	1	8	6	2		0	0
			Total	C	O			
12	B	1	8	6	2		0	0
			Total	C	O			
12	B	1	8	6	2		0	0

- Molecule 13 is beta-D-glucopyranose (three-letter code: BGC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
13	A	1	12	6	6	0	0
			Total	C	O		
13	A	1	12	6	6	0	0
			Total	C	O		
13	B	1	12	6	6	0	0
			Total	C	O		
13	B	1	12	6	6	0	0

- Molecule 14 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
14	A	626	626	626	0	0
			Total	O		
14	B	689	689	689	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-glucosidase 1

Chain A:



• Molecule 1: Beta-glucosidase 1

Chain B:



• Molecule 2: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:



• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 67% 33%

NAG1 NAG2 BMA3

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 100%

NAG1 NAG2 BMA3

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 100%

NAG1 NAG2 BMA3

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N: 100%

NAG1 NAG2 BMA3

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 50% 50%

NAG1 NAG2

• Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 10% 90%

NAG1 NAG2 BMA3 MAN4 MAN5 MAN6 MAN7 MAN8 MAN9 MAN10

• Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M: 



• Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 

• Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O: 

• Molecule 7: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:

• Molecule 8: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 

• Molecule 9: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 



• Molecule 10: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	82.48Å 121.76Å 221.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.99 – 2.30 45.99 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (45.99-2.30) 99.1 (45.99-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.22	Depositor
$< I/\sigma(I) >$ [1]	2.44 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.143 , 0.200 0.145 , 0.202	Depositor DCC
R_{free} test set	4977 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	22.6	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 38.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	15202	wwPDB-VP
Average B, all atoms (Å^2)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, NAG, MAN, BMA, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.92	2/6550 (0.0%)	0.94	11/8930 (0.1%)
1	B	0.98	5/6538 (0.1%)	0.98	21/8913 (0.2%)
All	All	0.95	7/13088 (0.1%)	0.96	32/17843 (0.2%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	541	CYS	CB-SG	9.81	1.99	1.82
1	B	541	CYS	CB-SG	8.09	1.96	1.82
1	A	541	CYS	CA-CB	7.66	1.70	1.53
1	B	208	TYR	CE1-CZ	6.49	1.47	1.38
1	B	541	CYS	CA-CB	6.28	1.67	1.53

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	329	ARG	NE-CZ-NH2	-20.97	109.81	120.30
1	A	329	ARG	NE-CZ-NH2	-17.88	111.36	120.30
1	B	329	ARG	NE-CZ-NH1	16.60	128.60	120.30
1	B	521	ARG	NE-CZ-NH2	-16.29	112.15	120.30
1	A	521	ARG	NE-CZ-NH2	-16.13	112.23	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6387	0	6097	53	0
1	B	6375	0	6082	42	0
2	C	61	0	52	0	0
3	D	39	0	34	1	0
3	G	39	0	34	0	0
3	L	39	0	34	0	0
3	N	39	0	34	0	0
4	E	28	0	25	4	0
5	F	116	0	97	0	0
5	M	116	0	97	0	0
6	H	83	0	70	2	0
6	O	83	0	70	1	0
7	I	83	0	70	2	0
8	J	83	0	70	0	0
9	K	72	0	61	2	0
10	P	94	0	79	2	0
11	A	28	0	26	0	0
11	B	42	0	39	1	0
12	A	16	0	28	8	0
12	B	16	0	28	2	0
13	A	24	0	24	2	0
13	B	24	0	24	3	0
14	A	626	0	0	11	0
14	B	689	0	0	7	0
All	All	15202	0	13175	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:THR:HG21	1:A:333:MET:HE2	1.48	0.95
1:A:63:THR:CG2	1:A:333:MET:HE2	2.01	0.90
12:A:941:MRD:HMC2	14:B:1103:HOH:O	1.74	0.88
1:A:556:VAL:HG13	1:A:560:TYR:HB3	1.57	0.83
1:A:63:THR:HG21	1:A:333:MET:CE	2.08	0.83

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	830/841 (99%)	794 (96%)	36 (4%)	0	100	100
1	B	828/841 (98%)	802 (97%)	26 (3%)	0	100	100
All	All	1658/1682 (99%)	1596 (96%)	62 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	671/677 (99%)	663 (99%)	8 (1%)	71	84
1	B	670/677 (99%)	663 (99%)	7 (1%)	76	87
All	All	1341/1354 (99%)	1326 (99%)	15 (1%)	73	86

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	701	THR
1	B	700	LEU
1	B	70	LEU
1	B	853	HIS
1	B	407	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	540	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

81 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	C	1	2,1	14,14,15	1.15	1 (7%)	17,19,21	1.08	1 (5%)
2	NAG	C	2	2	14,14,15	0.89	1 (7%)	17,19,21	1.47	5 (29%)
2	BMA	C	3	2	11,11,12	1.16	1 (9%)	15,15,17	1.60	4 (26%)
2	MAN	C	4	2	11,11,12	1.07	1 (9%)	15,15,17	3.22	8 (53%)
2	MAN	C	5	2	11,11,12	0.88	0	15,15,17	2.20	4 (26%)
3	NAG	D	1	3,1	14,14,15	1.02	1 (7%)	17,19,21	1.52	2 (11%)
3	NAG	D	2	3	14,14,15	0.75	0	17,19,21	1.60	5 (29%)
3	BMA	D	3	3	11,11,12	0.87	0	15,15,17	2.96	9 (60%)
4	NAG	E	1	4,1	14,14,15	0.72	0	17,19,21	1.44	4 (23%)
4	NAG	E	2	4	14,14,15	0.67	0	17,19,21	2.20	8 (47%)
5	NAG	F	1	5,1	14,14,15	0.93	0	17,19,21	1.39	2 (11%)
5	MAN	F	10	5	11,11,12	1.06	1 (9%)	15,15,17	2.24	4 (26%)
5	NAG	F	2	5	14,14,15	0.70	0	17,19,21	1.67	4 (23%)
5	BMA	F	3	5	11,11,12	1.09	1 (9%)	15,15,17	1.20	1 (6%)
5	MAN	F	4	5	11,11,12	0.65	0	15,15,17	1.94	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	MAN	F	5	5	11,11,12	0.79	0	15,15,17	1.33	2 (13%)
5	MAN	F	6	5	11,11,12	1.01	1 (9%)	15,15,17	2.11	6 (40%)
5	MAN	F	7	5	11,11,12	1.07	1 (9%)	15,15,17	1.40	3 (20%)
5	MAN	F	8	5	11,11,12	0.55	0	15,15,17	0.88	0
5	MAN	F	9	5	11,11,12	0.75	0	15,15,17	1.25	2 (13%)
3	NAG	G	1	3,1	14,14,15	0.94	1 (7%)	17,19,21	1.40	2 (11%)
3	NAG	G	2	3	14,14,15	0.80	1 (7%)	17,19,21	1.42	1 (5%)
3	BMA	G	3	3	11,11,12	0.74	0	15,15,17	1.37	2 (13%)
6	NAG	H	1	6,1	14,14,15	0.72	0	17,19,21	1.51	2 (11%)
6	NAG	H	2	6	14,14,15	0.84	1 (7%)	17,19,21	2.26	4 (23%)
6	BMA	H	3	6	11,11,12	0.78	0	15,15,17	0.74	0
6	MAN	H	4	6	11,11,12	0.85	0	15,15,17	1.49	3 (20%)
6	MAN	H	5	6	11,11,12	1.09	1 (9%)	15,15,17	2.14	4 (26%)
6	MAN	H	6	6	11,11,12	0.75	0	15,15,17	1.52	2 (13%)
6	MAN	H	7	6	11,11,12	0.68	0	15,15,17	1.50	4 (26%)
7	NAG	I	1	7,1	14,14,15	1.00	1 (7%)	17,19,21	1.92	3 (17%)
7	NAG	I	2	7	14,14,15	0.75	0	17,19,21	1.19	2 (11%)
7	BMA	I	3	7	11,11,12	1.18	0	15,15,17	2.31	6 (40%)
7	MAN	I	4	7	11,11,12	0.87	0	15,15,17	1.61	2 (13%)
7	MAN	I	5	7	11,11,12	0.97	0	15,15,17	1.52	2 (13%)
7	MAN	I	6	7	11,11,12	0.65	0	15,15,17	1.51	1 (6%)
7	MAN	I	7	7	11,11,12	0.82	0	15,15,17	2.11	5 (33%)
8	NAG	J	1	8,1	14,14,15	0.93	0	17,19,21	2.01	3 (17%)
8	NAG	J	2	8	14,14,15	0.78	1 (7%)	17,19,21	1.48	2 (11%)
8	BMA	J	3	8	11,11,12	1.03	1 (9%)	15,15,17	1.54	4 (26%)
8	MAN	J	4	8	11,11,12	0.61	0	15,15,17	1.52	3 (20%)
8	MAN	J	5	8	11,11,12	0.98	1 (9%)	15,15,17	1.22	2 (13%)
8	MAN	J	6	8	11,11,12	0.84	0	15,15,17	1.03	0
8	MAN	J	7	8	11,11,12	1.20	1 (9%)	15,15,17	2.39	7 (46%)
9	NAG	K	1	9,1	14,14,15	0.90	0	17,19,21	1.74	3 (17%)
9	NAG	K	2	9	14,14,15	1.16	1 (7%)	17,19,21	1.51	3 (17%)
9	BMA	K	3	9	11,11,12	0.71	0	15,15,17	1.56	3 (20%)
9	MAN	K	4	9	11,11,12	0.65	0	15,15,17	2.43	2 (13%)
9	MAN	K	5	9	11,11,12	0.81	0	15,15,17	0.91	0
9	MAN	K	6	9	11,11,12	1.46	1 (9%)	15,15,17	1.93	5 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	NAG	L	1	3,1	14,14,15	1.04	1 (7%)	17,19,21	1.76	3 (17%)
3	NAG	L	2	3	14,14,15	0.88	1 (7%)	17,19,21	1.21	2 (11%)
3	BMA	L	3	3	11,11,12	0.62	0	15,15,17	2.40	5 (33%)
5	NAG	M	1	5,1	14,14,15	0.93	0	17,19,21	1.89	5 (29%)
5	MAN	M	10	5	11,11,12	1.12	2 (18%)	15,15,17	2.24	5 (33%)
5	NAG	M	2	5	14,14,15	0.77	0	17,19,21	0.94	0
5	BMA	M	3	5	11,11,12	0.96	0	15,15,17	1.19	2 (13%)
5	MAN	M	4	5	11,11,12	0.70	0	15,15,17	1.93	5 (33%)
5	MAN	M	5	5	11,11,12	0.78	0	15,15,17	1.30	2 (13%)
5	MAN	M	6	5	11,11,12	1.03	0	15,15,17	0.88	0
5	MAN	M	7	5	11,11,12	0.86	0	15,15,17	1.82	4 (26%)
5	MAN	M	8	5	11,11,12	1.34	2 (18%)	15,15,17	1.95	3 (20%)
5	MAN	M	9	5	11,11,12	0.76	0	15,15,17	1.23	1 (6%)
3	NAG	N	1	3,1	14,14,15	0.70	0	17,19,21	1.38	1 (5%)
3	NAG	N	2	3	14,14,15	0.93	0	17,19,21	1.76	5 (29%)
3	BMA	N	3	3	11,11,12	0.88	0	15,15,17	1.49	2 (13%)
6	NAG	O	1	6,1	14,14,15	1.10	1 (7%)	17,19,21	1.78	6 (35%)
6	NAG	O	2	6	14,14,15	1.29	2 (14%)	17,19,21	1.79	5 (29%)
6	BMA	O	3	6	11,11,12	0.67	0	15,15,17	0.89	1 (6%)
6	MAN	O	4	6	11,11,12	1.16	2 (18%)	15,15,17	1.57	2 (13%)
6	MAN	O	5	6	11,11,12	0.66	0	15,15,17	1.30	1 (6%)
6	MAN	O	6	6	11,11,12	0.79	0	15,15,17	1.21	1 (6%)
6	MAN	O	7	6	11,11,12	0.91	0	15,15,17	1.74	5 (33%)
10	NAG	P	1	10,1	14,14,15	0.73	1 (7%)	17,19,21	1.89	2 (11%)
10	NAG	P	2	10	14,14,15	0.69	0	17,19,21	1.12	1 (5%)
10	BMA	P	3	10	11,11,12	0.97	0	15,15,17	3.46	6 (40%)
10	MAN	P	4	10	11,11,12	0.68	0	15,15,17	1.61	2 (13%)
10	MAN	P	5	10	11,11,12	0.64	0	15,15,17	1.87	3 (20%)
10	MAN	P	6	10	11,11,12	0.57	0	15,15,17	1.51	5 (33%)
10	MAN	P	7	10	11,11,12	0.90	0	15,15,17	1.70	3 (20%)
10	MAN	P	8	10	11,11,12	1.02	0	15,15,17	2.04	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	2/2/19/22	0/1/1/1
2	MAN	C	5	2	-	2/2/19/22	0/1/1/1
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
4	NAG	E	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	5/6/23/26	0/1/1/1
5	NAG	F	1	5,1	-	0/6/23/26	0/1/1/1
5	MAN	F	10	5	-	2/2/19/22	0/1/1/1
5	NAG	F	2	5	-	0/6/23/26	0/1/1/1
5	BMA	F	3	5	-	0/2/19/22	0/1/1/1
5	MAN	F	4	5	-	0/2/19/22	0/1/1/1
5	MAN	F	5	5	-	0/2/19/22	0/1/1/1
5	MAN	F	6	5	-	0/2/19/22	0/1/1/1
5	MAN	F	7	5	-	0/2/19/22	0/1/1/1
5	MAN	F	8	5	-	0/2/19/22	0/1/1/1
5	MAN	F	9	5	-	0/2/19/22	0/1/1/1
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	BMA	G	3	3	-	0/2/19/22	0/1/1/1
6	NAG	H	1	6,1	-	0/6/23/26	0/1/1/1
6	NAG	H	2	6	-	0/6/23/26	0/1/1/1
6	BMA	H	3	6	-	0/2/19/22	0/1/1/1
6	MAN	H	4	6	-	1/2/19/22	0/1/1/1
6	MAN	H	5	6	-	2/2/19/22	0/1/1/1
6	MAN	H	6	6	-	2/2/19/22	0/1/1/1
6	MAN	H	7	6	-	0/2/19/22	0/1/1/1
7	NAG	I	1	7,1	-	1/6/23/26	0/1/1/1
7	NAG	I	2	7	-	0/6/23/26	0/1/1/1
7	BMA	I	3	7	-	0/2/19/22	0/1/1/1
7	MAN	I	4	7	-	2/2/19/22	0/1/1/1
7	MAN	I	5	7	-	0/2/19/22	0/1/1/1
7	MAN	I	6	7	-	2/2/19/22	0/1/1/1
7	MAN	I	7	7	-	2/2/19/22	0/1/1/1
8	NAG	J	1	8,1	-	0/6/23/26	0/1/1/1
8	NAG	J	2	8	-	0/6/23/26	0/1/1/1
8	BMA	J	3	8	-	0/2/19/22	0/1/1/1
8	MAN	J	4	8	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	MAN	J	5	8	-	2/2/19/22	0/1/1/1
8	MAN	J	6	8	-	2/2/19/22	0/1/1/1
8	MAN	J	7	8	-	1/2/19/22	0/1/1/1
9	NAG	K	1	9,1	-	0/6/23/26	0/1/1/1
9	NAG	K	2	9	-	1/6/23/26	0/1/1/1
9	BMA	K	3	9	-	0/2/19/22	0/1/1/1
9	MAN	K	4	9	-	2/2/19/22	0/1/1/1
9	MAN	K	5	9	-	0/2/19/22	0/1/1/1
9	MAN	K	6	9	-	1/2/19/22	0/1/1/1
3	NAG	L	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	L	2	3	-	1/6/23/26	0/1/1/1
3	BMA	L	3	3	-	2/2/19/22	0/1/1/1
5	NAG	M	1	5,1	-	0/6/23/26	0/1/1/1
5	MAN	M	10	5	-	0/2/19/22	0/1/1/1
5	NAG	M	2	5	-	0/6/23/26	0/1/1/1
5	BMA	M	3	5	-	0/2/19/22	0/1/1/1
5	MAN	M	4	5	-	0/2/19/22	0/1/1/1
5	MAN	M	5	5	-	0/2/19/22	0/1/1/1
5	MAN	M	6	5	-	0/2/19/22	0/1/1/1
5	MAN	M	7	5	-	0/2/19/22	0/1/1/1
5	MAN	M	8	5	-	0/2/19/22	0/1/1/1
5	MAN	M	9	5	-	0/2/19/22	0/1/1/1
3	NAG	N	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	N	2	3	-	0/6/23/26	0/1/1/1
3	BMA	N	3	3	-	2/2/19/22	0/1/1/1
6	NAG	O	1	6,1	-	0/6/23/26	0/1/1/1
6	NAG	O	2	6	-	0/6/23/26	0/1/1/1
6	BMA	O	3	6	-	0/2/19/22	0/1/1/1
6	MAN	O	4	6	-	0/2/19/22	0/1/1/1
6	MAN	O	5	6	-	0/2/19/22	0/1/1/1
6	MAN	O	6	6	-	1/2/19/22	0/1/1/1
6	MAN	O	7	6	-	2/2/19/22	0/1/1/1
10	NAG	P	1	10,1	-	2/6/23/26	0/1/1/1
10	NAG	P	2	10	-	0/6/23/26	0/1/1/1
10	BMA	P	3	10	-	2/2/19/22	0/1/1/1
10	MAN	P	4	10	-	1/2/19/22	0/1/1/1
10	MAN	P	5	10	-	1/2/19/22	0/1/1/1
10	MAN	P	6	10	-	2/2/19/22	0/1/1/1
10	MAN	P	7	10	-	2/2/19/22	0/1/1/1
10	MAN	P	8	10	-	2/2/19/22	0/1/1/1

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	K	6	MAN	C2-C3	4.12	1.58	1.52
2	C	1	NAG	C1-C2	3.66	1.57	1.52
6	O	2	NAG	C2-N2	-3.53	1.40	1.46
5	M	8	MAN	C2-C3	3.28	1.57	1.52
2	C	4	MAN	C2-C3	3.22	1.57	1.52

The worst 5 of 253 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
10	P	3	BMA	C1-O5-C5	8.95	124.18	112.19
2	C	4	MAN	C1-O5-C5	8.40	123.44	112.19
3	D	3	BMA	C1-O5-C5	7.57	122.33	112.19
9	K	4	MAN	C1-O5-C5	7.46	122.18	112.19
6	H	2	NAG	C2-N2-C7	-7.23	113.21	122.90

There are no chirality outliers.

5 of 54 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	I	4	MAN	O5-C5-C6-O6
3	L	3	BMA	O5-C5-C6-O6
5	F	10	MAN	O5-C5-C6-O6
2	C	5	MAN	O5-C5-C6-O6
2	C	5	MAN	C4-C5-C6-O6

There are no ring outliers.

8 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	K	1	NAG	2	0
4	E	2	NAG	4	0
10	P	3	BMA	2	0
7	I	2	NAG	2	0
6	O	3	BMA	1	0
10	P	4	MAN	1	0
6	H	3	BMA	2	0
3	D	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain C



Oligosaccharide Chain D



Oligosaccharide Chain G



Oligosaccharide Chain L



Oligosaccharide Chain N



Oligosaccharide Chain E



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain F



Oligosaccharide Chain M



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain H



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain O



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain I



Oligosaccharide Chain J



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain K	
	
Bond lengths	**Bond angles**
	
Torsions	**Rings**

Oligosaccharide Chain P



5.6 Ligand geometry (i)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
12	MRD	A	941	-	7,7,7	0.83	0	9,10,10	0.66	0
11	NAG	A	939	1	14,14,15	0.59	0	17,19,21	1.83	5 (29%)
13	BGC	A	942	-	12,12,12	0.71	0	17,17,17	1.00	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
12	MRD	B	948	-	7,7,7	0.66	0	9,10,10	0.82	0
13	BGC	A	943	-	12,12,12	0.96	0	17,17,17	2.52	10 (58%)
11	NAG	B	908	1	14,14,15	0.79	0	17,19,21	1.76	6 (35%)
11	NAG	A	906	1	14,14,15	0.94	1 (7%)	17,19,21	1.41	3 (17%)
12	MRD	B	949	-	7,7,7	0.51	0	9,10,10	0.81	0
12	MRD	A	940	-	7,7,7	0.53	0	9,10,10	0.45	0
13	BGC	B	950	-	12,12,12	0.78	0	17,17,17	1.13	2 (11%)
13	BGC	B	951	-	12,12,12	1.04	1 (8%)	17,17,17	2.53	8 (47%)
11	NAG	B	947	1	14,14,15	0.75	0	17,19,21	1.24	2 (11%)
11	NAG	B	946	1	14,14,15	0.59	0	17,19,21	1.60	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	MRD	A	941	-	-	4/5/5/5	-
11	NAG	A	939	1	-	0/6/23/26	0/1/1/1
13	BGC	A	942	-	-	0/2/22/22	0/1/1/1
12	MRD	B	948	-	-	2/5/5/5	-
13	BGC	A	943	-	-	2/2/22/22	0/1/1/1
11	NAG	B	908	1	-	6/6/23/26	0/1/1/1
11	NAG	A	906	1	-	1/6/23/26	0/1/1/1
12	MRD	B	949	-	-	2/5/5/5	-
12	MRD	A	940	-	-	1/5/5/5	-
13	BGC	B	950	-	-	0/2/22/22	0/1/1/1
13	BGC	B	951	-	-	2/2/22/22	0/1/1/1
11	NAG	B	947	1	-	2/6/23/26	0/1/1/1
11	NAG	B	946	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	B	951	BGC	C1-C2	2.55	1.58	1.52
11	A	906	NAG	C1-C2	2.13	1.55	1.52

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
13	B	951	BGC	O5-C1-C2	4.87	118.86	110.30
13	A	943	BGC	O5-C1-C2	4.41	118.06	110.30
13	A	943	BGC	C3-C4-C5	-4.20	102.62	110.23
11	B	946	NAG	O5-C1-C2	-3.92	105.22	111.29
11	B	908	NAG	C8-C7-N2	3.57	122.03	116.12

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
12	A	941	MRD	C2-C3-C4-O4
12	A	941	MRD	C2-C3-C4-C5
12	B	949	MRD	C2-C3-C4-O4
12	B	949	MRD	C2-C3-C4-C5
13	A	943	BGC	O5-C5-C6-O6

There are no ring outliers.

9 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	A	941	MRD	6	0
13	A	942	BGC	1	0
12	B	948	MRD	1	0
13	A	943	BGC	1	0
11	B	908	NAG	1	0
12	B	949	MRD	1	0
12	A	940	MRD	2	0
13	B	950	BGC	1	0
13	B	951	BGC	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	834/841 (99%)	-0.38	11 (1%)	77	81	13, 22, 37, 67	0
1	B	832/841 (98%)	-0.52	1 (0%)	95	97	12, 19, 32, 62	0
All	All	1666/1682 (99%)	-0.45	12 (0%)	87	91	12, 20, 35, 67	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	573	LEU	3.1
1	A	110	VAL	3.0
1	A	95	LEU	2.8
1	A	109	GLY	2.4
1	A	144	ALA	2.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
9	MAN	K	6	11/12	0.49	0.25	69,83,89,91	0
3	BMA	G	3	11/12	0.54	0.31	79,91,96,96	0
6	MAN	O	7	11/12	0.69	0.29	72,78,86,87	0
2	MAN	C	4	11/12	0.70	0.23	70,78,89,90	0
7	MAN	I	7	11/12	0.77	0.25	51,63,68,69	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
9	MAN	K	4	11/12	0.79	0.25	73,77,80,83	0
10	MAN	P	7	11/12	0.79	0.26	51,61,66,77	0
3	BMA	N	3	11/12	0.84	0.24	66,74,79,83	0
2	MAN	C	5	11/12	0.84	0.17	55,60,69,70	0
10	MAN	P	8	11/12	0.84	0.25	66,72,81,81	0
3	BMA	D	3	11/12	0.86	0.23	40,48,60,62	0
3	NAG	G	2	14/15	0.86	0.17	52,64,80,85	0
7	MAN	I	6	11/12	0.87	0.22	56,60,68,69	0
3	BMA	L	3	11/12	0.87	0.30	61,76,90,93	0
6	MAN	H	7	11/12	0.88	0.27	62,64,74,76	0
9	MAN	K	5	11/12	0.90	0.34	80,84,89,92	0
5	MAN	M	10	11/12	0.90	0.14	39,47,51,55	0
6	BMA	H	3	11/12	0.91	0.17	32,38,45,56	0
10	MAN	P	6	11/12	0.91	0.22	51,59,66,74	0
7	BMA	I	3	11/12	0.91	0.14	37,44,51,52	0
2	BMA	C	3	11/12	0.91	0.12	43,56,61,62	0
8	MAN	J	7	11/12	0.92	0.11	36,39,42,42	0
10	BMA	P	3	11/12	0.92	0.18	36,44,55,61	0
9	BMA	K	3	11/12	0.92	0.13	49,54,62,75	0
6	MAN	H	4	11/12	0.92	0.15	34,36,43,44	0
5	MAN	F	10	11/12	0.92	0.15	37,42,54,57	0
7	MAN	I	4	11/12	0.93	0.13	30,36,44,58	0
6	BMA	O	3	11/12	0.93	0.17	31,37,44,52	0
3	NAG	N	2	14/15	0.93	0.18	41,47,57,60	0
6	NAG	O	2	14/15	0.93	0.17	23,30,37,38	0
6	NAG	H	2	14/15	0.94	0.13	21,29,35,41	0
3	NAG	G	1	14/15	0.94	0.09	29,37,50,54	0
7	NAG	I	2	14/15	0.95	0.21	30,32,42,51	0
6	NAG	H	1	14/15	0.95	0.12	22,28,42,46	0
3	NAG	L	2	14/15	0.95	0.14	21,32,46,50	0
5	MAN	M	6	11/12	0.95	0.14	30,33,36,36	0
10	NAG	P	2	14/15	0.95	0.15	25,33,41,43	0
3	NAG	D	2	14/15	0.95	0.14	26,33,36,42	0
10	MAN	P	4	11/12	0.95	0.21	25,38,53,57	0
8	BMA	J	3	11/12	0.95	0.08	26,31,33,35	0
8	MAN	J	5	11/12	0.95	0.20	33,42,54,77	0
7	NAG	I	1	14/15	0.95	0.11	24,26,32,34	0
5	MAN	F	5	11/12	0.96	0.10	23,27,36,40	0
9	NAG	K	2	14/15	0.96	0.10	25,29,34,41	0
5	MAN	F	6	11/12	0.96	0.18	34,35,44,44	0
5	MAN	F	7	11/12	0.96	0.10	19,24,26,26	0
3	NAG	D	1	14/15	0.96	0.07	22,27,30,36	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	NAG	M	2	14/15	0.96	0.09	17,22,28,33	0
7	MAN	I	5	11/12	0.96	0.13	35,37,44,48	0
6	NAG	O	1	14/15	0.96	0.09	19,28,41,42	0
3	NAG	N	1	14/15	0.96	0.09	26,36,45,53	0
10	MAN	P	5	11/12	0.96	0.12	32,36,39,44	0
4	NAG	E	2	14/15	0.96	0.13	30,37,45,51	0
8	MAN	J	4	11/12	0.96	0.13	30,34,40,41	0
6	MAN	O	4	11/12	0.96	0.14	31,34,38,41	0
6	MAN	H	6	11/12	0.97	0.15	27,30,38,42	0
2	NAG	C	1	14/15	0.97	0.08	22,25,28,28	0
5	NAG	F	2	14/15	0.97	0.11	25,27,29,29	0
5	BMA	F	3	11/12	0.97	0.09	23,25,27,30	0
2	NAG	C	2	14/15	0.97	0.09	25,33,43,46	0
10	NAG	P	1	14/15	0.97	0.10	19,23,26,28	0
5	BMA	M	3	11/12	0.97	0.08	22,24,25,28	0
6	MAN	O	5	11/12	0.97	0.11	32,33,36,46	0
6	MAN	O	6	11/12	0.97	0.10	25,31,35,37	0
5	MAN	M	5	11/12	0.97	0.10	25,26,32,32	0
8	MAN	J	6	11/12	0.97	0.15	34,36,41,50	0
6	MAN	H	5	11/12	0.97	0.12	26,28,39,51	0
9	NAG	K	1	14/15	0.97	0.09	20,25,27,27	0
5	MAN	M	9	11/12	0.98	0.10	25,27,29,35	0
4	NAG	E	1	14/15	0.98	0.07	22,24,26,30	0
5	NAG	M	1	14/15	0.98	0.09	21,25,27,27	0
8	NAG	J	1	14/15	0.98	0.08	15,18,21,22	0
8	NAG	J	2	14/15	0.98	0.07	18,23,28,34	0
5	NAG	F	1	14/15	0.98	0.10	24,27,31,31	0
5	MAN	F	4	11/12	0.98	0.09	19,21,23,24	0
5	MAN	M	4	11/12	0.98	0.06	18,19,21,23	0
5	MAN	F	8	11/12	0.98	0.10	24,26,28,29	0
5	MAN	F	9	11/12	0.98	0.11	29,32,34,35	0
5	MAN	M	7	11/12	0.98	0.07	16,19,19,21	0
5	MAN	M	8	11/12	0.98	0.09	19,21,22,28	0
3	NAG	L	1	14/15	0.99	0.06	15,19,23,25	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain D:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain G:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain L:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain N:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain E:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain F:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain M:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain O:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain I:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain J:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain K:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around Chain P:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
13	BGC	A	943	12/12	0.87	0.20	36,46,50,53	0
13	BGC	B	951	12/12	0.87	0.22	31,43,51,52	0
11	NAG	A	906	14/15	0.88	0.36	58,69,76,76	0
11	NAG	B	908	14/15	0.88	0.30	49,59,68,69	0
12	MRD	A	941	8/8	0.89	0.22	34,39,47,47	0
11	NAG	B	946	14/15	0.93	0.26	50,64,70,78	0
12	MRD	B	949	8/8	0.94	0.15	45,47,49,54	0
11	NAG	A	939	14/15	0.95	0.19	32,46,50,56	0
12	MRD	A	940	8/8	0.96	0.10	33,38,39,40	0
12	MRD	B	948	8/8	0.97	0.13	30,37,40,41	0
11	NAG	B	947	14/15	0.97	0.21	29,34,43,45	0
13	BGC	A	942	12/12	0.98	0.21	19,22,24,30	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
13	BGC	B	950	12/12	0.99	0.18	14,18,20,26	0

6.5 Other polymers (i)

There are no such residues in this entry.