

# Full wwPDB X-ray Structure Validation Report (i)

#### Sep 10, 2023 – 03:37 AM EDT

PDB ID : 4IIN

Title: Crystal structure of a putative 3-oxoacyl-[acyl-carrier protein] reductase from

Helicobacter pylori 26695 complexed with NAD+

Authors: Hou, J.; Osinski, T.; Zheng, H.; Shumilin, I.; Shabalin, I.; Shatsman, S.;

Anderson, W.F.; Minor, W.; Center for Structural Genomics of Infectious

Diseases (CSGID)

Deposited on : 2012-12-20

Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$ 

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4: 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1




## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	271	80%	7%	13%
1	В	271	81%	5%	14%
1	С	271	80%	7%	12%
1	D	271	81%	6%	13%



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7321 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 3-ketoacyl-acyl carrier protein reductase (FabG).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	236	Total	С	N	О	S	0	0	0
1	A	230	1737	1095	297	336	9	0	U	
1	В	234	Total	С	N	О	S	0	0	0
1	Б	234	1731	1092	296	334	9	0	U	
1	C	238	Total	С	N	О	S	0	0	0
1		230	1753	1105	299	340	9	0	U	
1	D	236	Total	С	N	О	S	0	0	0
1	ש	∠30	1742	1100	298	335	9		U	

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	expression tag	UNP O25286
A	-22	HIS	-	expression tag	UNP O25286
A	-21	HIS	-	expression tag	UNP O25286
A	-20	HIS	-	expression tag	UNP O25286
A	-19	HIS	-	expression tag	UNP O25286
A	-18	HIS	-	expression tag	UNP O25286
A	-17	HIS	-	expression tag	UNP O25286
A	-16	SER	-	expression tag	UNP O25286
A	-15	SER	-	expression tag	UNP O25286
A	-14	GLY	-	expression tag	UNP O25286
A	-13	VAL	-	expression tag	UNP O25286
A	-12	ASP	-	expression tag	UNP O25286
A	-11	LEU	-	expression tag	UNP O25286
A	-10	GLY	-	expression tag	UNP O25286
A	-9	THR	_	expression tag	UNP O25286
A	-8	GLU	-	expression tag	UNP O25286
A	-7	ASN		expression tag	UNP O25286
A	-6	LEU	-	expression tag	UNP O25286
A	-5	TYR	-	expression tag	UNP O25286
A	-4	PHE	-	expression tag	UNP O25286
A	-3	GLN	_	expression tag	UNP O25286



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP O25286
A	-1	ASN	_	expression tag	UNP O25286
A	0	ALA	-	expression tag	UNP O25286
В	-23	MET	-	expression tag	UNP O25286
В	-22	HIS	-	expression tag	UNP O25286
В	-21	HIS	-	expression tag	UNP O25286
В	-20	HIS	-	expression tag	UNP O25286
В	-19	HIS	-	expression tag	UNP O25286
В	-18	HIS	-	expression tag	UNP O25286
В	-17	HIS	-	expression tag	UNP O25286
В	-16	SER	-	expression tag	UNP O25286
В	-15	SER	-	expression tag	UNP O25286
В	-14	GLY	-	expression tag	UNP O25286
В	-13	VAL	-	expression tag	UNP O25286
В	-12	ASP	-	expression tag	UNP O25286
В	-11	LEU	-	expression tag	UNP O25286
В	-10	GLY	-	expression tag	UNP O25286
В	-9	THR	-	expression tag	UNP O25286
В	-8	GLU	-	expression tag	UNP O25286
В	-7	ASN	-	expression tag	UNP O25286
В	-6	LEU	-	expression tag	UNP O25286
В	-5	TYR	_	expression tag	UNP O25286
В	-4	PHE	-	expression tag	UNP O25286
В	-3	GLN	-	expression tag	UNP O25286
В	-2	SER	_	expression tag	UNP O25286
В	-1	ASN	_	expression tag	UNP O25286
В	0	ALA	-	expression tag	UNP O25286
С	-23	MET	-	expression tag	UNP O25286
С	-22	HIS	-	expression tag	UNP O25286
С	-21	HIS	-	expression tag	UNP O25286
С	-20	HIS	-	expression tag	UNP O25286
С	-19	HIS	-	expression tag	UNP O25286
С	-18	HIS	-	expression tag	UNP O25286
С	-17	HIS	-	expression tag	UNP O25286
С	-16	SER	-	expression tag	UNP O25286
С	-15	SER	-	expression tag	UNP O25286
С	-14	GLY	-	expression tag	UNP O25286
С	-13	VAL	-	expression tag	UNP O25286
С	-12	ASP	-	expression tag	UNP O25286
С	-11	LEU	-	expression tag	UNP O25286
С	-10	GLY	-	expression tag	UNP O25286
С	-9	THR	-	expression tag	UNP O25286




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	-8	GLU	-	expression tag	UNP O25286
С	-7	ASN	-	expression tag	UNP O25286
С	-6	LEU	-	expression tag	UNP O25286
С	-5	TYR	_	expression tag	UNP O25286
С	-4	PHE	-	expression tag	UNP O25286
С	-3	GLN	-	expression tag	UNP O25286
С	-2	SER	-	expression tag	UNP O25286
С	-1	ASN	-	expression tag	UNP O25286
С	0	ALA	-	expression tag	UNP O25286
D	-23	MET	-	expression tag	UNP O25286
D	-22	HIS	-	expression tag	UNP O25286
D	-21	HIS	-	expression tag	UNP O25286
D	-20	HIS	-	expression tag	UNP O25286
D	-19	HIS	-	expression tag	UNP O25286
D	-18	HIS	-	expression tag	UNP O25286
D	-17	HIS	-	expression tag	UNP O25286
D	-16	SER	-	expression tag	UNP O25286
D	-15	SER	-	expression tag	UNP O25286
D	-14	GLY	-	expression tag	UNP O25286
D	-13	VAL	-	expression tag	UNP O25286
D	-12	ASP	-	expression tag	UNP O25286
D	-11	LEU	-	expression tag	UNP O25286
D	-10	GLY	-	expression tag	UNP O25286
D	-9	THR	-	expression tag	UNP O25286
D	-8	GLU	-	expression tag	UNP O25286
D	-7	ASN	-	expression tag	UNP O25286
D	-6	LEU	-	expression tag	UNP O25286
D	-5	TYR	-	expression tag	UNP O25286
D	-4	PHE	-	expression tag	UNP O25286
D	-3	GLN	-	expression tag	UNP O25286
D	-2	SER	-	expression tag	UNP O25286
D	-1	ASN	-	expression tag	UNP O25286
D	0	ALA	-	expression tag	UNP O25286

• Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).






Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Λ	1	Total	С	N	О	Р	0	0
2	Λ	1	44	21	7	14	2	U	
2	В	1	Total	С	N	О	Р	0	0
	Б	1	44	21	7	14	2	U	0
2	С	1	Total	С	N	О	Р	0	0
		1	44	21	7	14	2	U	U
9	D	1	Total	С	N	О	Р	0	0
	ש	1	44	21	7	14	2	U	0

• Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

$\mathbf{N}$	$\mathbf{lol}$	Chain	Residues	Atoms	ZeroOcc	AltConf
	3	A	1	Total K 1 1	0	0
	3	В	1	Total K 1 1	0	0

 $\bullet$  Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $\mathrm{C_2H_3O_2}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total C O 4 2 2	0	0
4	С	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

#### • Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	42	Total O 42 42	0	0
5	В	39	Total O 39 39	0	0
5	С	35	Total O 35 35	0	0
5	D	52	Total O 52 52	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3-ketoacyl-acyl carrier protein reductase (FabG)







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	128.92Å 112.09Å 70.66Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.87 - 2.40	Depositor
Resolution (A)	43.83 - 2.38	EDS
% Data completeness	93.4 (43.87-2.40)	Depositor
(in resolution range)	92.1 (43.83-2.38)	EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	7.21 (at 2.39Å)	Xtriage
Refinement program	REFMAC	Depositor
P. P.	0.217 , 0.255	Depositor
$R, R_{free}$	0.220 , $0.257$	DCC
$R_{free}$ test set	1934 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.5	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33, 34.9	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.45, < L^2>=0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	7321	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, K, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	$\mid \text{RMSZ} \mid \# Z  > 5 \mid$		RMSZ	# Z >5	
1	A	0.67	0/1759	0.73	1/2370 (0.0%)	
1	В	0.71	0/1753	0.75	0/2360	
1	С	0.65	0/1775	0.71	0/2391	
1	D	0.70	0/1764	0.71	0/2376	
All	All	0.68	0/7051	0.73	1/9497 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

$\mathbf{Mol}$	Chain	$\operatorname{Res}$	Type	${f Atoms}$	$\mathbf{Z}$	$Observed(^{o})$	$ \operatorname{Ideal}(^o) $
1	A	179	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	204	TYR	Peptide



## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1737	0	1719	13	0
1	В	1731	0	1722	8	0
1	С	1753	0	1740	11	0
1	D	1742	0	1727	9	0
2	A	44	0	26	5	0
2	В	44	0	26	3	0
2	С	44	0	26	1	0
2	D	44	0	26	4	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	В	4	0	3	1	0
4	С	4	0	3	0	0
4	D	4	0	3	0	0
5	A	42	0	0	0	0
5	В	39	0	0	0	0
5	С	35	0	0	0	0
5	D	52	0	0	1	0
All	All	7321	0	7021	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:16:GLY:HA2	1:C:190:THR:HG21	1.81	0.61
1:A:17:ILE:HB	2:A:301:NAD:H51N	1.82	0.59
1:D:141:ALA:O	2:D:301:NAD:H6N	2.06	0.56
1:A:37:ARG:NH2	1:A:113:ASN:OD1	2.43	0.52
1:A:90:ASN:OD1	2:A:301:NAD:H4D	2.09	0.51
1:B:74:ILE:HG21	1:B:128:VAL:HG21	1.92	0.51
1:B:74:ILE:CG2	1:B:128:VAL:HG21	2.41	0.51
1:D:62:PHE:HB2	1:D:69:ASP:HB3	1.93	0.50
1:B:62:PHE:HB2	1:B:69:ASP:HB3	1.94	0.50
1:C:189:GLU:O	1:C:190:THR:C	2.51	0.50



Continued from previous page...

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap(Å)
1:D:0:ALA:N	5:D:405:HOH:O	2.41	0.49
1:C:62:PHE:HB2	1:C:69:ASP:HB3	1.94	0.49
1:A:62:PHE:HB2	1:A:69:ASP:HB3	1.96	0.48
2:B:301:NAD:C4N	4:B:303:ACT:C	2.92	0.47
1:A:74:ILE:HG21	1:A:128:VAL:HG21	1.96	0.47
1:A:74:ILE:CG2	1:A:128:VAL:HG21	2.44	0.47
1:C:74:ILE:CG2	1:C:128:VAL:HG21	2.45	0.47
1:C:74:ILE:HG21	1:C:128:VAL:HG21	1.98	0.46
1:A:141:ALA:O	2:A:301:NAD:H6N	2.16	0.45
2:C:301:NAD:O1N	2:C:301:NAD:N7N	2.48	0.45
1:A:35:ASN:OD1	2:A:301:NAD:H2A	2.17	0.45
1:A:155:TYR:HH	2:A:301:NAD:HO2N	1.65	0.43
1:D:90:ASN:OD1	2:D:301:NAD:H4D	2.18	0.43
1:B:17:ILE:HB	2:B:301:NAD:H51N	2.00	0.43
1:C:126:LEU:O	1:C:130:SER:HB3	2.19	0.42
1:D:185:PRO:HB2	2:D:301:NAD:C5N	2.49	0.42
1:A:130:SER:OG	1:A:176:ARG:NH1	2.43	0.42
1:B:37:ARG:HG3	2:B:301:NAD:C2A	2.50	0.42
1:C:10:ILE:HD12	1:C:32:VAL:CG1	2.50	0.42
1:D:126:LEU:O	1:D:130:SER:HB3	2.20	0.42
1:C:207:ASN:O	1:C:209:PRO:HD3	2.20	0.41
1:D:183:VAL:HG12	1:D:185:PRO:HD3	2.02	0.41
1:A:130:SER:CB	1:A:176:ARG:HH11	2.32	0.41
1:A:217:LYS:O	1:A:221:GLU:HG3	2.20	0.41
1:B:217:LYS:O	1:B:221:GLU:HG3	2.21	0.41
1:D:17:ILE:HB	2:D:301:NAD:H51N	2.03	0.41
1:A:102:MET:O	1:B:123:ARG:NH1	2.44	0.41
1:C:144:ILE:HD12	1:C:149:ASN:HB2	2.03	0.41
1:C:39:ASN:ND2	1:C:42:VAL:HG23	2.37	0.40
1:D:144:ILE:HD12	1:D:149:ASN:HB2	2.03	0.40
1:C:217:LYS:O	1:C:221:GLU:HG3	2.21	0.40
1:B:39:ASN:ND2	1:B:42:VAL:HG23	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Chain Analysed Favoured		Allowed	Outliers	Perce	ntiles
1	A	232/271~(86%)	226 (97%)	6 (3%)	0	100	100
1	В	230/271 (85%)	223 (97%)	7 (3%)	0	100	100
1	C	234/271 (86%)	228 (97%)	6 (3%)	0	100	100
1	D	232/271~(86%)	224 (97%)	7 (3%)	1 (0%)	34	48
All	All	928/1084 (86%)	901 (97%)	26 (3%)	1 (0%)	51	68

#### All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
1	D	205	VAL	

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outlie		Percentiles	
1	A	177/219 (81%)	174 (98%)	3 (2%)	60	78
1	В	177/219 (81%)	174 (98%)	3 (2%)	60	78
1	C	179/219 (82%)	176 (98%)	3 (2%)	60	78
1	D	$176/219 \; (80\%)$	173 (98%)	3 (2%)	60	78
All	All	709/876 (81%)	697 (98%)	12 (2%)	60	78

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\operatorname{Res}$	Type
1	A	70	PHE
1	A	85	SER
1	A	132	SER
1	В	38	SER
1	В	70	PHE



Continued from previous page...

Mol	Chain	Res	Type
1	В	85	SER
1	С	70	PHE
1	С	85	SER
1	С	190	THR
1	D	56	LYS
1	D	70	PHE
1	D	85	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chain I		Chain Dog Lie		Chain Res Link			Bond lengths		Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2		
2	NAD	С	301	-	42,48,48	1.05	5 (11%)	50,73,73	1.32	7 (14%)		
4	ACT	D	302	-	3,3,3	0.64	0	3,3,3	1.30	0		
4	ACT	С	302	-	3,3,3	0.82	0	3,3,3	0.88	0		



Mol Type		Chain Res Link			Bo	Bond lengths			Bond angles		
MIOI	Type   Chain   Res	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2		
4	ACT	В	303	-	3,3,3	0.83	0	3,3,3	0.52	0	
2	NAD	A	301	-	42,48,48	1.14	5 (11%)	50,73,73	1.29	6 (12%)	
2	NAD	В	301	-	42,48,48	1.04	4 (9%)	50,73,73	1.52	9 (18%)	
2	NAD	D	301	-	42,48,48	0.96	3 (7%)	50,73,73	1.57	6 (12%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	С	301	-	-	5/26/62/62	0/5/5/5
2	NAD	В	301	-	-	6/26/62/62	0/5/5/5
2	NAD	D	301	-	-	9/26/62/62	0/5/5/5
2	NAD	A	301	-	-	10/26/62/62	0/5/5/5

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(\AA)$	Ideal(Å)
2	A	301	NAD	O4B-C1B	3.08	1.45	1.41
2	A	301	NAD	C2A-N3A	2.73	1.36	1.32
2	В	301	NAD	C5A-C4A	2.59	1.47	1.40
2	A	301	NAD	C5A-C4A	2.53	1.47	1.40
2	С	301	NAD	O2D-C2D	2.52	1.48	1.43
2	С	301	NAD	O4D-C1D	2.49	1.44	1.41
2	С	301	NAD	C5A-C4A	2.47	1.47	1.40
2	С	301	NAD	C2A-N3A	2.43	1.36	1.32
2	D	301	NAD	C5A-C4A	2.39	1.47	1.40
2	В	301	NAD	C2D-C1D	-2.29	1.50	1.53
2	D	301	NAD	C2A-N3A	2.18	1.35	1.32
2	С	301	NAD	O4B-C1B	2.18	1.44	1.41
2	В	301	NAD	C2N-N1N	-2.12	1.32	1.35
2	D	301	NAD	O4D-C1D	2.07	1.44	1.41
2	A	301	NAD	O4D-C1D	2.03	1.43	1.41
2	A	301	NAD	C7N-N7N	2.02	1.36	1.33
2	В	301	NAD	C5A-N7A	-2.00	1.32	1.39

All (28) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
2	D	301	NAD	C1B-N9A-C4A	-4.75	118.30	126.64
2	В	301	NAD	O4B-C1B-C2B	-4.59	100.21	106.93
2	D	301	NAD	N3A-C2A-N1A	-4.21	122.10	128.68
2	С	301	NAD	N3A-C2A-N1A	-4.09	122.29	128.68
2	D	301	NAD	C3N-C7N-N7N	3.92	122.46	117.75
2	A	301	NAD	N3A-C2A-N1A	-3.75	122.81	128.68
2	В	301	NAD	N6A-C6A-N1A	3.62	126.08	118.57
2	A	301	NAD	O7N-C7N-C3N	-3.47	115.48	119.63
2	В	301	NAD	C5A-C6A-N6A	-3.02	115.76	120.35
2	A	301	NAD	C4A-C5A-N7A	-2.91	106.37	109.40
2	D	301	NAD	O7N-C7N-C3N	-2.80	116.28	119.63
2	A	301	NAD	C3N-C7N-N7N	2.80	121.11	117.75
2	D	301	NAD	C2A-N1A-C6A	2.79	123.52	118.75
2	С	301	NAD	C1B-N9A-C4A	-2.69	121.91	126.64
2	В	301	NAD	C6N-N1N-C2N	2.68	124.42	121.97
2	С	301	NAD	C3D-C2D-C1D	-2.59	97.08	100.98
2	D	301	NAD	C6N-N1N-C2N	-2.43	119.76	121.97
2	В	301	NAD	C3N-C7N-N7N	2.36	120.58	117.75
2	В	301	NAD	N3A-C2A-N1A	-2.30	125.08	128.68
2	В	301	NAD	C3D-C2D-C1D	-2.29	97.53	100.98
2	С	301	NAD	N6A-C6A-N1A	2.26	123.27	118.57
2	В	301	NAD	C3N-C2N-N1N	-2.18	118.30	120.43
2	С	301	NAD	O2D-C2D-C1D	2.13	118.73	110.85
2	С	301	NAD	O2B-C2B-C1B	2.13	118.70	110.85
2	A	301	NAD	O5D-C5D-C4D	2.12	116.28	108.99
2	С	301	NAD	C3B-C2B-C1B	-2.11	97.80	100.98
2	A	301	NAD	O2D-C2D-C1D	2.09	118.56	110.85
2	В	301	NAD	O7N-C7N-C3N	-2.01	117.23	119.63

There are no chirality outliers.

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	NAD	C5D-O5D-PN-O1N
2	A	301	NAD	C5D-O5D-PN-O2N
2	A	301	NAD	O4D-C1D-N1N-C2N
2	В	301	NAD	C5D-O5D-PN-O1N
2	В	301	NAD	C5D-O5D-PN-O2N
2	В	301	NAD	O4D-C1D-N1N-C2N
2	С	301	NAD	C5D-O5D-PN-O1N
2	С	301	NAD	C5D-O5D-PN-O2N
2	D	301	NAD	C5D-O5D-PN-O1N
2	D	301	NAD	C5D-O5D-PN-O2N



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	D	301	NAD	O4D-C1D-N1N-C2N
2	D	301	NAD	O4D-C1D-N1N-C6N
2	A	301	NAD	O4B-C4B-C5B-O5B
2	A	301	NAD	C3B-C4B-C5B-O5B
2	A	301	NAD	C4B-C5B-O5B-PA
2	A	301	NAD	PN-O3-PA-O1A
2	A	301	NAD	PN-O3-PA-O2A
2	С	301	NAD	O4B-C4B-C5B-O5B
2	С	301	NAD	PA-O3-PN-O2N
2	В	301	NAD	O4B-C4B-C5B-O5B
2	D	301	NAD	O4B-C4B-C5B-O5B
2	A	301	NAD	C5D-O5D-PN-O3
2	В	301	NAD	C5D-O5D-PN-O3
2	С	301	NAD	C5D-O5D-PN-O3
2	D	301	NAD	C5D-O5D-PN-O3
2	D	301	NAD	C2D-C1D-N1N-C2N
2	D	301	NAD	C2D-C1D-N1N-C6N
2	A	301	NAD	PA-O3-PN-O1N
2	В	301	NAD	PA-O3-PN-O1N
2	D	301	NAD	PA-O3-PN-O2N

There are no ring outliers.

5 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	301	NAD	1	0
4	В	303	ACT	1	0
2	A	301	NAD	5	0
2	В	301	NAD	3	0
2	D	301	NAD	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient




equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\# \mathrm{RSRZ} {>} 2$	$OWAB(Å^2)$	Q < 0.9
1	A	236/271 (87%)	-0.21	1 (0%) 92 91	11, 23, 46, 58	0
1	В	$234/271 \ (86\%)$	-0.32	1 (0%) 92 91	8, 20, 39, 59	0
1	С	238/271 (87%)	-0.28	1 (0%) 92 91	10, 21, 42, 67	0
1	D	$236/271 \ (87\%)$	-0.22	2 (0%) 86 84	10, 20, 40, 59	0
All	All	944/1084 (87%)	-0.26	5 (0%) 91 89	8, 21, 42, 67	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	38	SER	3.1
1	D	190	THR	2.9
1	С	130	SER	2.3
1	В	38	SER	2.1
1	A	190	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, $95^{th}$ percentile and maximum values of B factors of atoms in the group.	The column
labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.	

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	NAD	A	301	44/44	0.78	0.32	31,57,85,92	0
4	ACT	D	302	4/4	0.83	0.25	33,34,36,36	0
4	ACT	В	303	4/4	0.87	0.24	30,37,37,39	0
2	NAD	С	301	44/44	0.92	0.17	20,38,54,56	0
2	NAD	В	301	44/44	0.95	0.12	15,19,24,26	0
4	ACT	С	302	4/4	0.95	0.10	35,36,36,37	0
2	NAD	D	301	44/44	0.95	0.12	21,28,32,35	0
3	K	A	302	1/1	0.98	0.07	17,17,17,17	0
3	K	В	302	1/1	0.99	0.10	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








# 

#### Electron density around NAD B 301:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











# 6.5 Other polymers (i)

There are no such residues in this entry.

