



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 21, 2023 – 12:26 AM EDT

PDB ID : 5IOL  
Title : Crystal structure of Nucleoside Diphosphate Kinase from Schistosoma mansoni  
Authors : Torini, J.R.S.; Romanello, L.; Bird, L.E.; Nettleship, J.E.; Owens, R.J.; Aller, P.; DeMarco, R.; Brandao-Neto, J.; Pereira, H.M.  
Deposited on : 2016-03-08  
Resolution : 1.74 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	 93% 7%
1	B	150	 91% 8%
1	C	150	 93% 6%
1	D	150	 90% 10%
1	E	150	 93% 7%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	150	 91% 9%
1	G	150	 95% . .
1	H	150	 97% .
1	I	150	 97% . .
1	J	150	 94% 5% .
1	K	150	 94% 5% .
1	L	150	 97% .

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 16239 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	149	1189	752	209	219	9	0	2	0
1	B	149	1183	748	211	215	9	0	1	0
1	C	149	1179	745	209	216	9	0	0	0
1	D	150	1183	747	210	217	9	0	0	0
1	E	149	1179	745	209	216	9	0	0	0
1	F	150	1191	752	213	217	9	0	1	0
1	G	149	1179	745	209	216	9	0	0	0
1	H	150	1183	747	210	217	9	0	0	0
1	I	149	1179	745	209	216	9	0	0	0
1	J	149	1187	750	212	216	9	0	1	0
1	K	149	1179	745	209	216	9	0	0	0
1	L	150	1183	747	210	217	9	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP G4VJY9
A	1	PRO	-	expression tag	UNP G4VJY9
B	0	GLY	-	expression tag	UNP G4VJY9
B	1	PRO	-	expression tag	UNP G4VJY9
C	0	GLY	-	expression tag	UNP G4VJY9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	PRO	-	expression tag	UNP G4VJY9
D	0	GLY	-	expression tag	UNP G4VJY9
D	1	PRO	-	expression tag	UNP G4VJY9
E	0	GLY	-	expression tag	UNP G4VJY9
E	1	PRO	-	expression tag	UNP G4VJY9
F	0	GLY	-	expression tag	UNP G4VJY9
F	1	PRO	-	expression tag	UNP G4VJY9
G	0	GLY	-	expression tag	UNP G4VJY9
G	1	PRO	-	expression tag	UNP G4VJY9
H	0	GLY	-	expression tag	UNP G4VJY9
H	1	PRO	-	expression tag	UNP G4VJY9
I	0	GLY	-	expression tag	UNP G4VJY9
I	1	PRO	-	expression tag	UNP G4VJY9
J	0	GLY	-	expression tag	UNP G4VJY9
J	1	PRO	-	expression tag	UNP G4VJY9
K	0	GLY	-	expression tag	UNP G4VJY9
K	1	PRO	-	expression tag	UNP G4VJY9
L	0	GLY	-	expression tag	UNP G4VJY9
L	1	PRO	-	expression tag	UNP G4VJY9

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	135	Total O 135 135	0	0
2	B	153	Total O 153 153	0	0
2	C	176	Total O 176 176	0	0
2	D	169	Total O 169 169	0	0
2	E	172	Total O 172 172	0	0
2	F	182	Total O 182 182	0	0
2	G	187	Total O 187 187	0	0
2	H	178	Total O 178 178	0	0
2	I	172	Total O 172 172	0	0
2	J	204	Total O 204 204	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	K	162	Total 162	O 162	0	0
2	L	155	Total 155	O 155	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleoside diphosphate kinase





- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase



- Molecule 1: Nucleoside diphosphate kinase




- Molecule 1: Nucleoside diphosphate kinase

Chain F:  91% 9%



- Molecule 1: Nucleoside diphosphate kinase

Chain G:  95%



- Molecule 1: Nucleoside diphosphate kinase

Chain H:  97%



- Molecule 1: Nucleoside diphosphate kinase

Chain I:  97%



- Molecule 1: Nucleoside diphosphate kinase

Chain J:  94% 5%



- Molecule 1: Nucleoside diphosphate kinase

Chain K:  94% 5%



- Molecule 1: Nucleoside diphosphate kinase

Chain L:  97%





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	211.76Å 82.27Å 122.37Å 90.00° 104.89° 90.00°	Depositor
Resolution (Å)	59.13 – 1.74 59.13 – 1.74	Depositor EDS
% Data completeness (in resolution range)	98.2 (59.13-1.74) 98.2 (59.13-1.74)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 1.74Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.163 , 0.191 0.164 , 0.191	Depositor DCC
$R_{free}$ test set	10268 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.0	Xtrriage
Anisotropy	0.345	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 43.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	16239	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/1222	0.47	0/1649
1	B	0.38	0/1213	0.49	0/1636
1	C	0.29	0/1206	0.46	0/1627
1	D	0.29	0/1210	0.48	0/1633
1	E	0.28	0/1206	0.47	0/1627
1	F	0.29	0/1221	0.47	0/1647
1	G	0.29	0/1206	0.47	0/1627
1	H	0.30	0/1210	0.48	0/1633
1	I	0.29	0/1206	0.49	0/1627
1	J	0.31	0/1217	0.50	0/1641
1	K	0.29	0/1206	0.48	0/1627
1	L	0.29	0/1210	0.48	0/1633
All	All	0.30	0/14533	0.48	0/19607

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1189	0	1185	6	0
1	B	1183	0	1179	8	0
1	C	1179	0	1172	6	0
1	D	1183	0	1175	12	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1179	0	1172	9	0
1	F	1191	0	1188	8	0
1	G	1179	0	1172	4	0
1	H	1183	0	1175	4	0
1	I	1179	0	1172	2	0
1	J	1187	0	1185	6	0
1	K	1179	0	1172	6	0
1	L	1183	0	1175	2	0
2	A	135	0	0	2	0
2	B	153	0	0	5	0
2	C	176	0	0	4	0
2	D	169	0	0	8	0
2	E	172	0	0	5	0
2	F	182	0	0	4	0
2	G	187	0	0	1	0
2	H	178	0	0	2	0
2	I	172	0	0	1	0
2	J	204	0	0	3	0
2	K	162	0	0	2	0
2	L	155	0	0	1	0
All	All	16239	0	14122	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:46:GLN:NE2	2:D:202:HOH:O	2.06	0.88
1:D:98:PRO:O	2:D:201:HOH:O	1.94	0.84
1:D:97:CYS:SG	2:L:342:HOH:O	2.37	0.81
1:A:53:LYS:NZ	2:A:201:HOH:O	2.17	0.78
1:E:92:LYS:HA	1:E:109:VAL:HG22	1.69	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/150 (99%)	145 (97%)	3 (2%)	1 (1%)	22	8
1	B	148/150 (99%)	144 (97%)	3 (2%)	1 (1%)	22	8
1	C	147/150 (98%)	143 (97%)	3 (2%)	1 (1%)	22	8
1	D	148/150 (99%)	144 (97%)	3 (2%)	1 (1%)	22	8
1	E	147/150 (98%)	143 (97%)	3 (2%)	1 (1%)	22	8
1	F	149/150 (99%)	145 (97%)	3 (2%)	1 (1%)	22	8
1	G	147/150 (98%)	144 (98%)	2 (1%)	1 (1%)	22	8
1	H	148/150 (99%)	145 (98%)	2 (1%)	1 (1%)	22	8
1	I	147/150 (98%)	143 (97%)	3 (2%)	1 (1%)	22	8
1	J	148/150 (99%)	144 (97%)	3 (2%)	1 (1%)	22	8
1	K	147/150 (98%)	143 (97%)	3 (2%)	1 (1%)	22	8
1	L	148/150 (99%)	144 (97%)	3 (2%)	1 (1%)	22	8
All	All	1773/1800 (98%)	1727 (97%)	34 (2%)	12 (1%)	22	8

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	113	VAL
1	B	113	VAL
1	C	113	VAL
1	L	113	VAL
1	D	113	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/129 (102%)	130 (99%)	1 (1%)	81	72
1	B	129/129 (100%)	128 (99%)	1 (1%)	81	72
1	C	129/129 (100%)	127 (98%)	2 (2%)	62	44
1	D	129/129 (100%)	127 (98%)	2 (2%)	62	44
1	E	129/129 (100%)	129 (100%)	0	100	100
1	F	130/129 (101%)	129 (99%)	1 (1%)	81	72
1	G	129/129 (100%)	129 (100%)	0	100	100
1	H	129/129 (100%)	129 (100%)	0	100	100
1	I	129/129 (100%)	129 (100%)	0	100	100
1	J	130/129 (101%)	130 (100%)	0	100	100
1	K	129/129 (100%)	128 (99%)	1 (1%)	81	72
1	L	129/129 (100%)	129 (100%)	0	100	100
All	All	1552/1548 (100%)	1544 (100%)	8 (0%)	88	83

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	149	GLU
1	F	134	GLN
1	D	118	ASP
1	C	149	GLU
1	D	134	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	39	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	149/150 (99%)	-0.31	1 (0%) 87 91	17, 25, 45, 53	0
1	B	149/150 (99%)	-0.33	1 (0%) 87 91	15, 24, 44, 58	0
1	C	149/150 (99%)	-0.30	0 100 100	16, 25, 40, 47	0
1	D	150/150 (100%)	-0.33	2 (1%) 77 82	14, 22, 43, 49	0
1	E	149/150 (99%)	-0.29	1 (0%) 87 91	16, 25, 40, 51	0
1	F	150/150 (100%)	-0.23	0 100 100	16, 23, 40, 58	0
1	G	149/150 (99%)	-0.29	0 100 100	14, 21, 35, 52	0
1	H	150/150 (100%)	-0.17	2 (1%) 77 82	13, 21, 40, 62	0
1	I	149/150 (99%)	-0.40	0 100 100	14, 20, 35, 47	0
1	J	149/150 (99%)	-0.21	0 100 100	13, 19, 35, 46	0
1	K	149/150 (99%)	-0.38	0 100 100	17, 23, 42, 53	0
1	L	150/150 (100%)	-0.34	1 (0%) 87 91	14, 21, 43, 54	0
All	All	1792/1800 (99%)	-0.30	8 (0%) 92 94	13, 22, 42, 62	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	0	GLY	3.5
1	B	90	ALA	2.6
1	H	1	PRO	2.6
1	D	91	THR	2.3
1	L	57	PHE	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.