



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2023 – 12:46 AM JST

PDB ID : 6ION
Title : The complex of C4.4A with its antibody 11H10 Fab fragment
Authors : Huang, M.D.; Jiang, Y.B.; Yuan, C.; Lin, L.
Deposited on : 2018-10-30
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	215	
2	H	220	
3	A	201	
4	B	2	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4603 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti-C4.4A antibody 11H10, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	1630	1015	274	335	6	0	0	0

- Molecule 2 is a protein called anti-C4.4A antibody 11H10, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	208	1575	1010	256	305	4	0	0	0

- Molecule 3 is a protein called Ly6/PLAUR domain-containing protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	185	1342	814	241	268	19	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	B	2	28	16	2	10	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	A	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: anti-C4.4A antibody 11H10, light chain



- Molecule 2: anti-C4.4A antibody 11H10, heavy chain



- Molecule 3: Ly6/PLAUR domain-containing protein 3





- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B: 100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.86Å 63.96Å 107.71Å 90.00° 96.10° 90.00°	Depositor
Resolution (Å)	40.14 – 2.75 40.14 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.0 (40.14-2.75) 99.0 (40.14-2.75)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.77Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.197 , 0.255 0.196 , 0.256	Depositor DCC
R_{free} test set	926 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	54.2	Xtrriage
Anisotropy	0.712	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4603	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.57	0/1669	0.88	4/2273 (0.2%)
2	H	0.59	1/1621 (0.1%)	0.98	3/2228 (0.1%)
3	A	0.72	2/1362 (0.1%)	1.11	8/1844 (0.4%)
All	All	0.62	3/4652 (0.1%)	0.99	15/6345 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	116	VAL	CB-CG2	-5.32	1.41	1.52
3	A	78	CYS	CB-SG	-5.11	1.73	1.81
3	A	83	CYS	CB-SG	-5.10	1.73	1.81

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	69	LEU	CB-CG-CD2	-11.17	92.01	111.00
3	A	171	ARG	CG-CD-NE	-7.70	95.64	111.80
3	A	78	CYS	CA-CB-SG	-7.47	100.56	114.00
1	L	182	LEU	CA-CB-CG	6.50	130.24	115.30
3	A	83	CYS	CA-CB-SG	-6.36	102.55	114.00
2	H	52	HIS	N-CA-CB	6.23	121.82	110.60
1	L	39	ARG	CG-CD-NE	-6.05	99.09	111.80
1	L	182	LEU	CB-CG-CD2	-6.01	100.78	111.00
3	A	87	LEU	CA-CB-CG	5.61	128.20	115.30
1	L	105	LEU	CB-CG-CD1	-5.59	101.50	111.00
3	A	18	LYS	CA-CB-CG	-5.38	101.56	113.40
2	H	120	LYS	CA-CB-CG	-5.30	101.74	113.40
3	A	161	ARG	NE-CZ-NH2	-5.24	117.68	120.30
2	H	179	LEU	CB-CG-CD2	5.18	119.81	111.00
3	A	82	ARG	NE-CZ-NH1	-5.09	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1630	0	1523	75	0
2	H	1575	0	1513	44	0
3	A	1342	0	1271	41	0
4	B	28	0	25	2	0
5	A	28	0	26	0	0
All	All	4603	0	4358	155	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (155) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:109:ARG:HG2	1:L:141:TYR:CD2	1.80	1.17
1:L:191:ASN:ND2	1:L:213:ASN:ND2	1.98	1.11
1:L:109:ARG:HG2	1:L:141:TYR:HD2	1.13	1.09
1:L:191:ASN:HD21	1:L:213:ASN:ND2	1.56	1.03
1:L:191:ASN:ND2	1:L:213:ASN:HD21	1.56	1.00
1:L:141:TYR:HD1	1:L:142:PRO:HA	1.27	0.98
1:L:141:TYR:CD1	1:L:142:PRO:HA	1.99	0.96
1:L:141:TYR:HD1	1:L:142:PRO:CA	1.80	0.93
1:L:36:TYR:HE1	1:L:89:GLN:HG2	1.41	0.86
1:L:109:ARG:CG	1:L:141:TYR:HD2	1.87	0.85
1:L:191:ASN:HD21	1:L:213:ASN:HD22	1.21	0.84
1:L:119:PHE:HB2	1:L:134:VAL:HG23	1.61	0.83
1:L:36:TYR:CE1	1:L:89:GLN:HG2	2.17	0.78
1:L:134:VAL:HG21	2:H:129:LEU:HD13	1.66	0.78
2:H:126:VAL:HG11	2:H:211:VAL:HG11	1.68	0.76
1:L:8:PRO:HG2	1:L:11:LEU:HD13	1.68	0.74
3:A:147:VAL:HG12	3:A:181:LEU:HD13	1.68	0.74
3:A:4:TYR:HB3	3:A:19:MET:HE2	1.69	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:47:LEU:HD13	3:A:156:VAL:HG12	1.70	0.73
2:H:168:VAL:HG22	2:H:186:VAL:HG12	1.71	0.73
1:L:191:ASN:CG	1:L:213:ASN:ND2	2.42	0.73
2:H:99:ARG:HH21	2:H:101:ARG:HD2	1.54	0.73
3:A:62:ARG:CZ	3:A:177:PRO:HD3	2.19	0.73
1:L:58:ILE:HD12	1:L:59:PRO:HD2	1.73	0.70
1:L:107:ILE:H	1:L:167:GLN:HE22	1.38	0.70
2:H:88:ASP:O	2:H:116:VAL:HG21	1.92	0.70
1:L:12:SER:HB2	1:L:106:GLU:OE2	1.94	0.68
3:A:147:VAL:HG13	3:A:160:VAL:HG21	1.76	0.67
1:L:49:LYS:HE3	1:L:53:GLU:OE1	1.94	0.67
1:L:49:LYS:HB2	1:L:55:ILE:HD11	1.75	0.67
2:H:142:THR:HG22	2:H:187:THR:HG23	1.76	0.67
3:A:111:TYR:CE1	3:A:190:ARG:HB3	2.31	0.66
2:H:11:ILE:O	2:H:12:LEU:HD23	1.97	0.65
2:H:92:THR:OG1	2:H:116:VAL:HG22	1.97	0.65
3:A:149:LEU:HD12	3:A:179:PHE:HD2	1.62	0.65
4:B:2:NAG:O7	4:B:2:NAG:O3	2.14	0.64
2:H:49:TRP:HE1	2:H:52:HIS:HD2	1.45	0.63
2:H:52:HIS:CE1	2:H:60:TYR:HB2	2.34	0.63
3:A:135:SER:O	3:A:137:HIS:HD2	1.81	0.63
1:L:90:GLN:HE21	1:L:98:THR:HB	1.63	0.61
1:L:58:ILE:HD12	1:L:59:PRO:CD	2.30	0.61
3:A:34:ALA:HB2	3:A:75:LEU:HD12	1.81	0.61
2:H:89:THR:HA	2:H:116:VAL:CG2	2.31	0.61
3:A:149:LEU:HD12	3:A:179:PHE:CD2	2.36	0.61
1:L:34:HIS:HE1	1:L:91:SER:HB3	1.67	0.59
1:L:141:TYR:CD1	1:L:142:PRO:CA	2.68	0.58
2:H:11:ILE:HD11	2:H:152:PRO:HG3	1.85	0.58
2:H:49:TRP:NE1	2:H:52:HIS:HD2	2.00	0.57
3:A:24:CYS:HB3	3:A:28:VAL:CG2	2.34	0.57
3:A:108:VAL:CG2	3:A:140:LYS:HG2	2.35	0.57
4:B:1:NAG:H5	4:B:2:NAG:O5	2.04	0.57
1:L:85:ASP:OD2	1:L:104:LYS:HE2	2.05	0.56
2:H:37:GLY:HA2	2:H:52:HIS:HA	1.87	0.56
3:A:35:VAL:HG12	3:A:87:LEU:HD13	1.88	0.56
1:L:61:ARG:HD2	1:L:77:SER:O	2.06	0.56
3:A:108:VAL:HG23	3:A:140:LYS:HG2	1.87	0.56
2:H:99:ARG:NH2	2:H:101:ARG:HD2	2.21	0.56
1:L:49:LYS:HG2	1:L:50:TYR:CD1	2.41	0.56
2:H:148:LYS:HA	2:H:181:THR:HG23	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:62:ARG:NH2	3:A:177:PRO:HD3	2.21	0.56
2:H:89:THR:HA	2:H:116:VAL:HG23	1.88	0.56
1:L:95:PRO:O	2:H:52:HIS:NE2	2.36	0.55
1:L:141:TYR:HD1	1:L:142:PRO:N	2.04	0.55
2:H:18:LEU:O	2:H:83:LYS:HA	2.06	0.55
2:H:101:ARG:O	2:H:103:ARG:N	2.36	0.55
1:L:55:ILE:O	1:L:58:ILE:HG22	2.08	0.54
2:H:124:PRO:HB3	2:H:150:TYR:HB3	1.88	0.54
3:A:25:ALA:O	3:A:28:VAL:HG22	2.08	0.54
1:L:109:ARG:HG3	1:L:110:ALA:O	2.09	0.53
2:H:126:VAL:HG13	2:H:213:LYS:HD2	1.91	0.52
3:A:83:CYS:SG	3:A:84:ASN:N	2.80	0.52
3:A:8:GLN:HB3	3:A:48:ALA:HB3	1.90	0.52
3:A:24:CYS:HB3	3:A:28:VAL:HG23	1.90	0.52
2:H:116:VAL:HG23	2:H:116:VAL:O	2.09	0.52
3:A:1:LEU:HD12	3:A:81:ASP:OD1	2.10	0.52
2:H:49:TRP:CZ2	2:H:52:HIS:CD2	2.97	0.52
3:A:135:SER:O	3:A:137:HIS:CD2	2.62	0.52
1:L:13:VAL:HG21	1:L:19:VAL:HG21	1.92	0.51
3:A:19:MET:HE3	3:A:50:ARG:HD2	1.91	0.51
1:L:50:TYR:OH	2:H:103:ARG:HD2	2.11	0.50
1:L:4:MET:SD	1:L:90:GLN:HB3	2.51	0.50
2:H:99:ARG:O	2:H:105:PHE:HA	2.11	0.50
2:H:150:TYR:CE1	2:H:155:VAL:HG13	2.46	0.50
2:H:88:ASP:OD2	2:H:89:THR:N	2.40	0.50
1:L:124:GLU:O	1:L:127:THR:OG1	2.29	0.49
1:L:185:ASP:O	1:L:189:ARG:HG3	2.12	0.49
3:A:9:LYS:HG2	3:A:158:LEU:HD21	1.95	0.49
2:H:49:TRP:HZ2	2:H:52:HIS:CD2	2.31	0.49
3:A:33:GLU:HG3	3:A:87:LEU:HD12	1.94	0.48
1:L:109:ARG:CD	1:L:141:TYR:HD2	2.27	0.48
3:A:7:VAL:CG2	3:A:156:VAL:HG21	2.42	0.48
3:A:62:ARG:NE	3:A:177:PRO:HD3	2.29	0.48
1:L:184:LYS:O	1:L:188:GLU:HG2	2.14	0.48
2:H:2:VAL:HG13	2:H:27:PHE:HD1	1.78	0.48
1:L:58:ILE:CD1	1:L:59:PRO:HD2	2.41	0.48
2:H:160:ASN:HD22	2:H:164:LEU:HD13	1.78	0.47
1:L:21:PHE:O	1:L:72:THR:HA	2.14	0.47
1:L:36:TYR:HE1	1:L:89:GLN:CG	2.20	0.47
1:L:119:PHE:HB2	1:L:134:VAL:CG2	2.36	0.47
1:L:182:LEU:HD22	1:L:186:GLU:HB3	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:148:THR:HG23	3:A:157:SER:HB3	1.97	0.47
1:L:95:PRO:HB2	2:H:52:HIS:HE1	1.79	0.47
2:H:53:ILE:HD13	2:H:73:LYS:HG2	1.97	0.47
1:L:193:TYR:HB2	1:L:210:PHE:CE1	2.50	0.47
2:H:104:TYR:CD1	2:H:104:TYR:N	2.83	0.47
3:A:5:SER:HA	3:A:50:ARG:O	2.15	0.47
1:L:147:VAL:C	1:L:148:LYS:HD2	2.36	0.46
2:H:127:TYR:HD1	2:H:146:LEU:CD2	2.27	0.46
3:A:112:SER:O	3:A:127:PRO:HD2	2.15	0.46
1:L:188:GLU:HA	1:L:212:ARG:CZ	2.45	0.46
1:L:191:ASN:OD1	1:L:213:ASN:ND2	2.49	0.46
3:A:33:GLU:HG3	3:A:87:LEU:CD1	2.47	0.45
1:L:191:ASN:CG	1:L:213:ASN:HD21	2.08	0.45
3:A:145:GLY:HA3	3:A:183:GLY:HA3	1.99	0.45
1:L:28:ASN:OD1	1:L:68:GLY:HA2	2.17	0.45
1:L:95:PRO:HB2	2:H:52:HIS:CE1	2.52	0.45
1:L:109:ARG:NH1	1:L:110:ALA:O	2.50	0.45
1:L:187:TYR:CZ	1:L:212:ARG:HD3	2.52	0.45
3:A:17:ASN:O	3:A:18:LYS:HG3	2.16	0.45
1:L:13:VAL:HG21	1:L:19:VAL:CG2	2.46	0.44
1:L:50:TYR:O	1:L:51:ALA:HB3	2.18	0.44
1:L:137:LEU:HD23	1:L:145:ILE:CD1	2.47	0.44
3:A:34:ALA:CB	3:A:75:LEU:HD12	2.46	0.44
3:A:147:VAL:CG1	3:A:181:LEU:HD13	2.42	0.44
1:L:3:LEU:HD22	1:L:4:MET:N	2.33	0.43
1:L:151:ILE:HD11	1:L:180:LEU:HD21	2.00	0.43
1:L:26:ASN:N	1:L:26:ASN:OD1	2.51	0.43
3:A:196:ARG:HH11	3:A:196:ARG:HD3	1.67	0.43
1:L:89:GLN:HB3	1:L:99:PHE:CD2	2.53	0.43
2:H:2:VAL:HG13	2:H:27:PHE:CD1	2.53	0.42
2:H:49:TRP:HE1	2:H:52:HIS:CD2	2.32	0.42
1:L:65:SER:OG	1:L:72:THR:HG22	2.19	0.42
1:L:105:LEU:HD12	1:L:105:LEU:HA	1.66	0.42
1:L:34:HIS:O	1:L:88:CYS:HA	2.20	0.42
1:L:21:PHE:CD1	1:L:73:LEU:HD23	2.55	0.41
1:L:58:ILE:HD12	1:L:58:ILE:HA	1.72	0.41
3:A:24:CYS:HB3	3:A:28:VAL:HG21	2.01	0.41
2:H:83:LYS:HB2	2:H:83:LYS:HE3	1.84	0.41
2:H:59:LYS:HD3	2:H:71:ILE:O	2.20	0.41
1:L:13:VAL:HG12	1:L:106:GLU:O	2.21	0.41
1:L:19:VAL:HG21	1:L:78:VAL:HG21	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:83:ILE:HG23	1:L:105:LEU:O	2.21	0.41
1:L:89:GLN:HE22	2:H:104:TYR:HA	1.86	0.41
3:A:116:LEU:HB3	3:A:120:ALA:HB3	2.03	0.41
1:L:13:VAL:O	1:L:107:ILE:HA	2.21	0.41
3:A:143:PHE:O	3:A:161:ARG:HA	2.21	0.41
2:H:104:TYR:HE1	3:A:137:HIS:CE1	2.38	0.41
2:H:159:TRP:HZ3	2:H:215:ILE:HD13	1.87	0.40
2:H:195:SER:OG	2:H:196:GLU:N	2.54	0.40
1:L:139:ASN:HA	1:L:173:THR:CB	2.51	0.40
3:A:12:ASP:O	3:A:15:SER:HB3	2.20	0.40
3:A:114:VAL:O	3:A:114:VAL:HG12	2.22	0.40
1:L:49:LYS:HG2	1:L:50:TYR:HD1	1.85	0.40
1:L:33:ILE:HG22	1:L:51:ALA:HA	2.03	0.40
1:L:139:ASN:HA	1:L:173:THR:HB	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	211/215 (98%)	204 (97%)	7 (3%)	0	100	100
2	H	204/220 (93%)	198 (97%)	4 (2%)	2 (1%)	15	27
3	A	181/201 (90%)	168 (93%)	13 (7%)	0	100	100
All	All	596/636 (94%)	570 (96%)	24 (4%)	2 (0%)	41	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	102	LEU
2	H	15	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	185/192 (96%)	177 (96%)	8 (4%)	29	48
2	H	176/191 (92%)	167 (95%)	9 (5%)	24	41
3	A	151/164 (92%)	145 (96%)	6 (4%)	31	51
All	All	512/547 (94%)	489 (96%)	23 (4%)	27	46

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	8	PRO
1	L	14	SER
1	L	141	TYR
1	L	148	LYS
1	L	156	ARG
1	L	169	SER
1	L	177	SER
1	L	200	LYS
2	H	19	SER
2	H	48	GLU
2	H	67	SER
2	H	83	LYS
2	H	101	ARG
2	H	161	SER
2	H	208	SER
2	H	212	ASP
2	H	213	LYS
3	A	15	SER
3	A	78	CYS
3	A	149	LEU
3	A	169	CYS
3	A	197	ASN
3	A	200	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	34	HIS
1	L	167	GLN
1	L	213	ASN
2	H	160	ASN
3	A	137	HIS
3	A	187	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	1	4,3	14,14,15	1.52	2 (14%)	17,19,21	1.82	2 (11%)
4	NAG	B	2	4	14,14,15	0.90	1 (7%)	17,19,21	0.99	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1	4,3	-	2/6/23/26	0/1/1/1
4	NAG	B	2	4	-	4/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1	NAG	C1-C2	4.24	1.58	1.52
4	B	1	NAG	O5-C1	3.63	1.49	1.43
4	B	2	NAG	C1-C2	2.88	1.56	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1	NAG	C1-O5-C5	6.65	121.20	112.19
4	B	2	NAG	C1-O5-C5	2.92	116.15	112.19
4	B	1	NAG	O4-C4-C5	2.51	115.52	109.30

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1	NAG	O5-C5-C6-O6
4	B	2	NAG	C4-C5-C6-O6
4	B	2	NAG	O5-C5-C6-O6
4	B	1	NAG	C4-C5-C6-O6
4	B	2	NAG	C1-C2-N2-C7
4	B	2	NAG	C3-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	2	NAG	2	0
4	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	301	3	14,14,15	1.83	1 (7%)	17,19,21	1.24	3 (17%)
5	NAG	A	302	3	14,14,15	0.84	1 (7%)	17,19,21	1.45	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	301	3	-	1/6/23/26	0/1/1/1
5	NAG	A	302	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	301	NAG	O5-C1	-5.69	1.34	1.43
5	A	302	NAG	O5-C1	2.92	1.48	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	302	NAG	C1-O5-C5	5.10	119.11	112.19
5	A	301	NAG	C1-O5-C5	2.39	115.43	112.19
5	A	301	NAG	O3-C3-C2	2.16	113.93	109.47
5	A	301	NAG	C1-C2-N2	2.12	114.10	110.49

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	301	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	213/215 (99%)	0.73	25 (11%) 4 5	37, 89, 140, 172	0
2	H	208/220 (94%)	0.76	28 (13%) 3 3	34, 75, 137, 162	0
3	A	185/201 (92%)	0.49	10 (5%) 25 31	36, 57, 97, 132	0
All	All	606/636 (95%)	0.67	63 (10%) 6 7	34, 70, 135, 172	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	164	LEU	7.7
1	L	116	VAL	6.8
2	H	191	SER	5.3
1	L	213	ASN	4.2
2	H	217	PRO	4.1
2	H	192	THR	4.0
3	A	25	ALA	4.0
2	H	144	GLY	4.0
3	A	200	TYR	4.0
2	H	131	PRO	3.8
3	A	26	PRO	3.8
1	L	118	ILE	3.7
1	L	117	SER	3.7
3	A	27	GLY	3.7
1	L	145	ILE	3.6
1	L	203	THR	3.6
2	H	193	TRP	3.5
1	L	185	ASP	3.4
2	H	141	VAL	3.3
2	H	208	SER	3.3
2	H	189	PRO	3.2
2	H	140	MET	3.1
1	L	119	PHE	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	29	ASP	3.1
1	L	144	ASP	3.0
2	H	205	PRO	3.0
1	L	201	THR	2.9
1	L	197	ALA	2.8
1	L	155	GLU	2.8
2	H	162	GLY	2.8
2	H	213	LYS	2.7
2	H	188	VAL	2.7
1	L	195	CYS	2.7
2	H	215	ILE	2.7
1	L	127	THR	2.6
2	H	161	SER	2.5
1	L	179	THR	2.5
1	L	147	VAL	2.4
2	H	165	SER	2.4
2	H	143	LEU	2.4
3	A	152	ALA	2.3
1	L	133	VAL	2.3
2	H	2	VAL	2.3
1	L	182	LEU	2.3
2	H	190	SER	2.3
1	L	176	MET	2.2
3	A	199	THR	2.2
1	L	207	VAL	2.2
2	H	198	VAL	2.2
1	L	136	PHE	2.2
1	L	123	SER	2.2
2	H	163	SER	2.2
2	H	168	VAL	2.1
1	L	11	LEU	2.1
2	H	206	ALA	2.1
2	H	145	CYS	2.1
1	L	191	ASN	2.1
3	A	12	ASP	2.1
1	L	150	LYS	2.0
3	A	28	VAL	2.0
2	H	90	ALA	2.0
3	A	23	LYS	2.0
2	H	124	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	2	14/15	0.79	0.34	90,99,108,111	0
4	NAG	B	1	14/15	0.93	0.22	54,70,86,93	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	301	14/15	0.76	0.28	95,105,111,111	0
5	NAG	A	302	14/15	0.86	0.21	52,87,93,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.