



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 30, 2024 – 10:27 PM EST

PDB ID : 1IPP  
Title : HOMING ENDONUCLEASE/DNA COMPLEX  
Authors : Flick, K.E.; Jurica, M.S.; Monnat Jr., R.J.; Stoddard, B.L.  
Deposited on : 1998-03-19  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	21	
1	D	21	
2	A	163	
2	B	163	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*TP\*TP\*GP\*AP\*CP\*TP\*CP\*TP\*CP\*TP\*TP\*AP\*AP\*GP\*AP\*GP\*AP\*GP\*TP\*CP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	21	Total	C	N	O	P	0	0	0
			427	206	76	125	20			
1	D	21	Total	C	N	O	P	0	0	0
			427	206	76	125	20			

- Molecule 2 is a protein called INTRON-ENCODED ENDONUCLEASE I-PPOI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	162	Total	C	N	O	S	0	0	0
			1245	786	232	219	8			
2	B	162	Total	C	N	O	S	0	0	0
			1245	786	232	219	8			

- Molecule 3 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cd	0	0
			2	2		
3	B	2	Total	Cd	0	0
			2	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	73	Total O 73 73	0	0
5	D	69	Total O 69 69	0	0
5	A	202	Total O 202 202	0	0
5	B	203	Total O 203 203	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DNA (5'-D(\*TP\*TP\*GP\*AP\*CP\*TP\*CP\*TP\*CP\*TP\*TP\*AP\*AP\*GP\*AP\*GP\*AP\*GP\*TP\*CP\*A)-3')

Chain C: 



- Molecule 1: DNA (5'-D(\*TP\*TP\*GP\*AP\*CP\*TP\*CP\*TP\*CP\*TP\*TP\*AP\*AP\*GP\*AP\*GP\*AP\*GP\*TP\*CP\*A)-3')

Chain D: 




- Molecule 2: INTRON-ENCODED ENDONUCLEASE I-PPOI

Chain A: 



- Molecule 2: INTRON-ENCODED ENDONUCLEASE I-PPOI

Chain B: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.90Å 113.90Å 88.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.20	Depositor
% Data completeness (in resolution range)	90.2 (50.00-2.20)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	4.80	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, $R_{free}$	0.212 , 0.316	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3897	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	C	3.30	2/477 (0.4%)	1.75	2/732 (0.3%)
1	D	3.07	2/477 (0.4%)	1.67	3/732 (0.4%)
2	A	0.58	0/1286	0.78	0/1760
2	B	0.59	0/1286	0.81	1/1760 (0.1%)
All	All	1.73	4/3526 (0.1%)	1.15	6/4984 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	214	DG	O5'-C5'	70.05	3.17	1.42
1	D	214	DG	O5'-C5'	64.57	3.03	1.42
1	D	214	DG	P-O5'	-7.42	1.52	1.59
1	C	214	DG	P-O5'	-6.81	1.52	1.59

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	214	DG	P-O5'-C5'	-39.67	57.42	120.90
1	D	214	DG	P-O5'-C5'	-36.51	62.49	120.90
1	D	214	DG	O5'-C5'-C4'	15.01	148.52	111.00
1	C	214	DG	O5'-C5'-C4'	14.46	147.16	111.00
1	D	214	DG	O5'-P-OP2	5.69	117.53	110.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	210	DT	Sidechain
1	C	217	DA	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	427	0	241	21	0
1	D	427	0	240	18	0
2	A	1245	0	1191	74	0
2	B	1245	0	1191	89	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	202	0	0	25	0
5	B	203	0	0	31	0
5	C	73	0	0	4	0
5	D	69	0	0	9	0
All	All	3897	0	2863	178	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 178 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:9:LEU:HD22	2:B:157:ARG:HH11	1.13	1.13
2:B:95:THR:HG23	2:B:119:ASN:HD22	1.19	1.04
1:C:219:DT:H2''	1:C:220:DC:H5''	1.41	1.02
2:B:95:THR:CG2	2:B:119:ASN:HD22	1.77	0.96
1:D:214:DG:H2''	5:D:223:HOH:O	1.66	0.94

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	160/163 (98%)	152 (95%)	7 (4%)	1 (1%)	25	26
2	B	160/163 (98%)	152 (95%)	7 (4%)	1 (1%)	25	26
All	All	320/326 (98%)	304 (95%)	14 (4%)	2 (1%)	25	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	3	LEU
2	B	52	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	133/134 (99%)	114 (86%)	19 (14%)	3	2
2	B	133/134 (99%)	115 (86%)	18 (14%)	4	3
All	All	266/268 (99%)	229 (86%)	37 (14%)	3	3

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	83	LEU
2	B	144	LEU
2	B	84	LEU
2	B	99	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	A	93	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	156	GLN
2	B	141	GLN
2	B	27	HIS
2	B	129	ASN
2	A	155	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	1
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	213:DA	O3'	214:DG	P	4.01
1	D	213:DA	O3'	214:DG	P	3.88

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.