



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 03:11 am BST

PDB ID : 2IQ5  
Title : Unliganded Crystal Structure of the Uridine Phosphorylase from Salmonella Typhimurium at 1.90 Å Resolution  
Authors : Timofeev, V.I.; Dontsova, M.V.; Gabdoulkhakov, A.G.; Pavlyuk, B.P.; Mikhailov, A.M.  
Deposited on : 2006-10-13  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	 5% 88% 10%
1	B	252	 8% 71% 17% 11%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3723 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	1874	1171	330	361	12	0	0	0
1	B	225	1674	1050	291	322	11	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	89	Total 89	O 89	0	0
2	B	86	Total 86	O 86	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	151.66 Å 151.66 Å 47.92 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.90 28.66 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (10.00-1.90) 99.5 (28.66-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 1.91 Å)	Xtrriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.201 , 0.247 0.202 , 0.249	Depositor DCC
$R_{free}$ test set	1602 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.1	Xtrriage
Anisotropy	0.535	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 54.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3723	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1904	0.61	0/2581
1	B	0.47	0/1699	0.60	0/2303
All	All	0.47	0/3603	0.60	0/4884

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1874	0	1872	12	0
1	B	1674	0	1679	27	0
2	A	89	0	0	0	0
2	B	86	0	0	7	0
All	All	3723	0	3551	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1034:ILE:CG1	1:A:1064:VAL:HG11	2.13	0.77

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1081:LEU:HG	1:B:1086:ILE:HD13	1.73	0.70
1:A:1144:ALA:HA	1:A:1244:ILE:HD12	1.76	0.67
1:B:1004:SER:N	2:B:1337:HOH:O	2.28	0.67
1:A:1240:HIS:O	1:A:1244:ILE:HG12	1.95	0.65
1:B:1239:SER:O	1:B:1243:LYS:HG3	1.98	0.64
1:B:1048:ARG:HD2	2:B:1282:HOH:O	2.05	0.56
1:B:1158:SER:HB3	1:B:1200:ALA:HB2	1.89	0.54
1:B:1147:ILE:HD13	1:B:1147:ILE:H	1.72	0.53
1:B:1047:HIS:HE1	2:B:1318:HOH:O	1.92	0.52
1:B:1155:VAL:HG22	2:B:1305:HOH:O	2.10	0.52
1:B:1184:MET:C	1:B:1186:GLU:H	2.13	0.52
1:B:1142:GLU:HB2	2:B:1326:HOH:O	2.12	0.48
1:B:1240:HIS:HA	1:B:1243:LYS:HE2	1.96	0.48
1:B:1163:TYR:HB2	1:B:1164:PRO:HD3	1.94	0.48
1:B:1155:VAL:HG23	1:B:1192:VAL:HA	1.96	0.47
1:A:1012:THR:HG22	1:A:1015:ASP:CG	2.34	0.47
1:B:1008:HIS:HD2	1:B:1080:GLU:OE2	1.97	0.47
1:B:1247:GLU:O	1:B:1251:ARG:HG2	2.14	0.47
1:B:1167:GLU:HG2	1:B:1184:MET:SD	2.54	0.47
1:A:1201:THR:O	1:A:1205:MET:HG2	2.17	0.45
1:B:1078:VAL:HG21	1:B:1202:LEU:HD23	1.99	0.45
1:A:1038:MET:HG2	1:A:1057:LEU:HD13	2.00	0.44
1:B:1063:ILE:HG21	1:B:1086:ILE:HD11	1.98	0.44
1:B:1095:THR:HG23	2:B:1324:HOH:O	2.17	0.43
1:A:1025:PRO:O	1:A:1066:SER:HA	2.18	0.43
1:B:1225:GLN:HB3	2:B:1319:HOH:O	2.19	0.43
1:A:1030:ARG:NH1	1:A:1238:GLU:OE1	2.51	0.43
1:A:1230:ASN:HB2	1:A:1233:THR:OG1	2.19	0.42
1:A:1049:GLU:HG3	1:B:1049:GLU:CD	2.40	0.42
1:B:1240:HIS:ND1	1:B:1241:ALA:N	2.67	0.42
1:B:1096:GLY:HA2	1:B:1221:VAL:O	2.20	0.42
1:B:1094:THR:HB	1:B:1220:ILE:HD12	2.01	0.41
1:B:1161:THR:HB	1:B:1164:PRO:HD2	2.01	0.41
1:A:1007:PHE:CD1	1:A:1007:PHE:N	2.89	0.41
1:B:1100:PRO:HA	1:B:1224:THR:HB	2.03	0.41
1:A:1007:PHE:HZ	1:A:1083:GLN:HE22	1.69	0.40
1:B:1009:LEU:HD23	1:B:1081:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/252 (98%)	242 (98%)	5 (2%)	1 (0%)	34	24
1	B	219/252 (87%)	213 (97%)	4 (2%)	2 (1%)	17	7
All	All	467/504 (93%)	455 (97%)	9 (2%)	3 (1%)	25	15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1163	TYR
1	B	1166	GLN
1	B	1185	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/201 (98%)	189 (96%)	7 (4%)	35	26
1	B	176/201 (88%)	166 (94%)	10 (6%)	20	11
All	All	372/402 (92%)	355 (95%)	17 (5%)	27	17

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1007	PHE
1	A	1037	LEU
1	A	1048	ARG

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1084	LEU
1	A	1178	ARG
1	A	1186	GLU
1	A	1196	GLU
1	B	1011	LEU
1	B	1027	ASP
1	B	1081	LEU
1	B	1092	ILE
1	B	1145	LYS
1	B	1147	ILE
1	B	1196	GLU
1	B	1202	LEU
1	B	1224	THR
1	B	1238	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1083	GLN
1	A	1122	HIS
1	A	1226	GLN
1	B	1008	HIS
1	B	1047	HIS
1	B	1188	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/252 (99%)	0.44	12 (4%) 30 33	20, 29, 52, 60	0
1	B	225/252 (89%)	0.63	20 (8%) 9 11	18, 29, 56, 88	0
All	All	475/504 (94%)	0.53	32 (6%) 17 20	18, 29, 54, 88	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1225	GLN	13.0
1	B	1168	ARG	11.0
1	B	1167	GLU	9.5
1	B	1163	TYR	8.1
1	A	1007	PHE	8.0
1	A	1230	ASN	7.4
1	B	1166	GLN	7.0
1	B	1184	MET	6.6
1	B	1169	TYR	6.5
1	B	1162	PHE	4.7
1	B	1237	THR	4.5
1	B	1239	SER	4.5
1	B	1223	ARG	4.5
1	B	1164	PRO	4.4
1	B	1224	THR	4.4
1	A	1044	LEU	4.2
1	A	1231	ALA	4.2
1	B	1186	GLU	3.9
1	A	1101	HIS	3.8
1	A	1233	THR	3.4
1	B	1238	GLU	3.2
1	A	1232	GLU	3.1
1	A	1048	ARG	3.0
1	B	1195	TYR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	1014	ASN	2.7
1	B	1222	ASN	2.6
1	A	1234	MET	2.5
1	B	1220	ILE	2.2
1	A	1033	LYS	2.2
1	A	1238	GLU	2.2
1	B	1004	SER	2.1
1	B	1165	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.