

# wwPDB X-ray Structure Validation Summary Report (i)

#### Aug 27, 2023 – 06:41 AM EDT

PDB ID : 3IQ6

Title: Crystal structure of a tetrameric Zn-bound cytochrome cb562 complex with

covalently and non-covalently stabilized interfaces

Authors: Brodin, J.N.; Tezcan, F.A.

Deposited on : 2009-08-19

Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution

Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	106	69%	28%	<del>.</del>
1	В	106	69%	29%	•
1	С	106	68%	31%	<del>.</del>
1	D	106	74%	25%	•
1	Е	106	75%	23%	<del>-</del>



Continued from previous page...

Mol	Chain	Length	Quality of chai	n
1	F	106	58%	41%
1	G	106	7% 65%	34%
1	Н	106	70%	30%



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Soluble cytochrome b562.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace						
1	A	106	Total	С	N	О	S	0	0	0						
1	A	100	818	505	145	162	6	0	0	0						
1	В	106	Total	С	N	О	S	0	0	0						
1	Б	100	818	505	145	162	6	0	0	0						
1	С	106	Total	С	N	О	S	0	0	0						
1		100	818	505	145	162	6	0		0						
1	D	106	Total	С	N	О	S	0	0	0						
1	D		818	505	145	162	6		U							
1	Е	106	Total	С	N	О	S	0	0	0						
1	15	100	818	505	145	162	6	0		0						
1	F	106	Total	С	N	О	S	0	0	0						
1	Г	I.	I.	I.	Г	Г	Г	100	818	505	145	162	6	0	U	U
1	1 G	C	C	C	С	106	Total	С	N	О	S	0	0	0		
1		G 100	818	505	145	162	6	0		0						
1	1 H	106	Total	С	N	О	S	0	0	0						
1		106	818	505	145	162	6			U						

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	ALA	ARG	engineered mutation	UNP P0ABE7
A	38	ALA	LEU	engineered mutation	UNP P0ABE7
A	41	TRP	GLN	engineered mutation	UNP P0ABE7
A	42	SER	LYS	engineered mutation	UNP P0ABE7
A	59	HIS	LYS	engineered mutation	UNP P0ABE7
A	66	TRP	ASP	engineered mutation	UNP P0ABE7
A	69	ILE	VAL	engineered mutation	UNP P0ABE7
A	73	HIS	ASP	engineered mutation	UNP P0ABE7
A	77	HIS	LYS	engineered mutation	UNP P0ABE7
A	96	CYS	THR	engineered mutation	UNP P0ABE7
A	98	CYS	ARG	engineered mutation	UNP P0ABE7
A	101	CYS	TYR	engineered mutation	UNP P0ABE7
В	34	ALA	ARG	engineered mutation	UNP P0ABE7



Continued from previous page...

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
В	38	ALA	LEU	engineered mutation	UNP P0ABE7
В	41	TRP	GLN	engineered mutation	UNP P0ABE7
В	42	SER	LYS	engineered mutation	UNP P0ABE7
В	59	HIS	LYS	engineered mutation	UNP P0ABE7
В	66	TRP	ASP	engineered mutation	UNP P0ABE7
В	69	ILE	VAL	engineered mutation	UNP P0ABE7
В	73	HIS	ASP	engineered mutation	UNP P0ABE7
В	77	HIS	LYS	engineered mutation	UNP P0ABE7
В	96	CYS	THR	engineered mutation	UNP P0ABE7
В	98	CYS	ARG	engineered mutation	UNP P0ABE7
В	101	CYS	TYR	engineered mutation	UNP P0ABE7
С	34	ALA	ARG	engineered mutation	UNP P0ABE7
С	38	ALA	LEU	engineered mutation	UNP P0ABE7
С	41	TRP	GLN	engineered mutation	UNP P0ABE7
С	42	SER	LYS	engineered mutation	UNP P0ABE7
С	59	HIS	LYS	engineered mutation	UNP P0ABE7
С	66	TRP	ASP	engineered mutation	UNP P0ABE7
С	69	ILE	VAL	engineered mutation	UNP P0ABE7
С	73	HIS	ASP	engineered mutation	UNP P0ABE7
С	77	HIS	LYS	engineered mutation	UNP P0ABE7
С	96	CYS	THR	engineered mutation	UNP P0ABE7
С	98	CYS	ARG	engineered mutation	UNP P0ABE7
С	101	CYS	TYR	engineered mutation	UNP P0ABE7
D	34	ALA	ARG	engineered mutation	UNP P0ABE7
D	38	ALA	LEU	engineered mutation	UNP P0ABE7
D	41	TRP	GLN	engineered mutation	UNP P0ABE7
D	42	SER	LYS	engineered mutation	UNP P0ABE7
D	59	HIS	LYS	engineered mutation	UNP P0ABE7
D	66	TRP	ASP	engineered mutation	UNP P0ABE7
D	69	ILE	VAL	engineered mutation	UNP P0ABE7
D	73	HIS	ASP	engineered mutation	UNP P0ABE7
D	77	HIS	LYS	engineered mutation	UNP P0ABE7
D	96	CYS	THR	engineered mutation	UNP P0ABE7
D	98	CYS	ARG	engineered mutation	UNP P0ABE7
D	101	CYS	TYR	engineered mutation	UNP P0ABE7
E	34	ALA	ARG	engineered mutation	UNP P0ABE7
Е	38	ALA	LEU	engineered mutation	UNP P0ABE7
E	41	TRP	GLN	engineered mutation	UNP P0ABE7
E	42	SER	LYS	engineered mutation	UNP P0ABE7
E	59	HIS	LYS	engineered mutation	UNP P0ABE7
E	66	TRP	ASP	engineered mutation	UNP P0ABE7
E	69	ILE	VAL	engineered mutation	UNP POABE7




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
Е	73	HIS	ASP	engineered mutation	UNP P0ABE7
Е	77	HIS	LYS	engineered mutation	UNP P0ABE7
Е	96	CYS	THR	engineered mutation	UNP P0ABE7
Е	98	CYS	ARG	engineered mutation	UNP P0ABE7
Е	101	CYS	TYR	engineered mutation	UNP P0ABE7
F	34	ALA	ARG	engineered mutation	UNP P0ABE7
F	38	ALA	LEU	engineered mutation	UNP P0ABE7
F	41	TRP	GLN	engineered mutation	UNP P0ABE7
F	42	SER	LYS	engineered mutation	UNP P0ABE7
F	59	HIS	LYS	engineered mutation	UNP P0ABE7
F	66	TRP	ASP	engineered mutation	UNP P0ABE7
F	69	ILE	VAL	engineered mutation	UNP P0ABE7
F	73	HIS	ASP	engineered mutation	UNP P0ABE7
F	77	HIS	LYS	engineered mutation	UNP P0ABE7
F	96	CYS	THR	engineered mutation	UNP P0ABE7
F	98	CYS	ARG	engineered mutation	UNP P0ABE7
F	101	CYS	TYR	engineered mutation	UNP P0ABE7
G	34	ALA	ARG	engineered mutation	UNP P0ABE7
G	38	ALA	LEU	engineered mutation	UNP P0ABE7
G	41	TRP	GLN	engineered mutation	UNP P0ABE7
G	42	SER	LYS	engineered mutation	UNP P0ABE7
G	59	HIS	LYS	engineered mutation	UNP P0ABE7
G	66	TRP	ASP	engineered mutation	UNP P0ABE7
G	69	ILE	VAL	engineered mutation	UNP P0ABE7
G	73	HIS	ASP	engineered mutation	UNP P0ABE7
G	77	HIS	LYS	engineered mutation	UNP P0ABE7
G	96	CYS	THR	engineered mutation	UNP P0ABE7
G	98	CYS	ARG	engineered mutation	UNP P0ABE7
G	101	CYS	TYR	engineered mutation	UNP P0ABE7
H	34	ALA	ARG	engineered mutation	UNP P0ABE7
H	38	ALA	LEU	engineered mutation	UNP P0ABE7
H	41	TRP	GLN	engineered mutation	UNP P0ABE7
Н	42	SER	LYS	engineered mutation	UNP P0ABE7
H	59	HIS	LYS	engineered mutation	UNP POABE7
H	66	TRP	ASP	engineered mutation	UNP POABE7
H	69	ILE	VAL	engineered mutation	UNP POABE7
Н	73	HIS	ASP	engineered mutation	UNP POABE7
H	77	HIS	LYS	engineered mutation	UNP POABE7
Н	96	CYS	THR	engineered mutation	UNP POABE7
H	98	CYS	ARG	engineered mutation	UNP POABE7
Н	101	CYS	TYR	engineered mutation	UNP P0ABE7

• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (for-



 $mula:\ C_{34}H_{32}FeN_4O_4\big).$ 



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf								
2	A	1	Total	С	Fe	N	О	0	0								
2	Λ	1	43	34	1	4	4		0								
2	В	1	Total	С	Fe	N	Ο	0	0								
	Б	1	43	34	1	4	4	0	U								
2	C	1	Total	С	Fe	N	О	0	0								
2		1	43	34	1	4	4		0								
2	D	1	Total	С	Fe	N	О	0	0								
	D	D					1	43	34	1	4	4	0	U			
2	E	Е	Е	E	E	E	E	E	9 E	1	Total	С	Fe	N	Ο	0	0
				1	43	34	1	4	4	0	0						
2	F	1	Total	С	Fe	N	Ο	0	0								
	I.	1	43	34	1	4	4	0	U								
2	G	1	Total	С	Fe	N	Ο	0	0								
	G	G	G	G	G	G	G	G	1	43	34	1	4	4		U	
2	2 H	1	Total	С	Fe	N	Ο	0	0								
	11	1	43	34	1	4	4		U								

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	В	1	Total Zn 1 1	0	0
3	С	1	Total Zn 1 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Zn 1 1	0	0
3	Е	1	Total Zn 1 1	0	0
3	F	1	Total Zn 1 1	0	0
3	G	1	Total Zn 1 1	0	0
3	Н	1	Total Zn 1 1	0	0

#### • Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	12	Total O 12 12	0	0
4	В	17	Total O 17 17	0	0
4	С	26	Total O 26 26	0	0
4	D	20	Total O 20 20	0	0
4	E	18	Total O 18 18	0	0
4	F	19	Total O 19 19	0	0
4	G	17	Total O 17 17	0	0
4	Н	16	Total O 16 16	0	0



# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Soluble cytochrome b562



• Molecule 1: Soluble cytochrome b562







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	63.27Å 76.77Å 177.57Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 - 2.35	Depositor
rtesolution (A)	88.78 - 2.35	EDS
% Data completeness	98.6 (50.00-2.35)	Depositor
(in resolution range)	99.2 (88.78-2.35)	EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$< I/\sigma(I) > 1$	2.92 (at 2.34Å)	Xtriage
Refinement program	CNS	Depositor
D D.	0.228 , 0.296	Depositor
$R, R_{free}$	0.220 , $0.285$	DCC
$R_{free}$ test set	2562 reflections (7.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.4	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.32, 42.5	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7041	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 16.40% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	$\mathbf{angles}$
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.34	0/833	0.55	0/1125
1	В	0.36	0/833	0.61	0/1125
1	С	0.39	0/833	0.58	0/1125
1	D	0.34	0/833	0.58	0/1125
1	Е	0.39	0/833	0.56	0/1125
1	F	0.36	0/833	0.60	0/1125
1	G	0.34	0/833	0.56	0/1125
1	Н	0.36	0/833	0.56	0/1125
All	All	0.36	0/6664	0.57	0/9000

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	818	0	788	26	0
1	В	818	0	788	17	0
1	С	818	0	788	29	0
1	D	818	0	788	19	0
1	Е	818	0	788	21	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	818	0	788	45	0
1	G	818	0	788	39	0
1	Н	818	0	788	27	0
2	A	43	0	30	3	0
2	В	43	0	30	2	0
2	С	43	0	30	3	0
2	D	43	0	30	1	0
2	Ε	43	0	30	4	0
2	F	43	0	30	2	0
2	G	43	0	30	1	0
2	Н	43	0	30	1	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
3	Ε	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	Н	1	0	0	0	0
4	A	12	0	0	2	0
4	В	17	0	0	0	0
4	С	26	0	0	0	0
4	D	20	0	0	0	0
4	Ε	18	0	0	0	0
4	F	19	0	0	0	0
4	G	17	0	0	1	0
4	Н	16	0	0	0	0
All	All	7041	0	6544	215	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 215 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:13:ASN:HD21	1:A:32:LYS:HE3	1.24	1.00
1:H:21:ASP:H	1:H:25:GLN:HE22	1.05	0.99
1:G:83:LYS:HB3	1:G:86:GLU:HG3	1.40	0.99
1:C:46:PRO:HA	1:C:49:GLU:HG3	1.44	0.97
1:C:21:ASP:H	1:C:25:GLN:HE22	1.07	0.96



There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	104/106 (98%)	98 (94%)	3 (3%)	3 (3%)	4	2
1	В	104/106 (98%)	100 (96%)	4 (4%)	0	100	100
1	С	104/106 (98%)	102 (98%)	2 (2%)	0	100	100
1	D	104/106 (98%)	101 (97%)	2 (2%)	1 (1%)	15	15
1	E	104/106~(98%)	102 (98%)	2 (2%)	0	100	100
1	F	104/106 (98%)	97 (93%)	7 (7%)	0	100	100
1	G	104/106~(98%)	98 (94%)	6 (6%)	0	100	100
1	Н	104/106 (98%)	100 (96%)	4 (4%)	0	100	100
All	All	832/848 (98%)	798 (96%)	30 (4%)	4 (0%)	29	32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	47	LYS
1	D	83	LYS
1	A	103	GLN
1	A	104	LYS

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	84/84 (100%)	80 (95%)	4 (5%)	25	30
1	В	84/84 (100%)	76 (90%)	8 (10%)	8	7
1	С	84/84 (100%)	83 (99%)	1 (1%)	71	82
1	D	84/84 (100%)	81 (96%)	3 (4%)	35	43
1	E	84/84 (100%)	81 (96%)	3 (4%)	35	43
1	F	84/84 (100%)	80 (95%)	4 (5%)	25	30
1	G	84/84 (100%)	82 (98%)	2 (2%)	49	59
1	Н	84/84 (100%)	82 (98%)	2 (2%)	49	59
All	All	672/672 (100%)	645 (96%)	27 (4%)	31	39

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	27	LYS
1	Е	66	TRP
1	G	86	GLU
1	Е	3	LEU
1	Е	92	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	80	ASN
1	F	88	GLN
1	Н	99	ASN
1	Е	88	GLN
1	F	22	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	HEM	Н	150	1	41,50,50	1.49	7 (17%)	45,82,82	1.65	11 (24%)
2	HEM	G	150	1	41,50,50	1.31	5 (12%)	45,82,82	1.41	5 (11%)
2	HEM	D	150	1	41,50,50	1.24	3 (7%)	45,82,82	1.93	8 (17%)
2	HEM	С	150	1	41,50,50	1.28	3 (7%)	45,82,82	1.37	7 (15%)
2	HEM	Е	150	1	41,50,50	1.64	9 (21%)	45,82,82	1.52	8 (17%)
2	HEM	В	150	1	41,50,50	1.25	5 (12%)	45,82,82	1.60	7 (15%)
2	HEM	A	150	1	41,50,50	1.36	4 (9%)	45,82,82	1.55	5 (11%)
2	HEM	F	150	1	41,50,50	1.63	10 (24%)	45,82,82	1.91	10 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	Н	150	1	-	4/12/54/54	-
2	HEM	G	150	1	-	4/12/54/54	-
2	HEM	D	150	1	-	5/12/54/54	-
2	HEM	С	150	1	-	3/12/54/54	-
2	HEM	Е	150	1	-	2/12/54/54	-
2	HEM	В	150	1	-	2/12/54/54	-
2	HEM	A	150	1	-	6/12/54/54	-
2	HEM	F	150	1	-	3/12/54/54	-



The worst 5 of 46 bond length outliers are listed below:
--

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
2	Н	150	HEM	C3C-CAC	-4.10	1.39	1.47
2	A	150	HEM	C3C-CAC	-3.73	1.40	1.47
2	Е	150	HEM	C3C-CAC	-3.57	1.40	1.47
2	Е	150	HEM	C3C-C2C	-3.54	1.35	1.40
2	Е	150	HEM	FE-NB	3.37	2.13	1.96

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	D	150	HEM	CBA-CAA-C2A	7.40	125.26	112.62
2	F	150	HEM	CBA-CAA-C2A	7.17	124.86	112.62
2	В	150	HEM	C4B-CHC-C1C	5.50	129.82	122.56
2	Н	150	HEM	CBA-CAA-C2A	5.29	121.65	112.62
2	G	150	HEM	C4B-CHC-C1C	5.19	129.41	122.56

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	150	HEM	C3A-C2A-CAA-CBA
2	A	150	HEM	C2B-C3B-CAB-CBB
2	D	150	HEM	C1A-C2A-CAA-CBA
2	D	150	HEM	C3A-C2A-CAA-CBA
2	F	150	HEM	C3A-C2A-CAA-CBA

There are no ring outliers.


8 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Н	150	HEM	1	0
2	G	150	HEM	1	0
2	D	150	HEM	1	0
2	С	150	HEM	3	0
2	Е	150	HEM	4	0
2	В	150	HEM	2	0
2	A	150	HEM	3	0
2	F	150	HEM	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will




also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	106/106 (100%)	0.26	3 (2%) 53 64	30, 53, 82, 90	0
1	В	106/106 (100%)	-0.03	0 100 100	28, 44, 60, 65	0
1	С	106/106 (100%)	-0.07	0 100 100	22, 36, 55, 65	0
1	D	106/106 (100%)	0.02	1 (0%) 84 90	29, 44, 65, 76	0
1	E	106/106 (100%)	-0.09	0 100 100	26, 39, 55, 64	0
1	F	106/106 (100%)	0.16	2 (1%) 66 76	27, 51, 82, 94	0
1	G	106/106 (100%)	0.32	7 (6%) 18 26	29, 51, 78, 87	0
1	Н	106/106 (100%)	-0.06	0 100 100	30, 45, 58, 64	0
All	All	848/848 (100%)	0.06	13 (1%) 73 81	22, 45, 70, 94	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	66	TRP	5.3
1	F	66	TRP	2.9
1	G	52	SER	2.8
1	D	54	ASP	2.7
1	A	1	ALA	2.6

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
2	HEM	G	150	43/43	0.94	0.15	42,58,76,82	0
2	HEM	A	150	43/43	0.95	0.18	44,60,79,83	0
2	HEM	F	150	43/43	0.96	0.12	24,35,53,61	0
2	HEM	Е	150	43/43	0.97	0.11	15,23,50,54	0
2	HEM	В	150	43/43	0.97	0.12	23,31,54,63	0
2	HEM	С	150	43/43	0.97	0.12	11,28,49,54	0
2	HEM	Н	150	43/43	0.97	0.13	32,38,59,69	0
2	HEM	D	150	43/43	0.98	0.11	23,41,59,64	0
3	ZN	Е	201	1/1	0.98	0.13	32,32,32,32	0
3	ZN	D	201	1/1	0.99	0.13	34,34,34,34	0
3	ZN	В	201	1/1	0.99	0.12	33,33,33,33	0
3	ZN	F	201	1/1	0.99	0.13	37,37,37,37	0
3	ZN	Н	201	1/1	0.99	0.14	38,38,38,38	0
3	ZN	A	201	1/1	1.00	0.12	30,30,30,30	0
3	ZN	G	201	1/1	1.00	0.12	35,35,35,35	0
3	ZN	С	201	1/1	1.00	0.14	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
































# 





# 6.5 Other polymers (i)

There are no such residues in this entry.

