



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2024 – 02:35 AM EDT

PDB ID : 2IS5
Title : Crystal structure of 3 residues truncated version of protein NMB1012 from *Neisseria meningitides*
Authors : Chang, C.; Li, H.; Abdullah, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2006-10-16
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	
1	B	166	
1	C	166	
1	D	166	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	160	1278	806	237	230	1	4	0	5	0
1	B	150	1177	742	219	211	1	4	0	6	0
1	C	134	1040	659	196	182	1	2	0	7	0
1	D	147	1174	742	220	207	1	4	0	5	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ASN	-	CLONING ARTIFACT	UNP Q7DDI9
A	1	ALA	-	CLONING ARTIFACT	UNP Q7DDI9
A	48	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
A	51	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
A	73	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
A	96	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
B	0	ASN	-	CLONING ARTIFACT	UNP Q7DDI9
B	1	ALA	-	CLONING ARTIFACT	UNP Q7DDI9
B	48	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
B	51	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
B	73	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
B	96	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
C	0	ASN	-	CLONING ARTIFACT	UNP Q7DDI9
C	1	ALA	-	CLONING ARTIFACT	UNP Q7DDI9
C	48	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
C	51	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
C	73	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
C	96	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
D	0	ASN	-	CLONING ARTIFACT	UNP Q7DDI9
D	1	ALA	-	CLONING ARTIFACT	UNP Q7DDI9
D	48	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	51	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
D	73	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9
D	96	MSE	MET	MODIFIED RESIDUE	UNP Q7DDI9

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 13 6 7	0	0
2	D	1	Total C O 13 6 7	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	132	Total O 132 132	0	0
3	B	97	Total O 97 97	0	0
3	C	73	Total O 73 73	0	0
3	D	111	Total O 111 111	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hypothetical protein





- Molecule 1: Hypothetical protein



- Molecule 1: Hypothetical protein



- Molecule 1: Hypothetical protein



PHE	
THR	
SER	
PHE	
GLY	
LYS	
G149	●
W150	
V151	
R152	
G156	●
A163	●
D164	●
ASN	

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.74Å 79.79Å 65.61Å 90.00° 105.40° 90.00°	Depositor
Resolution (Å)	39.80 – 1.85 39.80 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (39.80-1.85) 99.4 (39.80-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.56 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.197 , 0.229 0.195 , 0.229	Depositor DCC
R_{free} test set	2811 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtrriage
Anisotropy	0.489	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5108	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/1307	0.74	1/1755 (0.1%)
1	B	0.74	0/1202	0.72	0/1613
1	C	0.84	2/1056 (0.2%)	0.78	2/1419 (0.1%)
1	D	0.75	0/1199	0.76	2/1607 (0.1%)
All	All	0.80	2/4764 (0.0%)	0.75	5/6394 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	2
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	52	THR	N-CA	10.35	1.67	1.46
1	C	52	THR	CA-CB	6.44	1.70	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	17	GLY	N-CA-C	-8.43	92.03	113.10
1	A	23	PRO	N-CA-CB	6.38	110.95	103.30
1	D	18	GLY	N-CA-C	-6.28	97.40	113.10
1	C	52	THR	N-CA-CB	5.92	121.55	110.30
1	C	133	ARG	NE-CZ-NH2	-5.78	117.41	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	ALA	Peptide
1	D	16	GLU	Peptide
1	D	17	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1278	0	1235	19	0
1	B	1177	0	1121	30	0
1	C	1040	0	1010	19	0
1	D	1174	0	1134	17	0
2	C	13	0	5	1	0
2	D	13	0	5	0	0
3	A	132	0	0	3	0
3	B	97	0	0	4	0
3	C	73	0	0	0	0
3	D	111	0	0	1	0
All	All	5108	0	4510	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87[A]:ASN:HD21	1:A:150:TRP:HE1	1.04	1.02
1:A:54[B]:GLU:OE2	3:A:297:HOH:O	1.92	0.87
1:A:134[A]:LEU:HG	1:A:154:VAL:HG11	1.60	0.84
1:C:52:THR:HG23	1:C:55:GLN:H	1.44	0.83
1:A:87[A]:ASN:ND2	1:A:150:TRP:HE1	1.76	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/166 (96%)	158 (99%)	2 (1%)	0	100	100
1	B	145/166 (87%)	143 (99%)	2 (1%)	0	100	100
1	C	126/166 (76%)	125 (99%)	1 (1%)	0	100	100
1	D	143/166 (86%)	142 (99%)	1 (1%)	0	100	100
All	All	574/664 (86%)	568 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/126 (99%)	124 (99%)	1 (1%)	81	76
1	B	113/126 (90%)	109 (96%)	4 (4%)	36	18
1	C	99/126 (79%)	98 (99%)	1 (1%)	76	69
1	D	114/126 (90%)	112 (98%)	2 (2%)	59	45
All	All	451/504 (90%)	443 (98%)	8 (2%)	59	45

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	152	ARG
1	D	16	GLU
1	B	120	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	54	GLU
1	C	120	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	10	ASN
1	D	88	HIS
1	D	118	ASN
1	D	92	ASN
1	B	88	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CIT	D	501	-	12,12,12	1.17	1 (8%)	17,17,17	1.62	2 (11%)
2	CIT	C	502	-	12,12,12	1.27	1 (8%)	17,17,17	1.74	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	D	501	-	-	1/16/16/16	-
2	CIT	C	502	-	-	6/16/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	502	CIT	O2-C1	-2.08	1.23	1.30
2	D	501	CIT	O1-C1	2.02	1.28	1.22

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	501	CIT	O6-C6-C3	4.34	121.47	113.14
2	C	502	CIT	O6-C6-C3	3.55	119.95	113.14
2	D	501	CIT	C2-C3-C6	-2.92	103.58	110.03
2	C	502	CIT	O6-C6-O5	-2.55	115.70	123.86
2	C	502	CIT	O4-C5-C4	2.30	121.65	114.35

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	502	CIT	C2-C3-C6-O6
2	C	502	CIT	O7-C3-C6-O6
2	C	502	CIT	C2-C3-C6-O5
2	C	502	CIT	C4-C3-C6-O6
2	C	502	CIT	C4-C3-C6-O5

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	502	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	156/166 (93%)	0.69	14 (8%) 9 9	19, 29, 38, 48	1 (0%)
1	B	146/166 (87%)	0.75	15 (10%) 6 6	20, 32, 44, 58	4 (2%)
1	C	132/166 (79%)	0.77	17 (12%) 3 3	22, 32, 49, 52	7 (5%)
1	D	143/166 (86%)	0.72	15 (10%) 6 6	21, 32, 41, 81	2 (1%)
All	All	577/664 (86%)	0.73	61 (10%) 6 6	19, 31, 44, 81	14 (2%)

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	52	THR	5.7
1	A	27	GLY	5.7
1	D	19	TYR	5.7
1	B	2	SER	5.1
1	C	134[A]	LEU	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CIT	D	501	13/13	0.86	0.16	23,32,34,34	0
2	CIT	C	502	13/13	0.89	0.15	34,41,45,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.