



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2024 – 03:39 AM EST

PDB ID : 4ISR
Title : Binding domain of Botulinum neurotoxin DC in complex with rat synaptotagmin II
Authors : Berntsson, R.P.-A.; Peng, L.; Svensson, L.M.; Dong, M.; Stenmark, P.
Deposited on : 2013-01-17
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	431	 3% 89% 7%
1	B	431	 3% 91% 5%
1	C	431	 9% 90% 6%
2	D	21	 5% 67% 10% 24%
2	E	21	 52% 24% 24%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	21	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into five segments: a red segment (14%), a green segment (52%), a yellow segment (19%), an orange segment (5%), and a grey segment (24%).</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11056 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neurotoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	414	3398	2181	561	642	14	0	1	0
1	B	413	3378	2168	558	638	14	0	0	0
1	C	413	3376	2167	558	637	14	0	0	0

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	854	MET	-	expression tag	UNP Q9LBR1
A	855	HIS	-	expression tag	UNP Q9LBR1
A	856	HIS	-	expression tag	UNP Q9LBR1
A	857	HIS	-	expression tag	UNP Q9LBR1
A	858	HIS	-	expression tag	UNP Q9LBR1
A	859	HIS	-	expression tag	UNP Q9LBR1
A	860	HIS	-	expression tag	UNP Q9LBR1
A	861	TYR	-	expression tag	UNP Q9LBR1
A	862	PHE	-	expression tag	UNP Q9LBR1
B	854	MET	-	expression tag	UNP Q9LBR1
B	855	HIS	-	expression tag	UNP Q9LBR1
B	856	HIS	-	expression tag	UNP Q9LBR1
B	857	HIS	-	expression tag	UNP Q9LBR1
B	858	HIS	-	expression tag	UNP Q9LBR1
B	859	HIS	-	expression tag	UNP Q9LBR1
B	860	HIS	-	expression tag	UNP Q9LBR1
B	861	TYR	-	expression tag	UNP Q9LBR1
B	862	PHE	-	expression tag	UNP Q9LBR1
C	854	MET	-	expression tag	UNP Q9LBR1
C	855	HIS	-	expression tag	UNP Q9LBR1
C	856	HIS	-	expression tag	UNP Q9LBR1
C	857	HIS	-	expression tag	UNP Q9LBR1
C	858	HIS	-	expression tag	UNP Q9LBR1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	859	HIS	-	expression tag	UNP Q9LBR1
C	860	HIS	-	expression tag	UNP Q9LBR1
C	861	TYR	-	expression tag	UNP Q9LBR1
C	862	PHE	-	expression tag	UNP Q9LBR1

- Molecule 2 is a protein called Synaptotagmin-2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	D	16	Total	C	N	O	S	0	0	0
			138	89	21	27	1			
2	E	16	Total	C	N	O	S	0	0	0
			138	89	21	27	1			
2	F	16	Total	C	N	O	S	0	0	0
			138	89	21	27	1			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	176	Total	O	0	0
			176	176		
4	B	156	Total	O	0	0
			156	156		
4	C	90	Total	O	0	0
			90	90		
4	D	3	Total	O	0	0
			3	3		
4	E	2	Total	O	0	0
			2	2		
4	F	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Neurotoxin



- Molecule 1: Neurotoxin




- Molecule 1: Neurotoxin



- Molecule 2: Synaptotagmin-2






- Molecule 2: Synaptotagmin-2



- Molecule 2: Synaptotagmin-2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	165.24Å 57.78Å 169.56Å 90.00° 118.49° 90.00°	Depositor
Resolution (Å)	47.73 – 2.59 47.73 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.73-2.59) 99.6 (47.73-2.59)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.58Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.220 , 0.235 0.202 , 0.218	Depositor DCC
R_{free} test set	4426 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for -h-l,k,h 0.017 for l,k,-h-l 0.017 for h,-k,-h-l 0.019 for -h-l,-k,l 0.018 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11056	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3476	0.48	0/4704
1	B	0.30	0/3455	0.47	0/4675
1	C	0.30	0/3453	0.47	0/4673
2	D	0.35	0/140	0.50	0/183
2	E	0.35	0/140	0.55	0/183
2	F	0.35	0/140	0.49	0/183
All	All	0.30	0/10804	0.47	0/14601

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3398	0	3318	19	0
1	B	3378	0	3304	16	0
1	C	3376	0	3299	20	0
2	D	138	0	129	3	0
2	E	138	0	129	9	0
2	F	138	0	129	3	0
3	A	20	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	20	0	0	0	0
3	C	15	0	0	0	0
3	E	5	0	0	0	0
4	A	176	0	0	4	0
4	B	156	0	0	2	0
4	C	90	0	0	0	0
4	D	3	0	0	0	0
4	E	2	0	0	0	0
4	F	3	0	0	0	0
All	All	11056	0	10308	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 68 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:885:TYR:HB3	1:B:925:ILE:CD1	2.08	0.83
2:E:56:ASN:HA	2:E:57:GLU:HG3	1.61	0.83
2:D:56:ASN:HA	2:D:57:GLU:HB2	1.62	0.81
1:A:1125:ASN:HB2	4:A:1566:HOH:O	1.81	0.79
1:C:923:GLU:HB3	1:C:924:ASN:HB2	1.72	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	411/431 (95%)	392 (95%)	19 (5%)	0	100 100
1	B	409/431 (95%)	389 (95%)	18 (4%)	2 (0%)	29 52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	409/431 (95%)	386 (94%)	22 (5%)	1 (0%)	47	71
2	D	14/21 (67%)	14 (100%)	0	0	100	100
2	E	14/21 (67%)	14 (100%)	0	0	100	100
2	F	14/21 (67%)	13 (93%)	0	1 (7%)	1	1
All	All	1271/1356 (94%)	1208 (95%)	59 (5%)	4 (0%)	41	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1033	LEU
2	F	43	GLN
1	B	1032	GLU
1	C	925	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/395 (96%)	379 (100%)	1 (0%)	92	98
1	B	378/395 (96%)	378 (100%)	0	100	100
1	C	377/395 (95%)	375 (100%)	2 (0%)	88	96
2	D	15/19 (79%)	15 (100%)	0	100	100
2	E	15/19 (79%)	15 (100%)	0	100	100
2	F	15/19 (79%)	14 (93%)	1 (7%)	16	33
All	All	1180/1242 (95%)	1176 (100%)	4 (0%)	92	98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1137	ASP
1	C	1034	THR
1	C	1152	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	56	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1063	ASN
1	C	1152	ASN
2	E	43	GLN
1	C	1154	ASN
1	B	1223	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1304	-	4,4,4	0.33	0	6,6,6	0.13	0
3	SO4	B	1304	-	4,4,4	0.36	0	6,6,6	0.10	0
3	SO4	B	1301	-	4,4,4	0.30	0	6,6,6	0.12	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	1303	-	4,4,4	0.34	0	6,6,6	0.07	0
3	SO4	C	1302	-	4,4,4	0.32	0	6,6,6	0.16	0
3	SO4	A	1301	-	4,4,4	0.31	0	6,6,6	0.11	0
3	SO4	B	1303	-	4,4,4	0.34	0	6,6,6	0.11	0
3	SO4	C	1301	-	4,4,4	0.33	0	6,6,6	0.15	0
3	SO4	A	1302	-	4,4,4	0.34	0	6,6,6	0.12	0
3	SO4	C	1303	-	4,4,4	0.35	0	6,6,6	0.10	0
3	SO4	B	1302	-	4,4,4	0.36	0	6,6,6	0.09	0
3	SO4	E	101	-	4,4,4	0.34	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	414/431 (96%)	-0.01	11 (2%) 54 48	20, 31, 60, 89	0
1	B	413/431 (95%)	0.04	15 (3%) 42 35	17, 31, 74, 99	0
1	C	413/431 (95%)	0.36	38 (9%) 9 6	24, 45, 96, 135	0
2	D	16/21 (76%)	0.59	1 (6%) 20 15	29, 47, 76, 78	0
2	E	16/21 (76%)	0.75	0 100 100	27, 39, 64, 64	0
2	F	16/21 (76%)	1.42	3 (18%) 1 0	38, 53, 76, 77	0
All	All	1288/1356 (94%)	0.16	68 (5%) 26 20	17, 35, 82, 135	0

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	927	TYR	6.7
1	C	1050	PRO	5.9
1	B	912	ASP	4.9
1	B	925	ILE	4.9
1	C	922	ASN	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	1304	5/5	0.81	0.32	77,78,80,81	0
3	SO4	B	1302	5/5	0.82	0.20	61,62,62,64	0
3	SO4	A	1302	5/5	0.83	0.21	63,65,67,67	0
3	SO4	B	1303	5/5	0.87	0.30	73,73,74,75	0
3	SO4	A	1303	5/5	0.91	0.25	86,87,89,89	0
3	SO4	C	1302	5/5	0.92	0.18	62,63,63,64	0
3	SO4	A	1304	5/5	0.95	0.19	68,68,69,69	0
3	SO4	C	1303	5/5	0.95	0.24	70,70,71,71	0
3	SO4	E	101	5/5	0.97	0.15	62,62,64,64	0
3	SO4	C	1301	5/5	0.98	0.11	44,44,45,45	0
3	SO4	A	1301	5/5	0.98	0.16	33,33,34,34	0
3	SO4	B	1301	5/5	0.99	0.13	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.