



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 01:06 AM EDT

PDB ID : 5ISG  
Title : Crystal structure of mouse CARM1 in complex with inhibitor SA0707  
Authors : Cura, V.; Marechal, N.; Mailliot, J.; Troffer-Charlier, N.; Hassenboehler, P.;  
Wurtz, J.M.; Bonnefond, L.; Cavarelli, J.  
Deposited on : 2016-03-15  
Resolution : 2.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	<p>93% • 5%</p>
1	B	361	<p>93% • 5%</p>
1	C	361	<p>93% • 5%</p>
1	D	361	<p>94% • 5%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	D	504	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 22708 atoms, of which 11063 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	343	5466	1781	2707	455	508	15	0	1	0
1	B	343	5477	1784	2712	460	507	14	0	1	0
1	C	343	5467	1781	2708	455	508	15	0	1	0
1	D	343	5481	1787	2714	458	508	14	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	GLY	-	expression tag	UNP Q9WVG6
A	128	HIS	-	expression tag	UNP Q9WVG6
A	129	MET	-	expression tag	UNP Q9WVG6
B	127	GLY	-	expression tag	UNP Q9WVG6
B	128	HIS	-	expression tag	UNP Q9WVG6
B	129	MET	-	expression tag	UNP Q9WVG6
C	127	GLY	-	expression tag	UNP Q9WVG6
C	128	HIS	-	expression tag	UNP Q9WVG6
C	129	MET	-	expression tag	UNP Q9WVG6
D	127	GLY	-	expression tag	UNP Q9WVG6
D	128	HIS	-	expression tag	UNP Q9WVG6
D	129	MET	-	expression tag	UNP Q9WVG6

- Molecule 2 is 1,2-DIMETHOXYETHANE (three-letter code: DXE) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	16	4	10	2	0	0
2	A	1	16	4	10	2	0	0
2	B	1	16	4	10	2	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	10	2	6	2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C H O 10 2 6 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C H O 10 2 6 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C H O 10 2 6 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is 1-(2-METHOXY-ETHOXY)-2-{2-[2-(2-METHOXY-ETHOXY)-ETHOXY]}-ETHANE (three-letter code: PG6) (formula: C<sub>12</sub>H<sub>26</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	44	12	26	6	0	0


- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	A	1	17	4	10	3	0	0
5	D	1	17	4	10	3	0	0

- Molecule 6 is 5'-{[(3S)-3-amino-3-carboxypropyl](3-aminopropyl)amino}-5'-deoxyadenosine


(three-letter code: 6D6) (formula:  $C_{17}H_{28}N_8O_5$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			
6	A	1	Total	57	17	27	8	5	0	0
6	B	1	Total	57	17	27	8	5	0	0
6	C	1	Total	57	17	27	8	5	0	0
6	D	1	Total	57	17	27	8	5	0	0

- Molecule 7 is 1-METHOXY-2-(2-METHOXYETHOXY)ETHANE (three-letter code: M2M) (formula:  $C_6H_{14}O_3$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
7	A	1	23	6	14	3	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	105	Total	O	0	0
			105	105		
8	B	84	Total	O	0	0
			84	84		
8	C	81	Total	O	0	0
			81	81		
8	D	74	Total	O	0	0
			74	74		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone-arginine methyltransferase CARM1

Chain A: 




- Molecule 1: Histone-arginine methyltransferase CARM1

Chain B: 



- Molecule 1: Histone-arginine methyltransferase CARM1

Chain C: 



- Molecule 1: Histone-arginine methyltransferase CARM1

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.69Å 98.16Å 205.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.12 – 2.42 20.12 – 2.42	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.12-2.42) 98.4 (20.12-2.42)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.36 (at 2.41Å)	Xtrriage
Refinement program	PHENIX (dev_2313: ???)	Depositor
R, $R_{free}$	0.161 , 0.207 0.163 , 0.207	Depositor DCC
$R_{free}$ test set	2914 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.3	Xtrriage
Anisotropy	0.227	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 49.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	22708	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.00 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0099e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG6, 6D6, M2M, DXE, EDO, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/2829	0.45	0/3833
1	B	0.25	0/2836	0.44	0/3842
1	C	0.25	0/2829	0.45	0/3833
1	D	0.25	0/2837	0.44	0/3842
All	All	0.25	0/11331	0.44	0/15350

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2759	2707	2707	6	0
1	B	2765	2712	2711	4	0
1	C	2759	2708	2707	5	0
1	D	2767	2714	2714	2	0
2	A	12	20	20	0	0
2	B	6	10	10	0	0
3	A	16	6	24	1	0
3	B	16	6	24	0	0
3	C	24	6	36	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	16	6	24	0	0
4	A	18	26	26	0	0
5	A	7	10	10	0	0
5	D	7	10	10	0	0
6	A	30	27	0	1	0
6	B	30	27	0	0	0
6	C	30	27	0	0	0
6	D	30	27	0	0	0
7	A	9	14	14	0	0
8	A	105	0	0	2	0
8	B	84	0	0	1	0
8	C	81	0	0	1	0
8	D	74	0	0	1	0
All	All	11645	11063	11037	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 17 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:MET:O	6:A:509:6D6:N1	2.18	0.77
1:A:469:ASP:OD1	8:A:601:HOH:O	2.07	0.72
1:C:469:ASP:OD1	8:C:601:HOH:O	2.13	0.66
1:B:362:GLU:OE2	8:B:601:HOH:O	2.14	0.65
1:A:159:GLN:HE21	3:A:505:EDO:H12	1.69	0.57

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/361 (95%)	332 (97%)	10 (3%)	0	100	100
1	B	342/361 (95%)	330 (96%)	12 (4%)	0	100	100
1	C	342/361 (95%)	332 (97%)	10 (3%)	0	100	100
1	D	341/361 (94%)	331 (97%)	10 (3%)	0	100	100
All	All	1367/1444 (95%)	1325 (97%)	42 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/313 (96%)	300 (100%)	0	100	100
1	B	300/313 (96%)	298 (99%)	2 (1%)	84	92
1	C	300/313 (96%)	299 (100%)	1 (0%)	92	97
1	D	300/313 (96%)	298 (99%)	2 (1%)	84	92
All	All	1200/1252 (96%)	1195 (100%)	5 (0%)	91	96

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	258	GLU
1	B	308	PHE
1	C	258	GLU
1	D	162	ASN
1	D	258	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	159	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	A	504	-	3,3,3	0.48	0	2,2,2	0.49	0
3	EDO	C	505	-	3,3,3	0.48	0	2,2,2	0.51	0
6	6D6	D	506	-	28,32,32	1.86	6 (21%)	29,45,45	1.50	4 (13%)
3	EDO	D	504	-	3,3,3	0.48	0	2,2,2	0.46	0
6	6D6	A	509	-	28,32,32	1.89	6 (21%)	29,45,45	1.40	3 (10%)
2	DXE	A	502	-	5,5,5	0.42	0	4,4,4	0.20	0
3	EDO	C	506	-	3,3,3	0.47	0	2,2,2	0.51	0
3	EDO	A	503	-	3,3,3	0.47	0	2,2,2	0.50	0
3	EDO	A	505	-	3,3,3	0.48	0	2,2,2	0.49	0
3	EDO	B	503	-	3,3,3	0.47	0	2,2,2	0.51	0
3	EDO	D	501	-	3,3,3	0.47	0	2,2,2	0.49	0
3	EDO	B	502	-	3,3,3	0.46	0	2,2,2	0.48	0
7	M2M	A	510	-	8,8,8	0.60	0	7,7,7	0.30	0
5	PEG	D	505	-	6,6,6	0.48	0	5,5,5	0.48	0
5	PEG	A	508	-	6,6,6	0.49	0	5,5,5	0.49	0
3	EDO	C	503	-	3,3,3	0.47	0	2,2,2	0.47	0
3	EDO	B	505	-	3,3,3	0.47	0	2,2,2	0.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	D	502	-	3,3,3	0.48	0	2,2,2	0.47	0
3	EDO	C	501	-	3,3,3	0.48	0	2,2,2	0.48	0
3	EDO	D	503	-	3,3,3	0.48	0	2,2,2	0.47	0
4	PG6	A	507	-	17,17,17	0.52	0	16,16,16	0.38	0
6	6D6	B	506	-	28,32,32	1.90	6 (21%)	29,45,45	1.53	5 (17%)
2	DXE	B	501	-	5,5,5	0.42	0	4,4,4	0.22	0
3	EDO	B	504	-	3,3,3	0.48	0	2,2,2	0.52	0
6	6D6	C	507	-	28,32,32	1.86	6 (21%)	29,45,45	1.52	5 (17%)
2	DXE	A	501	-	5,5,5	0.42	0	4,4,4	0.21	0
3	EDO	A	506	-	3,3,3	0.47	0	2,2,2	0.50	0
3	EDO	C	502	-	3,3,3	0.48	0	2,2,2	0.45	0
3	EDO	C	504	-	3,3,3	0.48	0	2,2,2	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	504	-	-	0/1/1/1	-
3	EDO	C	505	-	-	1/1/1/1	-
6	6D6	D	506	-	-	7/17/37/37	0/3/3/3
3	EDO	D	504	-	-	1/1/1/1	-
6	6D6	A	509	-	-	5/17/37/37	0/3/3/3
2	DXE	A	502	-	-	1/3/3/3	-
3	EDO	C	506	-	-	0/1/1/1	-
3	EDO	A	503	-	-	0/1/1/1	-
3	EDO	A	505	-	-	0/1/1/1	-
3	EDO	B	503	-	-	0/1/1/1	-
3	EDO	D	501	-	-	1/1/1/1	-
3	EDO	B	502	-	-	1/1/1/1	-
7	M2M	A	510	-	-	4/6/6/6	-
5	PEG	D	505	-	-	2/4/4/4	-
5	PEG	A	508	-	-	1/4/4/4	-
3	EDO	C	503	-	-	1/1/1/1	-
3	EDO	B	505	-	-	1/1/1/1	-
3	EDO	D	502	-	-	1/1/1/1	-
3	EDO	C	501	-	-	1/1/1/1	-
3	EDO	D	503	-	-	1/1/1/1	-
4	PG6	A	507	-	-	9/15/15/15	-
6	6D6	B	506	-	-	9/17/37/37	0/3/3/3

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DXE	B	501	-	-	2/3/3/3	-
3	EDO	B	504	-	-	0/1/1/1	-
6	6D6	C	507	-	-	4/17/37/37	0/3/3/3
2	DXE	A	501	-	-	2/3/3/3	-
3	EDO	A	506	-	-	1/1/1/1	-
3	EDO	C	502	-	-	0/1/1/1	-
3	EDO	C	504	-	-	0/1/1/1	-

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	506	6D6	O3-C13	6.94	1.50	1.41
6	A	509	6D6	O3-C13	6.85	1.50	1.41
6	D	506	6D6	O3-C13	6.70	1.50	1.41
6	C	507	6D6	O3-C13	6.67	1.50	1.41
6	C	507	6D6	C10-C4	2.88	1.58	1.52

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	507	6D6	N8-C16-N7	-4.65	121.40	128.68
6	B	506	6D6	N8-C16-N7	-4.61	121.47	128.68
6	D	506	6D6	N8-C16-N7	-4.61	121.48	128.68
6	A	509	6D6	N8-C16-N7	-4.58	121.53	128.68
6	B	506	6D6	C3-N2-C6	-3.52	103.56	111.96

There are no chirality outliers.

5 of 56 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	509	6D6	C7-C12-C6-N2
6	A	509	6D6	O3-C12-C6-N2
6	B	506	6D6	C4-C10-C5-N3
6	B	506	6D6	C7-C12-C6-N2
6	B	506	6D6	O3-C12-C6-N2


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	509	6D6	1	0
3	A	505	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	343/361 (95%)	-0.41	4 (1%) 79 76	20, 33, 57, 78	0
1	B	343/361 (95%)	-0.38	5 (1%) 73 71	20, 37, 59, 85	0
1	C	343/361 (95%)	-0.25	8 (2%) 60 57	28, 41, 65, 81	0
1	D	343/361 (95%)	-0.26	7 (2%) 65 62	30, 42, 65, 97	0
All	All	1372/1444 (95%)	-0.32	24 (1%) 70 67	20, 39, 63, 97	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	135	ARG	6.0
1	C	478	THR	5.4
1	D	477[A]	TYR	5.4
1	B	135	ARG	5.4
1	A	478	THR	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	D	504	4/4	0.52	0.59	65,69,73,75	0
3	EDO	B	505	4/4	0.60	0.35	56,57,58,59	0
4	PG6	A	507	18/18	0.72	0.27	51,63,74,74	0
5	PEG	A	508	7/7	0.76	0.26	54,65,68,68	0
3	EDO	D	503	4/4	0.81	0.18	55,57,57,57	0
3	EDO	C	502	4/4	0.81	0.19	48,50,50,51	0
3	EDO	A	506	4/4	0.83	0.22	45,46,49,50	0
2	DXE	B	501	6/6	0.84	0.22	45,55,66,66	0
7	M2M	A	510	9/9	0.84	0.38	39,62,78,79	0
5	PEG	D	505	7/7	0.85	0.21	46,72,91,92	0
3	EDO	C	506	4/4	0.86	0.19	47,50,56,62	0
3	EDO	C	504	4/4	0.88	0.43	62,69,78,84	0
2	DXE	A	501	6/6	0.88	0.14	46,56,67,67	0
2	DXE	A	502	6/6	0.89	0.25	51,61,63,63	0
3	EDO	B	502	4/4	0.89	0.17	42,51,52,53	0
3	EDO	A	503	4/4	0.89	0.27	53,64,65,66	0
3	EDO	D	501	4/4	0.90	0.11	48,58,59,60	0
3	EDO	C	505	4/4	0.91	0.22	43,49,61,69	0
3	EDO	A	504	4/4	0.92	0.19	49,51,55,59	0
3	EDO	A	505	4/4	0.92	0.16	50,55,63,69	0
3	EDO	C	501	4/4	0.93	0.15	48,58,60,62	0
3	EDO	D	502	4/4	0.93	0.37	56,57,58,60	0
3	EDO	B	504	4/4	0.93	0.38	47,48,54,58	0
3	EDO	C	503	4/4	0.93	0.17	48,49,49,51	0
6	6D6	D	506	30/30	0.94	0.15	30,45,73,77	0
6	6D6	C	507	30/30	0.94	0.16	27,41,71,74	0
3	EDO	B	503	4/4	0.95	0.11	38,40,41,47	0
6	6D6	B	506	30/30	0.96	0.14	20,44,69,70	0
6	6D6	A	509	30/30	0.96	0.17	21,33,66,66	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around PG6 A 507:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 6D6 D 506:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around 6D6 C 507:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 6D6 B 506:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.