



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 07:46 pm BST

PDB ID : 2IT1  
Title : Structure of PH0203 protein from *Pyrococcus horikoshii*  
Authors : Lokanath, N.K.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2006-10-18  
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	
1	B	362	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6134 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 362aa long hypothetical maltose/maltodextrin transport ATP-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	362	2881	1867	476	532	6	0	0	0
1	B	361	2873	1862	475	531	5	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	B	1	5	4	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	172	172	172	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	203	Total 203	O 203	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 362aa long hypothetical maltose/maltodextrin transport ATP-binding protein



- Molecule 1: 362aa long hypothetical maltose/maltodextrin transport ATP-binding protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.23Å 85.32Å 82.60Å 90.00° 97.16° 90.00°	Depositor
Resolution (Å)	20.00 – 1.94 44.47 – 1.94	Depositor EDS
% Data completeness (in resolution range)	98.1 (20.00-1.94) 98.1 (44.47-1.94)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.75 (at 1.94Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.210 , 0.250 0.215 , 0.223	Depositor DCC
$R_{free}$ test set	2471 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.4	Xtrriage
Anisotropy	0.422	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6134	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/2936	0.66	0/3963
1	B	0.63	0/2928	0.70	0/3953
All	All	0.61	0/5864	0.68	0/7916

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2881	0	2990	29	0
1	B	2873	0	2978	44	0
2	B	5	0	0	0	0
3	A	172	0	0	2	0
3	B	203	0	0	3	0
All	All	6134	0	5968	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:341:VAL:HG21	1:A:346:LEU:HD11	1.60	0.83
1:A:254:THR:HG22	1:A:256:LYS:H	1.54	0.72
1:A:254:THR:HG22	1:A:255:GLU:N	2.04	0.72
1:B:341:VAL:HG23	1:B:341:VAL:O	1.94	0.67
1:A:4:ILE:HD11	1:A:6:LEU:HD11	1.77	0.66
1:B:272:THR:OG1	1:B:274:ILE:HG13	1.96	0.66
1:A:341:VAL:CG2	1:A:346:LEU:HD11	2.28	0.64
1:B:296:ILE:HB	1:B:339:ILE:HG12	1.78	0.64
1:B:237:PRO:HG3	1:B:311:ILE:CD1	2.28	0.63
1:A:254:THR:CG2	1:A:255:GLU:N	2.64	0.60
1:B:292:GLU:HG3	1:B:336:LYS:HG2	1.85	0.59
1:A:286:ALA:HB2	1:A:341:VAL:HG12	1.84	0.58
1:B:120:MET:SD	1:B:183:LEU:HD21	2.44	0.57
1:A:198:LEU:HD23	1:A:204:ILE:CD1	2.36	0.55
1:A:175:GLU:OE2	1:A:179:LEU:HD11	2.07	0.54
1:B:103:ARG:O	1:B:103:ARG:HG2	2.08	0.54
1:B:237:PRO:CG	1:B:311:ILE:HD11	2.37	0.53
1:B:77:VAL:HG12	1:B:154:VAL:CG1	2.39	0.53
1:A:313:THR:CG2	1:A:320:ILE:CG2	2.87	0.53
1:A:267:GLU:O	1:A:271:GLU:HG2	2.09	0.53
1:A:322:LYS:NZ	3:A:524:HOH:O	2.42	0.52
1:B:308:ARG:HG3	1:B:308:ARG:O	2.09	0.52
1:B:313:THR:HG23	1:B:320:ILE:HG13	1.92	0.52
1:B:237:PRO:HG3	1:B:311:ILE:HD11	1.91	0.52
1:B:254:THR:HG22	1:B:255:GLU:H	1.74	0.52
1:B:194:GLN:CG	1:B:234:LEU:HA	2.40	0.51
1:B:113:LYS:NZ	1:B:149:VAL:O	2.39	0.51
1:B:254:THR:HG22	1:B:255:GLU:N	2.26	0.50
1:B:173:ARG:HB2	1:B:200:MET:SD	2.50	0.50
1:B:160:PRO:HG2	1:B:161:LEU:HD13	1.93	0.50
1:B:237:PRO:CG	1:B:311:ILE:CD1	2.90	0.49
1:B:284:HIS:CE1	1:B:324:PHE:CE2	3.00	0.49
1:B:9:ILE:HB	1:B:22:ILE:HB	1.94	0.49
1:A:247:GLU:HB2	1:A:252:VAL:HG21	1.95	0.48
1:B:140:GLN:CG	3:B:473:HOH:O	2.61	0.47
1:B:194:GLN:HG2	1:B:234:LEU:HA	1.96	0.47
1:B:240:ASN:O	1:B:280:GLY:HA2	2.13	0.47
1:B:140:GLN:HG3	3:B:473:HOH:O	2.14	0.47
1:B:228:LYS:HD2	1:B:278:ILE:HD13	1.97	0.47
1:B:61:TYR:HA	1:B:65:LYS:O	2.15	0.47
1:A:228:LYS:HD2	1:A:278:ILE:HD13	1.97	0.47
1:B:2:VAL:HG21	1:B:62:PHE:HB3	1.97	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:281:PHE:HB3	1:B:346:LEU:HD23	1.98	0.46
1:A:77:VAL:HG12	1:A:154:VAL:CG1	2.46	0.46
1:B:289:VAL:HG21	1:B:340:LYS:HD3	1.97	0.46
1:A:343:GLU:HA	1:A:346:LEU:HD22	1.99	0.45
1:B:249:GLY:HA2	1:B:261:ILE:HD12	1.99	0.45
1:B:131:PRO:O	1:B:139:GLN:NE2	2.50	0.45
1:A:286:ALA:CB	1:A:341:VAL:HG12	2.48	0.44
1:A:116:GLU:HG3	3:B:523:HOH:O	2.17	0.44
1:A:100:LEU:HD11	1:A:113:LYS:HD3	1.98	0.44
1:A:297:VAL:HG12	1:A:336:LYS:HE2	1.99	0.44
1:A:158:ASP:HA	1:A:190:VAL:HB	2.00	0.44
1:A:116:GLU:HG2	1:A:120:MET:CE	2.48	0.44
1:A:176:LEU:HA	1:A:179:LEU:HD12	2.00	0.43
1:B:237:PRO:HG3	1:B:311:ILE:HD12	1.97	0.43
1:B:257:SER:OG	1:B:297:VAL:HG22	2.18	0.43
1:B:258:LYS:HD3	1:B:258:LYS:N	2.34	0.43
1:A:214:GLN:HG3	1:A:226:LYS:HE2	2.01	0.43
1:A:2:VAL:HG11	1:A:62:PHE:HB3	2.00	0.43
1:B:308:ARG:CG	1:B:308:ARG:O	2.66	0.42
1:A:358:GLU:HG3	3:A:413:HOH:O	2.19	0.42
1:B:292:GLU:HG3	1:B:336:LYS:CG	2.50	0.42
1:B:234:LEU:CD2	1:B:234:LEU:N	2.83	0.42
1:A:6:LEU:HD12	1:A:60:ILE:HG23	2.02	0.41
1:B:194:GLN:HG3	1:B:234:LEU:HA	2.01	0.41
1:B:100:LEU:HD11	1:B:149:VAL:HG22	2.02	0.41
1:B:176:LEU:HD23	1:B:176:LEU:O	2.21	0.41
1:B:236:ASN:HA	1:B:237:PRO:HA	1.90	0.41
1:A:116:GLU:HG2	1:A:120:MET:HE3	2.03	0.41
1:B:341:VAL:CG2	1:B:341:VAL:O	2.64	0.41
1:A:300:VAL:HG21	1:A:331:PHE:HB3	2.03	0.40
1:B:237:PRO:HG2	1:B:311:ILE:HD11	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/362 (99%)	352 (98%)	6 (2%)	2 (1%)	25	13
1	B	359/362 (99%)	347 (97%)	11 (3%)	1 (0%)	41	32
All	All	719/724 (99%)	699 (97%)	17 (2%)	3 (0%)	34	24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	317	ASN
1	A	317	ASN
1	A	14	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/315 (100%)	301 (96%)	14 (4%)	28	13
1	B	314/315 (100%)	302 (96%)	12 (4%)	33	18
All	All	629/630 (100%)	603 (96%)	26 (4%)	30	15

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ILE
1	A	6	LEU
1	A	64	GLU
1	A	125	LYS
1	A	149	VAL
1	A	165	ASP
1	A	208	ARG
1	A	213	LEU
1	A	248	ASP
1	A	258	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	327	GLU
1	A	330	HIS
1	A	346	LEU
1	A	355	LYS
1	B	5	LYS
1	B	15	ASN
1	B	94	LYS
1	B	149	VAL
1	B	151	GLU
1	B	194	GLN
1	B	234	LEU
1	B	256	LYS
1	B	258	LYS
1	B	264	GLN
1	B	267	GLU
1	B	320	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	192	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	B	401	-	4,4,4	0.11	0	6,6,6	0.15	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	362/362 (100%)	0.23	14 (3%)	39 47	23, 35, 52, 60	0
1	B	361/362 (99%)	0.16	14 (3%)	39 47	23, 34, 51, 63	0
All	All	723/724 (99%)	0.20	28 (3%)	39 47	23, 35, 51, 63	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	167	LEU	6.7
1	B	170	LEU	5.3
1	A	359	PHE	4.1
1	A	360	SER	4.0
1	B	171	GLU	3.8
1	A	1	MET	3.7
1	A	330	HIS	3.6
1	A	362	LEU	3.5
1	A	179	LEU	3.5
1	A	320	ILE	2.9
1	B	317	ASN	2.9
1	B	308	ARG	2.9
1	B	168	LEU	2.8
1	B	274	ILE	2.8
1	A	15	ASN	2.8
1	B	174	ALA	2.8
1	B	294	GLU	2.4
1	A	168	LEU	2.4
1	A	167	LEU	2.4
1	A	293	GLY	2.4
1	A	16	PHE	2.3
1	B	361	LYS	2.3
1	B	178	ARG	2.2
1	A	103	ARG	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	64	GLU	2.1
1	B	181	LYS	2.1
1	A	301	TYR	2.0
1	B	15	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	401	5/5	0.95	0.13	48,50,51,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.