



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2023 – 01:36 AM JST

PDB ID : 6IUA
Title : Crystal structure of importin-alpha1 bound to the 53BP1 nuclear localization signal (S1678D)
Authors : Matsuura, Y.
Deposited on : 2018-11-27
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	427	
2	B	22	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3911 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Importin subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	3251	2073	549	619	10	0	8	0

- Molecule 2 is a protein called Peptide from TP53-binding protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	22	185	109	41	35	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1678	ASP	SER	engineered mutation	UNP Q12888


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	450	450	450	0	0
3	B	25	25	25	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Importin subunit alpha-1



- Molecule 2: Peptide from TP53-binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.26Å 90.17Å 97.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.32 – 1.70 24.32 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (24.32-1.70) 99.9 (24.32-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.70Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.155 , 0.179 0.154 , 0.178	Depositor DCC
R_{free} test set	3838 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	22.8	Xtrriage
Anisotropy	0.242	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3911	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.65	0/3334	0.75	2/4542 (0.0%)
2	B	0.66	0/188	0.92	0/243
All	All	0.65	0/3522	0.76	2/4785 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	285	ARG	NE-CZ-NH1	-6.24	117.18	120.30
1	A	258	ARG	NE-CZ-NH1	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3251	0	3341	19	0
2	B	185	0	200	1	0
3	A	450	0	0	3	0
3	B	25	0	0	0	0
All	All	3911	0	3541	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:489:LEU:O	1:A:493:GLU:HG3	1.92	0.70
1:A:207:ASP:OD1	1:A:251:GLN:NE2	2.26	0.68
1:A:372:GLN:OE1	3:A:501:HOH:O	2.10	0.68
1:A:482:GLU:O	1:A:486:LYS:HG2	2.05	0.55
1:A:288:MET:HA	1:A:291:LYS:HE3	1.89	0.54
1:A:395:LYS:HG2	1:A:436:ILE:HD12	1.90	0.53
1:A:418[A]:HIS:HD2	3:A:832:HOH:O	1.92	0.52
1:A:262:HIS:O	1:A:268:LEU:HD21	2.10	0.51
1:A:287:GLU:O	1:A:291:LYS:HG3	2.12	0.49
1:A:494:LYS:HE2	1:A:495:TYR:OH	2.13	0.49
1:A:452:GLU:HG2	1:A:457:THR:HG21	1.97	0.46
1:A:462:ILE:O	1:A:466:GLU:HG3	2.15	0.46
1:A:435:LYS:NZ	3:A:508:HOH:O	2.51	0.43
1:A:231:TRP:CE2	2:B:1682[B]:ARG:HD3	2.54	0.43
1:A:272:CYS:HB3	1:A:312:PRO:HB2	2.00	0.42
1:A:425:LEU:HD22	1:A:440:ILE:HG23	2.02	0.42
1:A:477:GLN:O	1:A:485:TYR:HB2	2.20	0.41
1:A:494:LYS:HE2	1:A:495:TYR:CZ	2.55	0.41
1:A:273:TRP:CD2	1:A:312:PRO:HB3	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	428/427 (100%)	424 (99%)	4 (1%)	0	100	100
2	B	21/22 (96%)	21 (100%)	0	0	100	100
All	All	449/449 (100%)	445 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/358 (101%)	360 (99%)	2 (1%)	86	80
2	B	20/19 (105%)	20 (100%)	0	100	100
All	All	382/377 (101%)	380 (100%)	2 (0%)	88	83

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	288	MET
1	A	291	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/427 (98%)	0.36	40 (9%) 8 9	14, 23, 64, 98	0
2	B	22/22 (100%)	1.98	11 (50%) 0 0	21, 35, 102, 112	0
All	All	444/449 (98%)	0.44	51 (11%) 4 5	14, 24, 70, 112	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	485	TYR	9.7
2	B	1674	GLU	7.1
2	B	1673	SER	6.4
2	B	1676	GLU	6.2
1	A	478	ARG	6.1
1	A	455	GLY	6.1
1	A	432	LYS	5.5
2	B	1675	GLU	5.2
1	A	489	LEU	5.1
2	B	1665	SER	4.6
1	A	482	GLU	4.4
2	B	1677	ARG	4.3
2	B	1678	ASP	4.2
1	A	467	CYS	4.1
1	A	108	LYS	4.1
1	A	462	ILE	4.1
1	A	480	GLU	4.1
1	A	486	LYS	4.0
1	A	107	GLU	3.9
1	A	493	GLU	3.8
1	A	88	ASN	3.7
1	A	466	GLU	3.4
2	B	1672	THR	3.3
1	A	475	ALA	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	454	LEU	3.2
2	B	1679	PRO	3.2
1	A	483	SER	3.1
1	A	382	LEU	3.1
1	A	459	LYS	2.9
1	A	317	ILE	2.7
1	A	481	ASN	2.7
2	B	1686	SER	2.7
1	A	492	ILE	2.5
1	A	490	ASN	2.5
1	A	362	ILE	2.5
1	A	458	GLU	2.5
1	A	479	HIS	2.4
1	A	256	LEU	2.4
1	A	491	LEU	2.4
1	A	463	MET	2.4
1	A	457	THR	2.4
2	B	1671	ILE	2.3
1	A	471	ASP	2.3
1	A	124	PHE	2.3
1	A	476	LEU	2.2
1	A	477	GLN	2.2
1	A	465	GLU	2.2
1	A	109	GLN	2.2
1	A	144	LEU	2.2
1	A	474	GLU	2.1
1	A	468	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.